

Literatura

- ABELS H. Über Nachempfindungen im Gebiete des kinaesthetischen und statischen Sinnes. *Z Psych* 1906;43: 268.
- ALPINI D, CAPUTO D, CESARANI A. A paradoxical stabilization effect of visual-somatosensory deprivation on postural control and its possible relationship to cerebellar dysfunction. In *Proceedings of the Annual N.E.S. Conference*, Alghero, Sardinia, 2001.
- ALPINI D, CAPUTO D, HAHN A. Grading brainstem involvement in multiple sclerosis – By means of electro-oculography. *J Neuro/Virol* 2000; (Suppl 1).
- ALPINI D, CESARANI A, HAHN A. Tinnitus school: an educational approach to tinnitus management based on a stress-reaction tinnitus model. *Int Tinnitus J* 2007; 13(1): 63–68.
- ALPINI D, HAHN A, RIVA D. Static and dynamic postural control adaptations induced by playing ice hockey. *Sport Sci Health* 2008; 2(3): 85–92.
- BALDWIN D, KING TT, CHEVRETTON E, et al. Bilateral cerebellopontine angle tumors in neurofibromatosis type 2. *J Neurosurg* 1991; 74: 910–995.
- BAREŠ K, HAHN A, NAVARA M, et al. BERA findings in Deaf Patients Symposium für Schädelbasis – Chirurgie. Berlin, Charite, 1989. (prosím, rozlišit, co je název článku, co je název symposia, doplnit stránky)
- BATTAGLIA A, BURCHETTE R, CUEVA R. Combination therapy (intratympanic dexamethasone + high-dose prednisone taper) for the treatment of idiopathic sudden sensorineural hearing loss. *Otol Neurotol* 2008; 29(4): 453–460.
- BETKA J, ZVĚŘINA E, et al. Minimálně invazivní endoskopická a endoskopicky asistovaná chirurgie vestibulárního schwannomu. *Endoskopie* 2009; 18(2): 67–71.
- BLAKLEY BW. Update on intratympanic gentamicin for Ménière's disease. *Laryngoscope* 2000; 110: 236–240.
- BOCKENHEIMER S, SCHMIDT CL, ZOLLNER, C. Neuro-otological findings in patients with small acoustic neuromas. *Arch Otorhinolaryngol* 1984; 239: 31–39.
- BRANDT T, STRUPP M, NOVOZHILOV S, et al. Artificial neural network posturography detects the transition of vestibular neuritis to phobic postural vertigo. *J Neurol* 2012; 259: 182–184.
- BRYNIN R, YOMTOB C. Missed cervical spine fracture - chiropractic implications. *J Manipulative Physiol Ther* 1999; 22(9): 610–614.
- CLARK DB, HIRSCH BE, SMITH MG, et al. Panic in otolaryngology patients presenting with dizziness or hearing loss. *Am J Psychiat* 1994; 151: 1223–1225.
- CLAUSSEN CF. *Über die Aufzeichnung und Auswertung ausgewählter quantitativer Gleichgewichtsprüfungen* (Habil. Dis.). Schrift Berlin: Dissertations-druckstelle 1970.
- CLAUSSEN CF, AUST G, SCHÄFER W. *Atlas der Elektronystagografie*. Hamburg: Werner Rudat 1986.
- DEHLER R, DEHLER F, CLAUSSEN CF. Competitive kinesthetic interaction therapy. *Int Tinnitus J* 2000; 6(1): 29–35.
- EVANA RW. Some observations on whiplash injuries. *Neurol Clin* 1992, 10(4): 975–997.
- FITZGERALD DC. Head trauma: hearing loss and dizziness. *J Trauma* 1996; 40(3): 488–496.

- FOLMER RL, GRIEST SE. Chronic tinnitus resulting from head or neck injuries. *Laryngoscope* 2003; 113(5): 821–827.
- FORT GG. Ramsay Hunt syndrome. In FERRI FF. *Ferri's Clinical Advisor* 2011. Philadelphia: Mosby Elsevier 2010.
- GAGEY PM, TOUPET M. *L'amplitude des oscillations posturales dans la bande de fréquence 0.2 Hertz. Etude chez le sujet normal*. Paris, Publications de l'Institut de Posturologie 1998.
- GERSDORFF MC, NOUWEN J, DECAT M, et al. Labyrinthine fistula after holocholesteatomatous chronic otitis media. *Am J Otol* 2000; 21(1): 32–35.
- GIARDO L, ZANNI M, FERNANDEZ M, et al. Plasticity of GABA(a) system during aging: focus on vestibular compensation and possible pharmacological intervention. *Brain Res* 2002; 929(1): 76–86.
- GOEBEL JA, O'MARA W, GIANOLI G. Anatomic considerations in vestibular neuritis. *Otol Neurotol* 2001; 22: 512–518.
- GOLDBERG JM. *The Vestibular System: A Sixth Sense*. Oxford University Press 2012; 10–12.
- HAHN A. Audiovestibular findings in the course of Ménière's disease 1999 – Update. In STERKERS O, et al. *Ménière's Disease Update*. Hague: Kugler Publications 2000; 285–289.
- HAHN A. Diagnostická a terapeutická strategie v léčbě tinnitu. *Čes Ger Rev* 2005; 3(1): 24–26.
- HAHN A. Ginkgo biloba: local experiences. *Int Tinnitus J* 2000; 6(1): 54–55.
- HAHN A. *Otoneurologie*. Praha: Avicenum 2004.
- HAHN A. *Základní vyšetření v otoneurologii a diagnostika nádoru statoakustického nervu*. Praha: Vesmír 2000.
- HAHN A, NOVOTNÝ M, SHOTEKOV PM, et al. Comparison of innarizine/dimethylhydrinate fixed combination with the respective monotherapies for vertigo of various origins: a randomized, double-blind, active-controlled, multicentre study. *Clin Drug Investig* 2011; 31(6): 371–383.
- HAHN A, POLNICKÝ R, ŠTOLBOVÁ K, OTRUBA L. The Monitoring of the vestibular disorders in the population. In: CLAUSSEN CF, SAKATA E, ITOH A. *Vertigo, Nausea, Tinnitus and Hearing Loss in Central and Peripheral Vestibular Disease*. Amsterdam: Elsevier 1995; 209–212.
- HAHN A, PROCHÁZKOVÁ I, ŠEJNA J. Skutil Ginkgo biloba Extract EGb 761[®] and pentoxifylline in chronic tinnitus: a randomised, double-blind comparative study. In 86. *Jahrestagung der Deutschen Gesellschaft für Hals-Nasen-Ohren-Heilkunde, Kopf- und Hals-Chirurgie, Berlin, 13.–16. Mai 2015*. Stuttgart: Thieme Verlag 2015.
- HAHN A, RADKOVA L, ACHIEMERE G, et al. Multimodal therapy for chronic tinnitus. *Int Tinnitus J* 2008; 14(1): 69–75.
- HAHN A, RADKOVÁ L, SCHALEK P, SKUTIL J. The intra-ear catheter in the chronic tinnitus treatment. *Otorinolaringologia* 2007; 57(3): 109–111.
- HAHN A, ŠEJNA I, ŠTEFFLOVÁ B, et al. A fixed combination of cinnarizine/dimethylhydrinate for the treatment of patients with acute vertigo due to vestibular disorders: a randomized, referencecontrolled clinical study. *Clin Drug Investig* 2008; 28(2): 89–99.

- HAHN A, SCHALEK P, ŠEJNA I, ROSINA J. Combined tinnitus therapy with laser and EGb 761: further experiences. *Int Tinnitus J* 2012; 17(1): 47–50.
- HAHN A, SCHNEIDER D, HAID CT, et al. *Neurootological considerations for diagnosis of acoustic neurinomas*. Prag: Barány Society, 1992.
- HAHN A, SMOLÍK P. Psychiatric disturbances and psychotropic drugs in tinnitus patients. *Int Tinn J* 2002; 8(1): 45–49.
- HAHN A, ŠEJNA I, ŠTOLBOVÁ K, ČOČEK A. Combined laser – EGb 761 tinnitus therapy. *Acta Otolaryngol* 2001; 121(Suppl. 545): 92–93.
- HAHN A, ŠEJNA I, ŠTOLBOVÁ K, ČOČEK A. Visuo-vestibular biofeedback in patients with peripheral vestibular disorders. *Acta Otolaryngol* 2001; 121(Suppl. 545): 88–91.
- HAHN A, ŠTOLBOVÁ K. Ginkgo biloba: local experiences. *Int Tinnitus J* 2000; 6 (1): 54–55.
- HAIN TC, HELMINSKI JO, REIS IL, et al. Vibration does not improve results of the canalolith repositioning procedure. *Arch Otolaryngol Head Neck Surg* 2000; 126(5): 617–622.
- HAKUBA N, HATO N, ZSINOMORI Y, et al. Labyrinthine fistula as a late complication of middle ear surgery using the canal wall down technique. *Otol Neurotol* 2002; 23: 832–835.
- HAN DH, KIM DG, CHI JG, et al. Malignant triton tumor of the acoustic nerve. Case report. *J Neurosurg* 1992; 76: 874–877.
- HELMS J. Hörerhalt bei der Chirurgie von Akustikusneurinomen. *HNO-Informationen* 1992; 2: 5–8.
- HENRY JA, JASTREBOFF MM, JASTREBOFF PJ, et al. Guide to conducting tinnitus retraining initial and follow-up interviews. *J Rehab Res Dev* 2003; 40(2): 157–177.
- HOFFER ME, et al. Characterizing and treating dizziness after mild head trauma. *Otol Neurotol* 2004; 25: 135–138.
- HOLÝ R, NAVARA M, DOŠEL P, et al. Hyperbaric oxygen therapy in idiopathic sudden sensorineural hearing loss (ISSNHL) in association with combined treatment. *Undersea Hyperbaric M* 2011; 38: 135–140.
- HOSUK C, WON-HO CH. Perilymph fistula test. *N Engl J Med* 2012; 366: 48.
- JACOB RG, LILIENFELD SO, FURMAN JMR, et al. Panic disorder with vestibular dysfunction: further clinical observations and description of space and motion phobic stimuli. *J Anx Dis* 1989; 117(3): 130.
- JASTREBOFF PJ, JASTREBOFF MM. Tinnitus retraining therapy (TRT) as a method for treatment of tinnitus and hyperacusis patients. *J Am Acad Audiol* 2000; 11: 162–177.
- JIAN BJ, SHINTANI T, EMANUEL BA, et al. Convergence of limb, visceral, and vertical semicircular canal or otolith inputs onto vestibular nucleus neurons. *Exp Brain Res* 2002; 144(2): 247–257.
- KAPTEYN TS, DEWIT G. Posturography as an auxiliary in vestibular investigation. *Acta Otolaryngol* 1972; 73: 104–111.
- KETTNER NW, GUEBERT GM. The radiology of cervical spine injury. *J Manipulative Physiol Ther* 1991; 14(9): 518–526.
- KEVETTER GA, LEONARD RB. Molecular probes of the vestibular nerve. II. Characterization of neurons in Scarpa's ganglion to determine separate populations within the nerve. *Brain Res* 2002; 928(1–2): 18–29.

- KILPATRICK JK, SISMANIS A, SPENCER FR, et al. Low-dose oral methotrexate management of patients with bilateral Ménière's disease. *Ear Nose Throat J* 2000; 79(2): 82–83.
- KISELY M, TOTTH A, EMRI M, et al. Effect of pathologic and induced peripheral vestibular balance disturbance on the central nervous system. *Orv Hetil* 2002; 26(143) (21, Suppl 3): 1330–1332.
- KJAER D, PEDERSEN CB. Ménière's disease. The clinical effect of gentamicin treatment. *Ugeskr Laeger* 2000; 162(19): 2739–2743.
- KOHEN-RAZ, R. *Learning Disabilities and Postural Control*. London: Freund Publishers 1986.
- KONDO K, SAGARA H, HIROSAWA K, et al. Hair cell development in vivo and in vitro: analysis by using a monoclonal antibody specific to hair cells in the Chininner ear. *J Comp Neurol* 2002; 445(2): 176–198.
- KUHWEIDE R, VAN DE STEENE V, VLAMINCK S, CASSELMAN JW. Ramsay Hunt syndrome: pathophysiology of cochleovestibular symptoms. *J Laryngol Otol* 2002; 116(10): 844–848.
- KUSUMA S, LIOU S, HAYNES DS. Disequilibrium after cochlear implantation caused by a perilymph fistula. *Laryngoscope* 2005; 115: 2526.
- LANGE G, KELLER R. Bilateral malfunction of peripheral vestibular organs. Observations of 20 cases of Dandy syndrome. *Laryngorhinootologie* 2000; 79(2): 77–80.
- LAUGHLIN PJ, REDFERN MS. Spectral analysis of visually induced postural sway in healthy elderly young subjects. *IEEE Transact Rehabil Engineering* 2001; 9(1): 24–30.
- LEVINE SC, ANTONELLI PJ, et al. Relative value of diagnostic tests for small acoustic neuromas. *Am J Otol* 1991; 12(5): 341–346.
- LI JC, LI CJ, EPLEY J, et al. Cost-effective management of benign positional vertigo using canalith repositioning. *Otolaryngol Head Neck Surg* 2000; 122(3): 334–339.
- LONGRIDGE NS, MALLINSON AI. Low-dose intratympanic gentamicin treatment for dizziness in Ménière's disease. *J Otolaryngol* 2000; 29(1): 35–39.
- MAMIKOGLU B, WIET RJ, ESQUIVEL CR. Translabyrinthine approach for the management of large and giant vestibular schwannomas. *Otol Neurotol* 2002; 23(2): 224–227.
- MARKS S, ARENBERG IK, HOFFER ME. Round-Window Microcatheter administered microdose of gentamicin: An alternative in treatment of tinnitus associated with Ménière's disease. *Laryng Rhin Otol* 2000; 79(6): 327–331.
- MICHEL O, JAHNS T, JOOST-ENNEKING M, et al. The Stennert antiphlogistic-rheologic infusion schema in treatment of cochleovestibular disorders. *HNO* 2000; 48(3): 182–188.
- MORAD Y, AZARIA B, AVNI I, et al. Posturography as an indicator of fatigue due to sleep deprivation. *Aviat Space Environ Med* 2007; 78(9): 859–863.
- ÖDKVIST LM, ARLINGER S, BILLERMARK E, et al. Effects of middle ear pressure changes on clinical symptoms in patients with Ménière's disease: A clinical, multi-center, placebo-controlled study. *Acta Otolaryngol Suppl* 2000; 543: 99–101.
- OKUBO J, WANATABE I. Influence of foot position and visual field condition in examination for equilibrium function and sway of the center of gravity in normal persons. *Agressologie* 1979; 20: 127–132.

- OZUNLU A, MUS N, GULHAN M. Multiple sclerosis: a cause of sudden hearing loss. *Audiology* 1998; 37(1): 52–58.
- PANJABI MM, NIBU K, CHOLEWICKI J. Whiplash injuries and the potential for mechanical instability. *Eur Spine J* 1998; 7(6): 484–492.
- PLONTKE SK, LÖWENHEIM H, MERTENS J, et al. Randomized, double blind, placebo controlled trial on the safety and efficacy of continuous intra tympanic dexamethasone delivered via a round window catheter for severe to profound sudden idiopathic sensorineural hearing loss after failure of systemic therapy. *Laryngoscope* 2009; 119(2): 359–369.
- PROCHÁZKA M, HAHN A. Komplexní laserová rehabilitační terapie tinnitu: dlouhodobá dvojitá slepá studie na skupině 200 pacientů v průběhu 3 let. *Laser Partner* 2002; 51.
- ROBERTS LE. Ringing ears: the neuroscience of tinnitus. *J Neurosci* 2010; 30(45): 14972–14979.
- ROZSASI A, SIGG O, KECK T. Persistent inner ear injury after diving. *Otol Neurotol* 2003; 24: 195–200.
- SAKATA E, KITAGO Y, MURATA Y, TERAMOTO K. Behandlung der Ménièreschen Krankheit. Paukenhöhlen-infusion von Lidocain-und Steroidlösung. *Aurix Nasus Larynx (Tokyo)* 1986; 13(2): 79–89.
- SALLEY LH JR, GRIMM M, SISMANIS A, et al. Methotrexate in the management of immune mediated cochleovesibular disorders: clinical experience with 53 patients. *J Rheumatol* 2001; 28(5): 1037–1040.
- SHULMAN A. A final common pathway for tinnitus the medical temporal lobe system. *Int Tinnitus J* 1995; 1(2): 115–126.
- SHULMAN A, GOLDSTEIN B, STRASHUN AM. Final common pathway for tinnitus: theoretical and clinical implications of neuroanatomical substrates. *Int Tinnitus J* 2009; 15(1): 5–50.
- SCHOPS P, PFINGSTEN M, SIEBERT U. Reliability of manual medical examination techniques of the cervical spine. Study of quality assurance in manual diagnosis. *Orthop Ihre Grenzgeb* 2000; 138(1): 2–7.
- SCHUKNECHT HF. *Pathology of the Ear*. Cambridge, MA: Harvard University Press, 1974.
- SINGER RN. Interrelationship of physical, perceptual-motor and academic variables in elementary school children. *Perc Motor Skills* 1968; 27: 1323–1332.
- STERKERS JM, PERRE J, VIATA P, et al. The origin of acoustic neuromas. *Acta Otolaryngol (Stockh.)* 1987; 103: 427.
- STRUPP M, HUPERT D, FRENZEL C, et al. Long-term prophylactic treatment of attacks of vertigo in Ménière's disease comparison of a high with a low dosage of betahistine in an open trial. *Acta Otolaryngol* 2008; 128(5): 520–524.
- STRUTZ J. Value of the Stennert schema in therapy of sudden deafness. *HNO* 2000; 48(3): 165–167.
- TASS PA, ADAMCHIC H-J, FREUND T, et al. Counteracting tinnitus by acoustic coordinated reset neuromodulation. *Restor Neurol Neurosci* 2012; 30: 137–159.
- WEI BP, MUBIRU S, O'LEARY S. Steroids for idiopathic sudden sensorineural hearing loss. *Cochrane Database Syst Rev* 2006; 1: CD003998.

- WEI BP, MUBIRU S, O'LEARY S. Steroids for idiopathic sudden sensorineural hearing loss. *Cochrane Database Syst Rev* 2013; 7: CD003998.
- WELLING DB, NAGARAJA HN. Endolymphatic mastoid shunt: a reevaluation of efficacy. *Otolaryngol Head Neck Surg* 2000; 122(3): 340–345.
- WORDLICZEK J, SZCZEPANIK AM, BANACH M, et al. The effect of pentoxifylline on post-injury hyperalgesia in rats and postoperative pain in patients. *Life Sci* 2000; 66(12): 1155–1164.