

Literatura

- Arntz, A., & Jacob, G. (2012). *Schema Therapy in practice: An introductory guide to the schema mode approach*. Oxford: Wiley Blackwell.
- Bamelis, L. L., Evers, S. M., Spinhoven, P., & Arntz, A. (2013). Results of a multicenter randomized controlled trial of the clinical effectiveness of Schema Therapy for personality disorders. *American Journal of Psychiatry*, 170(12), 1503–1508. doi: 10.1176/appi.ajp.2013.12040518
- Bernstein, D. P., Arntz, A., & de Vos, M. (2007). Schema focused therapy in forensic settings: Theoretical model and recommendations for best clinical practice. *International Journal of Forensic Mental Health*, 6, 169–183.
- Cockram, M. D., Drummond, P. D., & Lee, W. C. (2010). Role and treatment of early maladaptive schemas in Vietnam veterans with PTSD. *Clinical Psychology and Psychotherapy*, 17, 165–182.
- Comtois, K. A., Russo, J., Snowden, M., Srebnik, D., Ries, R., & Roy-Byrne, P. (2003). Factors associated with high use of public mental health services by persons with borderline personality disorder. *Psychiatric Services*, 54(8), 1149–1154.
- Farrell, J., & Shaw, I. (2012). *Group Schema Therapy for borderline personality disorder: A step-by-step treatment manual with patient workbook*. Oxford: Wiley Blackwell.
- Farrell, J., Shaw, I., & Webber, M. (2009). A schema-focused approach to group psychotherapy for outpatients with borderline personality disorder: A randomized controlled trial. *Journal of Behavior Therapy and Experimental Psychiatry*, 40, 317–328. doi:10.1016/j.jbtep.2009.01.002
- Farrell, J., Young, J., & Shaw, I. (2011). Online manuscript. www.bpd-home-base.org/
- Giesen-Bloo, J., van Dyck, R., Spinhoven, P., van Tilburg, W., Dirksen, C., van Asselt, T., et al. (2006). Outpatient psychotherapy for borderline personality disorder: Randomized trial of schema-focused therapy vs. transference-focused psychotherapy. *Archives of General Psychiatry*, 63, 649–658.

Group Schema Therapy Mixed Personality Disorder Presentation ISST Conference N.Y. 2012.

- Jacob, G. A., & Arntz, A. (2013). Schema Therapy for personality disorders – a review. *International Journal of Cognitive Therapy*, 6(2), 172–185.
- Linehan, M. M. (1993). Cognitive-behavioral treatment of borderline personality disorder. New York: The Guilford Press.
- Lobbestael, J., van Vreeswijk, M., Spinhoven, P., Schouten, E., & Arntz, A. (2010). Reliability and validity of the short Schema Mode Inventory (SMI). *Behavioural and Cognitive Psychotherapy*, 38, 437–458.
- Muste, E., Weertman, A., & Claassen, A. M. (2009). Handboek Klinische Schematherapie. Houten: Bohn Stafleu van Loghum.
- Nadort, M., Arntz, A., Smit, J. H., Wensing, M., Giesen-Bloo, J., Eikelenboom, M., et al. (2009). Implementation of outpatient schema therapy for borderline personality disorder with versus without crisis support by the therapist outside office hours: A randomized trial. *Behaviour Research and Therapy*, 47(11), 961–973.
- Reiss, N., & Vogel, F. (2010). Stationäre Schematherapie bei Borderline-Persönlichkeitsstörung. In E. Roediger & G. Jacob (Eds.), *Fortschritte der Schematherapie. Konzepte und Anwendungen*, pp. 217–226. Göttingen: Hogrefe.
- Reiss, N., Dominiak, P., Harris, D., Knornschild, C., Schouten, E., & Jacob, G. A. (2012). Reliability and validity of the German version of the revised Schema Mode Inventory (SMI). *European Journal of Psychological Assessment*, 4, 297–304. doi: 10.1027/1015-5759/a000110
- Reiss, N., Lieb, K., Arntz, A., Shaw, I. A., & Farrell, J. M. (2013a). Responding to the treatment challenge of patients with severe BPD: Results of three pilot studies of inpatient schema therapy. *Behavioural and Cognitive Psychotherapy*, 1–13. doi:10.1017/S1352465813000027
- Reiss, N., Vogel, F., Nill, M., Graf-Morgenstern, M., Finkelmeier, B., & Lieb, K. (2013b). Behandlungszufriedenheit von Patientinnen mit Borderline Persönlichkeitsstörung bei stationärer Schematherapie. *Psychotherapie – Psychosomatik – Medizinische Psychologie*, 3, 93–100.
- Spinhoven, J., Giesen-Bloo, J., van Dyck, R., Kooiman, K., & Arntz, A. (2007). The therapeutic alliance in schema-focused therapy and transference-focused psychotherapy for borderline personality disorder. *Journal of Consulting and Clinical Psychology*, 75(1), 104–115.

- Van Asselt, A. D., Dirksen, C. D., Arntz, A., Giesen-Bloo, J. H., van Dyck, R., Spinhoven, P., et al. (2008). Outpatient psychotherapy for borderline personality disorder: Cost-effectiveness of schema-focused therapy vs. transference-focused psychotherapy. *The British Journal of Psychiatry*, 192(6), 450–457. doi:10.1192/bjp. bp.106.033597
- Yalom, I. D., & Leszcz, M. (2005). *The theory & practice of group psychotherapy* (5th ed.). New York: Basic Books.
- Young, J. E. (1990). *Cognitive therapy for personality disorders: A schema-focused approach*. Sarasota, FL: Professional Resource.
- Young, J. E., Klosko, J. S., & Weishaar, M. E. (2003). *Schema Therapy: A practitioner's guide*. New York: The Guilford Press.
- Zarbock, G., Rahn, V., Farrell, J., & Shaw, I. (2011). *Group Schema Therapy: An innovative approach to treating patients with personality disorder, developed by Farrell & Shaw*. DVD set. IVAH: Hamburg. www.bpd-home-base.org