

Literatura

- Adachi YU, Watanabe K, Higuchi H, et al. Flumazenil reduces the hypnotic dose of propofol in male patients under spinal anesthesia. *J Anesth.* 2002;16:9–12.
- Adams S, Werner C. Vom Razemat zum Enantiomer: (S)-Ketamin; Renaissance einer Substanz? *Anaesthesist.* 1997;46:1026–1042.
- Amrein R, Hetzel W. Pharmacology of dormicum (midazolam) and anexate (flumazenil). *Acta Anaesth Scand.* 1990;34(Suppl. 92):6–15.
- Amrein R, Hetzel W. Pharmacology of drugs frequently used in ICUs: Midazolam and flumazenil. *Intens Care Med.* 1991;17(Suppl. 1):1–10.
- Anttila M, Penttilä J, Helminen A, et al. Bioavailability of dexmedetomidine after extravascular doses in healthy subjects. *J Clin Pharmacol.* 2003;56:691–693.
- Bailey PL, Port JD, Pace NL, et al. The ED₅₀ of carfentanil for elk immobilization with and without the tranquilizer R51703. *J Wildl Manager.* 1985;49:931–934.
- Bartošová L. Otravy nervově paralytickými látkami: co o nich víme? *Českosl Fyziol.* 2006;55:66–75.
- Behne, M. The effect of flumazenil on fentanyl – induced respiratory depression. *Anaesthesist.* 1991;40:271–275.
- Belleville JP, Ward DS, Bloor BC, et al. Ventilatory effects of dexmedetomidine in humans. *Anesthesiology.* 1990;73:A1167.
- Benumof JL, Dagg R, Benumof R. Critical hemoglobin desaturation will occur before return to an unparalyzed state following 1 mg/kg intravenous succinylcholine. *Anesthesiology.* 1997;87:979–982.
- Bermejo S, et al. Absence of neuromuscular blockade after administration of intratracheal succinylcholine. Abstracts and Programme Euroanaesthesia 2004. *Eur J Anesthesiology.* 2004:139.
- Birch BR, Miller RA. An assessment of resedation following flumazenil – induced antagonism intravenous midazolam: comparison of psychomotor and amnesic recovery with a non-sedated reference group. *J Psychopharmacol.* 1995;9:103–111.
- Bloor BC, Fonarow G C. Intracoronary dexmedetomidine does not produce coronary vasoconstriction at therapeutic blood levels in chronically instrumented anaesthetized swine. *J Cardiothor Vasc Anesthesia.* 1994;8(Suppl. 3):4.
- Blouin RT, Conard PF, Perreault S, et al. The effect of flumazenil on midazolam – induced depression of the ventilatory response to hypoxia during isohypercarbia. *Anesthesiology.* 1993;78: 635–641.
- Bourke DL, Malit LA, Smith TC. Respiratory interactions of ketamine and morphine. *Anesthesiology.* 1987;66:153–156.
- Braestrup C, Albrechtsen R, Squires RF. High densities of benzodiazepine receptors in human cortical areas. *Nature.* 1977;269:702–704.
- Brenchley J, Ramlakhan S. Intranasal alfentanil for acute pain in children. *Emerg Med J.* 2006;23:488.
- Brodie B. Experiments and observations on the different modes in which death is produced by certain vegetable poison. London: W. Bulmer and Co., 1811.

- Buck M. Options for intranasal procedural sedation and analgesia in children. *Pediatr Pharm.* 2013;19:1–7.
- Capuano SV, Lerche NW, Valverde CR. Cardiovascular, respiratory, thermoregulatory, sedative, and analgesic effects of intravenous administration of medetomidine in rhesus macaques. *Lab Nim Sci.* 1999;49:537–544.
- Cistola AM, Golder FJ, Centonze LA, et al. Anesthetic and physiologic effects of tiletamine, zolazepam, ketamine, and xylazine combination (TKX) in feral cats undergoing surgical sterilization. *J Feline Med Surg.* 2004;6:297–303.
- Clanahan AS, Marschall RJ. Diazepam potentiates the coronary vasodilator actions of adenosine in anaesthetized dogs. *Brit J Pharmacol.* 1980;70:68–678.
- Claude Bernard. *Lecons sur les effets des substances toxiques et medicamenteuses.* Paris: J.-B.Bailliere et Fils, 1857:311–325.
- Corssen G, Domino EF. Dissociative anesthesia: Further pharmacologic studies and first clinical experience with the phencyclidine derivative Cl-581. *Anesth Analg.* 1996;45(29).
- Crozier TA, Beck D, Schlaeger M, et al. Endocrinological changes following etomidate, midazolam, or methohexitol for minor surgery. *Anesthesiology.* 1987;66:628–635.
- Dagher AM, Bradley EL Jr, Kissin I. Midazolam antagonizes the analgesic effects of morphine in rats. *Anesth Analg.* 1987;66:944–947.
- Dahaba AA, Bornemann H, Rehak PH, et al. Effect of flumazenil on bispectral index monitoring in unpremedicated patients. *Anesthesiology.* 2009;110:1036–1040.
- Dershawitz M, Walsh JL, Morishige RJ, et al. Pharmacokinetics and pharmacodynamics of inhaled versus intravenous morphine in healthy volunteers. *Anesthesiology.* 2000;93:619–628.
- Diaz JH. Intranasal ketamine preinduction of paediatric outpatients. *Pediatr Anaesth.* 1997;7:273–278.
- Doherty TJ, McDonell WN, Dyson DH, et al. Cardiorespiratory effects of a 5HT2 antagonist (R51703) in awake and anesthetized dogs. *Can J Vet Res.* 1996;60:172–178.
- Donati F. The right dose of succinylcholine. *Anesthesiology.* 2003;99:1037–1078.
- Duncalf D, Jeyaram C, Tarter R, et al. Ocular absorption of naloxone in narcotic addicts. *Int J Addict.* 1975;10:23–27.
- Dundee JW, Halliday NJr, Harper KW, et al. Midazolam: a review of its pharmacological properties and therapeutic use. *Drugs.* 1984;28:519–543.
- Dunton AW, Schwam E, Pitman V, et al. The relationship between dose and duration of action of intravenous flumazenil in reversing sedation induced by a continuous infusion of midazolam. *Eur J Anaesthesiol.* 1988;2:97–102.
- Dyck JB, Maze M, Haack C, et al. The pharmacokinetics and hemodynamic effects of intravenous and intramuscular dexmedetomidine hydrochloride in adult human volunteers. *Anesthesiology.* 1993b;78:813–820.
- Eddy DR, Storm WF, Gibbons JA, et al. Reversal of Zolpidem intoxication by sublingual flumazenil. <http://www.stormingmedia.us/39/3921/A392105.html>.
- Enserink M, Stone R. Toxicology: Questions swirl over knockout gas used in hostage crisis. *Science.* 2002;298:1150–1151.
- Erhardt W, Henke J, Haberstroh J. Anästhesie und Analgesie beim Klein – und Heimtier sowie bei Vögeln, Reptilien, Amphibien und Fischen. New York, Stuttgart: Schattauer, 2004.

- Faclam M, Schoch P, Haefely W. Relationship between benzodiazepine receptor occupancy and potentiation of GABA-stimulated chloride flux in vitro of four ligands of differing intrinsic efficacies. *J Pharmacol Exp Ther.* 1992;261:1106–1112.
- Farnsworth ST, Gauthier ME, McJames SW, et al. Ocular transmucosal absorption and toxicity of sufentanil in dogs. *Anesth Analg.* 1998;86:138–140.
- Farr C, Menzel J, Seeberger J, et al. Clinical pharmacology and therapeutic use of hyaluronidase with special consideration to hylase „Dessau“. *Transplantion Wiener Medizinische Wochenschrift* 1997;147:347–355.
- Flacke JW, Bloor BC, Flacke WE, et al. Reduced narcotic requirement by clonidine with improved hemodynamic and adrenergic stability in patients undergoing coronary bypass surgery. *Anesthesiology.* 1987;11–19.
- Flecknell PA. Laboratory animal anaesthesia. 2. Auflage. London, San Diego, New York, Boston, Sydney, Tokyo, Toronto: Academic Press, 1966.
- Furst SR, Weinger MB. Dexmedetomidine, a selective alpha₂-adrenergic agonist, does not potentiate the cardiorespiratory depression of alfentanil in the rat. *Anesthesiology.* 1990;72:882–888.
- Gath I, Weidenfeld J, Collins G, et al. Electrophysiological aspects of benzodiazepine antagonists, Ro15-1788 a Ro153505. *Br J Clin Pharm.* 1984;18:541–547.
- Geller E, Weinbroum A, Schiff B, et al. The effects of flumazenil on the process of recovery from halothane anaesthesia Europe. *J Anaesth.* 1988;1:151–153.
- Ghodse H, Taylor DR, Greaves JL, et al. The opiate addiction test: a clinical evaluation of a quick test for physical dependence on opiate drugs. *Br J Clin Pharmacol.* 1995;39:257–259.
- Girdler NM, Fairbrother KJ, Lyne JP, et al. A randomised crossover trial of post-operative cognitive and psychomotor recovery from benzodiazepine sedation: effects of reversal with flumazenil over a prolonged recovery period. *Brit Dental J.* 2002;192:335–339.
- Girdler NM, Lyne JP, Wallace R, et al. A randomised, controlled trial of cognitive and psychomotor recovery from midazolam sedation following reversal with oral flumazenil. *Anaesthesia.* 2002;57:868–876.
- Glisson SN, Haddad W, Kubak MA, et al. Anti-adrenergic activity of diazepam, midazolam and lorazepam in dogs. *Anesthesiology – ASA abstracts.* 1983;59:A328.
- Haefely W. Ein Benzodiazepin, das die Wirkung von Benzodiazepinen aufhebt. *Magazin Roche.* 1982;14:10–17.
- Havdala HS, Borison LR, Diamond BI. Ketamine anesthesia and analgesia: neurochemical differentiation. *Anesthesiology.* 1980;55:1–11.
- Hayashi Y, Sumikawa K, Maze M, et al. Dexmedetomidine prevents epinephrine-induced arrhythmias through stimulation of central alpha₂-adrenoceptors in halothane-anesthetized dogs. *Anesthesiology.* 1991;75:113–117.
- Hayes AH, Breslin DS, Mirakhur RK, et al. Frequency of hemoglobin desaturation with the use of succinylcholine during rapid sequence induction of anaesthesia. *Acta Anaesthesiol Scand.* 2001;45:746–749.
- Heier T, Feiner JR, Lin J, et al. Hemoglobin desaturation after succinylcholine-induced apnea: A study of the recovery of spontaneous ventilation in healthy volunteers. *Anesthesiology.* 2001;94:754–759.

- Heinrichs M, Baumgartner T, Kirschbaum C, et al. Social support and oxytocin interact to suppress cortisol and subjective responses to psychosocial stress. *Biol Psychiatry*. 2003;54:1389–1398.
- Hellebrekers LJ, De Boer EJW, Van Zuylen MA, et al. A comparison between medetomidine-ketamine and medetomidine-propofol-anaesthesia in rabbits. *Lab Anim*. 1997;31:58–69.
- Henderson JM, Brodsky DA, Fischer DM, et al. Pre-induction of anesthesia in pediatric patients with nasally administered sufentanil. *Anesthesiology*. 1988;68:671–675.
- Heniff MS, Moore GP, Trout A, et al. Comparison of routes of flumazenil administration to reverse midazolam-induced respiratory depression in a canine model. *Academic Emergency Med*. 1997;4:1115–1118.
- Hering W, Schüttler J. Intravenöse Anesthetika. In: Hempelmann G, Krier C, Schulte AM, et al. *Anästhesiologie Intensivmedizin Notfallmedizin Schmerztherapie*. Stuttgart: Georg Thieme Verlag, New York, 2001:210–227.
- Hess L. Antifibrilační účinek medetomidinu v časné fázi místní ischémie srdeční u psa. *Cor Vasa*. 1995;37:295–298.
- Hess L. Nalmefen. *Remedia*. 2013;23:298–301.
- Hess L. Nové trendy v anesteziologii lidoopů. *Veterinářství*. 1999;49:249–241.
- Hess L. The antifibrillatory effect of flumazenil (Anexate) and its combination with flunitrazepam (Rohypnol) in the early stage of local myocardial ischemia in the dog. *Cor Vasa*. 1995;37:34–36.
- Hess L, Málek J, Schreiberová J. Specifický antagonist benzodiazepinů flumazenil na začátku 21. století – je vůbec zapotřebí? *Anesteziol Intenziv Med*. 2006;17: 295–298.
- Hess L, Vodička R, Král J. První zkušenosti s pravotočivým izomerem ketaminu u primátů. *Zoo Praha. Gazella*. 1999;26:53–56.
- Hess L, Votava M, Málek J, et al. Sedative effects of intranasal oxytocin in rabbits and rhesus monkeys. *Physiol Res*. 2016;65(Suppl. 4):473–480.
- Hess L, Vrána M, Vránová Z, et al. A comparison of the antifibrillatory effects of midazolam and flunitrazepam in acute myocardial ischemia in the dog. *Curr Med Res Opinion*. 1987;8:527–530.
- Higgins MJ, Asbury AJ, Brodie MJ. Inhaled nebulised fentanyl for postoperative analgesia. *Anaesthesia*. 1991;46:973–976.
- Hildebrand PJ, Elwood RJ, McClean E, et al. Intramuscular and oral midazolam. Some factors influencing uptake. *Anaesthesia*. 1983;38:1220–1221.
- Hoffman W, Kochs E, Werner C, et al. Dexmedetomidine improves neurologic outcome from incomplete ischemia in the rat. *Anesthesiology*. 1991;75:328–332.
- Holtzman JR Jr, Sloan JW, Jing X, et al. Modification of morphine analgesia and tolerance by flumazenil in male and female rats. *Eur J Pharmacol*. 2003;470:149–56.
- Hommer D, Weingartner H, Breier A. Dissociation of benzodiazepine induced amnesia from sedation by flumazenil pretreatment. *Psychopharmacology*. 1993;112:455–460.
- Hornocker MG, Graighead JJ, Pfeiffer EW. Immobilizing mountain lions with sukcynylcholine chloride and pentobarbital sodium. *J Wildl Mgmt*. 1965;29:880–883.
- Hunkeler W, Möhler H, Pieri L. Selective antagonists of benzodiazepines. *Nature*. 1981;290:514–516.

- Huupponen R, Karhuvaara S, Anttila M, et al. Buccal delivery of an alpha 2-adrenergic receptor antagonist, atipamezole, in humans. *Clin Pharmacol Therapeutics*. 1995;58:506–511.
- Hynie S, Klenerová V. Centrální regulační úloha oxytocinu. *Psychiatrie*. 2008;12(Suppl. 2):4–10.
- Chen G, Ensor CR, Russell D, et al. The pharmacology of 1-(1-phenylcyclohexyl) piperidine HCl. *J Pharm Exp Ther*. 1959;127:241.
- Chiolero RL, Ravussin P, Anderes JP, et al. The effects of midazolam reversal by Ro 15-1788 on cerebral perfusion pressure in patients with severe head injury. *Intens Care Med*. 1988;14:196–200.
- Church J, Lodge D. N-methyl-D-aspartate (NMDA) antagonism in central to the actions of ketamine and other phencyclidine receptor ligand. In: Status of ketamine in anesthesiology. Ann Arbor: E. F. Domino, NPP Books, 1990:501–520.
- Jaakola ML. Dexmedetomidine premedication before intravenous regional anesthesia in minor outpatient hand surgery. *J Clin Anesth*. 1994;6:204–211.
- Jin Y, Wilson S, Elko EE, et al. Ocular hypotensive effects of medetomidine and its analogs. *J Ocul Pharmacol*. 1991;7:285–296.
- Johnstone M, Evans V, Baigel JR. Sernyl (Cl-395) in clinical anesthesia. *Brit J Anesth*. 1959;31:433.
- Kalema-Zikusoka G, Horne WA, Levine J, et al. Comparison of the cardiorespiratory effects of medetomidine-butorphanol-ketamine and medetomidine-butorphanol-midazolam in patas monkeys. *J Zoo Wildl Med*. 2003;34.
- Kalin NH. The neurobiology of fear. *Sci Amer*. 1993;268:94–101.
- Kallio A, Salonen M, Forssell H, et al. Medetomidine premedication in dental surgery – a double-blind cross-over study with a new alpha2-adrenoceptor agonist. *Acta Anaesthesiol*. 1990;Scand 34:171–175.
- Kampen J, et al. Onset time and effectiveness of drugs after intraosseous bolus administration in adults – a model investigation using succinylcholine in clinical setting. Abstracts and Programme Euroanaesthesia 2004. *Eur J Anaesthesiol*. 2004:181–182.
- Karakosta A, Andreoti B, Chapsa C. Flumazenil expedites recovery from sevoflurane/remifentanil anaesthesia when administered healthy unpremedicated patients. *Eur J Anaesthesiol*. 2010;27:955–959.
- Karhuvaara S, Kallio A, Salonen M, et al. Rapid reversal of alpha2-adrenoceptor agonist effects by atipamezole in human volunteers. *Br J Clin Pharmacol*. 1991;31:160–165.
- Karl HW, Keifer AT, Rosenberger JL, et al. Comparison of the safety and efficacy of intranasal midazolam or sufentanil for preinduction of anesthesia in pediatric patients. *Anesthesiology*. 1992;76:209–215.
- Katz N, Pillar G, Peled E, et al. Sublingual flumazenil for the residual effects of hypnotics: zolpidem and brotizolam. *Clin Pharmacol Drug Develop*. 2012:45–51.
- Kearns KS, Swenson B, Ramsay EC. Dosage trials with transmucosal carfentanil citrate in non-human primates. *Zoo Biology*. 1999;18:397–402.
- Kearns KS, Swenson B, Ramsay EC. Oral induction of anesthesia with droperidol and transmucosal carfentanil citrate in chimpanzees (*pan troglodytes*). *J Zoo Wildl Med*. 2000;31:185–189.

- Kelly AM, Kerr D, Dietze P, et al. Randomised trial of intranasal versus intramuscular naloxone in prehospital treatment for suspected opioid overdose. *MJA*. 2005;182:24–27.
- Kivistö KT, Kallio A, Neuvonen PJ. Pharmacokinetics and pharmacodynamics of transdermal dexmedetomidine. *Eur J Clin Pharmacol*. 1994;46:345–349.
- Klotz U, Ziegler G, Reimann, IW. Pharmacokinetics of the selective benzodiazepine antagonist Ro 15-1788 in man. *Eur J Clin Pharmacol*. 1984;27:115–117.
- Kobilka BK, Matsui H, Kobilka TS, et al. Cloning, sequencing, and expression of the gene coding for the human platelet alpha₂-adrenergic receptor. *Science*. 1987;30:650–654.
- Koch SC, Fitzgerald M, Hathway GJ. Midazolam potentiates nociceptive behavior, sensitizes cutaneous reflexes, and is devoid of sedative action in neonatal rats. *Anesthesiology*. 2008;108:122–129.
- Kopman AF, Zhaku B, Lai KS. The „intubating dose of succinylcholine“. The effect of decreasing doses on recovery time. *Anesthesiology*. 2003;99:1050–1054.
- Kreeger TJ, Seal US. Immobilization of gray wolves (*canis lupus*) with sufentanil citrate. *J Wildl Dis*. 1990;26:561–563.
- Kretz FJ, Althaus Ch, Striebel HW, et al. Pulsoxymetrische Untersuchungen und arterielle Blutgasanalysen nach Narkosen mit Midazolam und Fentanyl sowie postoperativer Antagonisierung mit Flumazenil. In: *Anexate (Flumazenil)*. Der erste spezifische Benzodiazepin-Antagonist, Editiones Roche 1989:185–192.
- Kulka PJ, Tryba M, Menzel C, et al. Preoperative clonidine improves postoperative renal function in CABG patients. *Br J Anaesth*. 1993;70(suppl 1):A143.
- Kuntze A. Klinische Beiträge zur Anästhesie und medikamentellen Immobilisation der Zootiere (Ursiden, Feliden, Ruminantier). Berlin und Hamburg: Paul Parey, 1967.
- Kurzová A, Málek J, Hess L, et al. Non-traditional administration of remifentanil in an experimental setting. *Physiol Res*. 2019;68(Suppl. 1):97–103.
- Lancouchová L. Otrava Sernylanem. *Rozhledy Chirurg*. 1979;58:717–720.
- Långsjö J, Kaskinoro K, Kaisti K, et al. Dexmedetomidine and propofol have a different initial site of action: 9AP3-7. *Eur J Anaesthesiol*. 2010;27:148.
- Larsen R. *Anesthesia*. 7., přepracované a rozšířené vydání. Praha: Grada Publishing, 2004.
- Lee EJHD, Williams KM. Chirality: Clinical pharmacokinetic and pharmacodynamic considerations. *Clin Pharmacokinet*. 1990;18:339–345.
- Levänen J, Mäkelä ML, Scheinin H. Dexmedetomidine premedication attenuates ketamine induced cardiostimulatory effects and postanesthetic delirium. *Anesthesiology*. 1995;82:1117–1125.
- Lewandowski AH, Bonar ChJ, Evans SE. Tiletamine-zolazepam, ketamine, and xylazine anesthesia of captive cheetah (*Acinonyx jubatus*). *J Zoo Wildl Med*. 2002;33:332–336.
- Masuda A, Wakasugi M, Joumura K, et al. Flumazenil antagonism of midazolam-induced respiratory depression. *Masui*. 1994;43:722–727.
- Maze M, Bonett F. Analgesics. Receptor ligands-α₂ adrenergic receptor agonists. In: Evers AS, Maze M. *Anesthetic pharmacology: physiologic principles and clinical practice. A Companion to Miller's Anesthesia*. Livingstone: Churchill, 2004.
- McGlone R, Fleet T, Durham S, et al. A comparison of intramuscular ketamine with high dose intramuscular midazolam with and without intranasal flumazenil in children before suturing. *Emerg Med J*. 2001;18:34–38.

- Mertes N, Goeters C, Kuhmann M, et al. Postoperative alpha₂-adrenergic stimulation attenuates protein catabolism. *Anesth Analg.* 1996;82:258–263.
- Möhler H, Okada T. Benzodiazepine receptors. Demonstrations in the central nervous system. *Science.* 1877;198:849–851.
- Möhler H, Rudolph V, Crestani F, et al. Sedatives anxiolytics and amnestic. In: Evers AS, Maze M. Anesthetic pharmacology: physiologic principles and clinical practice. A Companion to Miller's Anesthesia. Livingstone: Churchill, 2004.
- Moore EN, Spear JF. Acute animal models for the study of antiarrhythmic drugs for the prevention of sudden coronary death. In: Clinical pharmacology of antiarrhythmic therapy (Ed. Benedict R, Lucchesi JV, Dingell A, et al.). New York: Raven Press, 1984.
- Mora CT, Torjman M, White PF. Sedative and ventilatory effects of midazolam infusion: effect of flumazenil reversal. *Can J Anaesth.* 1995;42:677–684.
- Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia a delay of lethal cell injury in ischemic myocardium. *Circulation.* 1986;74:1124–1136.
- Mushin W. Intramuscular gallamine with hyaluronidase. *Anaesthesia.* 1956;111:135–138.
- Nave R, Herer P, Lavie P. The intrinsic effects of sarmazenil on sleep propensity and performance of sleep-deprived subjects. *Psychopharmacology (Berl.).* 1994;115: 366–370.
- Neave N, Reid C, Scholey AB, et al. Dose-dependent effects of flumazenil on cognition, mood, and cardio-respiratory physiology in healthy volunteers. *Br Dent J.* 2000;189:668–674.
- Nelson LE, Lu J, Guo T, et al. The alpha₂-adrenoceptor agonist dexmedetomidine converges on an endogenous sleep-promoting pathway to exert its sedative effects. *Anesthesiology.* 2003;98:428–436.
- Nielsen BN, Friis SM, Romsing J, et al. Intranasal sufentanil/ketamine analgesia in children. *Pediatric Anesthesia.* 2014;170–180.
- Ochs MW, Tucker MR, Owsley TG, et al. The effectiveness of flumazenil in reversing the sedation and amnesia produced by intravenous midazolam. *J Oral Maxillofac Surg.* 1990;48:240–245.
- Oliver FM, Sweatman TW, Unkel JH, et al. Comparative pharmacokinetics of submucosal vs. intravenous flumazenil (Romazicon) in an animal model. *Pediatr Dent.* 2000;22:489–493.
- Paddleford RR, Erhardt W. Allgemeinanästhetika. In: Anästhesie bei Kleintieren. Paddleford RR (ed.). Philadelphia: Saunders, 1999:78–88.
- Palmer RB, Mautz DS, Cox K, et al. Endotracheal flumazenil: a new route of administration for benzodiazepine antagonism. *Am J Emerg Med.* 1998;16:170–172.
- Pfenninger E, Baier Ch, Claus S, et al. Untersuchung zu psychometrischen Veränderungen sowie tur analgetischen Wirkung und kardiovaskulären Nebenwirkungen von Ketamin-Razemat versus S-(+)-Ketamin in subanästhetischer Dosierung. *Anesthesist.* 1994;43(Suppl. 2):68–75.
- Phillips JW. Adenosines role in the central actions of the benzodiazepines. *Prog Neuro-Psychopharmacol Biol Psychiat.* 1984;8:495–602.
- Pieri L, Schaffner R, Scherschlicht R, et al. Pharmacology of midazolam. *Arzneim-Forsch./Drug Res.* 1981;31:2180–2201.
- Polc P. Electrophysiology of benzodiazepine receptor ligands: Multiple mechanism and sites of action. *Neurobiology.* 1988;31:349–423.

- Polc P, Bonetti EP, Pieri L, et al. Caffeine antagonizes several central effects of diazepam. *Life Sci.* 1981;28:2265–2275.
- Port JD, Stanley TH, McJames S. Neurosciences and anesthetic action III. Anesthesiology. 1983;59(Suppl):A325.
- Port JD, Stanley TH, Steffey EM. Narcotic inhalation anesthesia. *Anesthesiology*. 1982;57(Suppl):A344.
- Port JD, Stanley TH, Steffey EP, et al. Intravenous carfentanil in the dog and rhesus monkey. *Anesthesiology*. 1984;61(Suppl):A378.
- Poulsen H, Hougs W. The effect of some curarizing drugs in unanesthetized man. I. *Acta Anaesth Scandinav.* 1957;1:15–39.
- Proctor LT, Schmeling WT, Roerig D, et al. Oral dexmedetomidine attenuates hemodynamic responses during emergence from general anesthesia in chronically instrumented dogs. *Anesthesiology*. 1991;74:108–114.
- Proctor LT, Schmeling WT, Warltier DC. Premedication with oral dexmedetomidine alters hemodynamic actions of intravenous anesthetic agents in chronically instrumented dogs. *Anesthesiology*. 1992;77:554–562.
- Prommer E, Thompson L. Intranasal fentanyl for pain control: current status with a focus on patient considerations. *Patient Prefer Adherence*. 2011;5:157164.
- Ramsay EC, Wetzel RW. Comparison of five regimens for oral administration of medication to induce sedation in dogs prior to euthanasia. *J Am Vet Med Assoc*. 1998;213:240–242.
- Reves JG, Fragen RJ, Vinik HR, et al. Midazolam: pharmacology and uses. *Anesthesiology*. 1985;62:310–324.
- Rey E, Delaunay L, Pons G, et al. Pharmacokinetics of midazolam in children: comparative study of intranasal and intravenous administration. *Eur J Clin Pharmacol*. 1991;41:355–357.
- Riches JR, Read RW, Black RM, Cooper NJ. Analysis of clothing and urine from Moscow theatre siege casualties reveals carfentanil and remifentanil use. *J Anal Toxicol*. 2012;36:647–656.
- Rivo J, Raphael J, Drenger B, et al. Flumazenil mimics whereas midazolam abolishes ischemic preconditioning in a rabbit heart model of ischemia-reperfusion. *Anesthesiology*. 2006;105:65–71.
- Roald OK, Forsman N, Steen PA. Partial reversal of the cerebral effects of isoflurane in the dog by the benzodiazepine antagonist flumazenil. *Acta Anaesthesiol Scand*. 1988;32:209–212.
- Roekaerts P, Prinzen F, Willigers H, et al. The effect of dexmedetomidine on systemic haemodynamics, regional myocardial function and blood flow during coronary artery stenosis in acute anaesthetized dogs. *J Cardiothor Vascul Anesthesia*. 1994;8(Suppl. 3):58.
- Salonen M, Reid K, Maze M. Synergistic interaction between α_2 -adrenergic agonistic and benzodiazepines in rats. *Anesthesiology*. 1992;76:1004–1011.
- Salonen M, Reid K, Maze M. Synergistic interaction between alpha₂-adrenergic agonists and benzodiazepines in rats. *Anesthesia*. 1994;8(Suppl. 3):1004–1011.
- Scott RC, Besag FMC, Boyd SG, et al. Buccal absorption of midazolam: pharmacokinetics and EEG pharmacodynamics. *Epilepsia*. 1998;39:290–294.

- Segal IS, Vickery RG, Walton JK, et al. Dexmedetomidine diminishes halothane anesthetic requirements in rats through a postsynaptic alpha₂-adrenergic receptor. *Anesthesiology*. 1988;69:818–823.
- Seifritz E, Hemmeter U, Trachsel L, et al. Effects of flumazenil on recovery sleep and hormonal secretion after sleep deprivation in male controls. *Psychopharmacology* (Berl.). 1995;120:449–456.
- Scheinin H, Aantaa R, Anttila M, et al. Reversal of the sedative and sympatholytic effects of dexmedetomidine with a specific alpha₂-adrenoceptor antagonist atipamezole. *Anesthesiology*. 1998;89:574–584.
- Schaller G. 1966. In: Harthoorn AM. *The chemical capture of animals* London: Baillière Tindall, 1976.
- Scheepers LD, Montgomery CJ, Kinahan AM, et al. Plasma concentrations of flumazenil following intranasal administration in children. *Canadian J Anesthesia*. 2000;47:120–124.
- Scheinin H, Karhuvaara S, Olkkola KT, et al. Pharmacodynamics and pharmacokinetics of intramuscular dexmedetomidine. *Clin Pharmacol Ther*. 1992;52:537–546.
- Scheinin M, Kallio A, Koulu M, et al. Sedative and cardiovascular effects of medetomidine, a novel selective alpha₂-adrenoceptor agonist, in healthy volunteers. *Br J Clin Pharmacol*. 1987;24:443–451.
- Scheinin M, Schwinn DA. The Locus Coeruleus. Site of hypnotic actions of alpha₂-adrenoreceptor agonists? *Anesthesiology*. 1992;76:873–875.
- Schort CE. Alpha₂-agents in animals – sedation, analgesia and anaesthesia. Santa Barbara USA: Veterinary Practice Publishing Company, 1992.
- Schwagmeier R, Alincic S, Striebel HW. Midazolam pharmacokinetics following intravenous and buccal administration. *Br J Clin Pharmacol*. 1998;46:203–206.
- Schwartz AE, Manaksha FR, Kanchuger MS. Flumazenil decreases the minimum alveolar concentration isoflurane in dogs. *Anesthesiology*. 1989;70:764–766.
- Schweiger IM, Szlam F, Hug CC J. Absence of agonistic or antagonistic effect of flumazenil (Ro 15-1788) in dogs anesthetized with enflurane, isoflurane, or fentanyl – enflurane. *Anesthesiology*. 1989;70:477–480.
- Schweizer E, Clary C, Dever AI, et al. The use of low-dose intranasal midazolam to treat panic disorder: a pilot study. *J Clin Psychiatry*. 1992;53:19–22.
- Sleeman JM, Carter W, Tobin T, et al. Immobilization of domestic goats (*Capra hircus*) using orally administered carfentanil citrate and detomidine hydrochloride. *J Zoo Wildl Med*. 1997;28:158–165.
- Sliva J, Hess L, Votava M, et al. Endogenous ligands of benzodiazepine binding site have inverse agonistic properties. *Med Hypotheses*. 2013;81:1075–1077.
- Spinadel L. *Klinická anesteziologie*. Praha: Naše vojsko, 1950:280.
- Squires RF, Braestrup C. Benzodiazepine receptors in rat brain. *Nature*. 1977;266:732–734.
- Stanley TH, Hague B, Mock DL, et al. Oral transmucosal fentanyl citrate (lollipop) premedication in human volunteers. *Anesth Analg*. 1989;69:21–27.
- Stegmann GF, Bester L. Sedative-hypnotic effects of midazolam in goats after intravenous and intramuscular administration. *Veterinary Anaesthesia Analgesia*. 2001;28:49–55.

- Steiger A, Guldner J, Lauer CJ, et al. Flumazenil exerts intrinsic activity on sleep EEG and nocturnal hormone secretion in normal controls. *Psychopharmacology (Berl.)*. 1994;113:334–338.
- Streisand JB, Varvel JR, Stanski DR, et al. Absorption and bioavailability of oral transmucosal fentanyl citrate. *Anesthesiology*. 1991;223–229.
- Striebel HW, Koenigs D, Krämer J. Postoperative pain management by intranasal demand-adapted fentanyl titration. *Anesthesiology*. 1992;77:281–285.
- Striebel HW, Kramer J, Luhmann I, et al. Pharmakokinetische Studie zur intranasalen Gabe von Fentanyl. *Der Schmerz*. 1993;7:122–125.
- Striebel HW, Pommerening J, Rieger A. Intranasal fentanyl titration for postoperative pain management in an unselected population. *Anaesthesia*. 1993;48:753–757.
- Suter-Zimmermann K. Transmucosal nasal drug delivery, systemic bioavailability of nasaly applied midazolam. Basel: Dissertation, 2008.
- Swaab D. Wir sind unser Gehirn. München: Droemer Verlag, 2011.
- Šustková-Fišerová M, Vávrová J, Kršiak M. Brain levels of GABA, glutamate and aspartate in sociable, aggressive and timid mice an in vivo microdialysis study. *Neuro Endocrinol Lett*. 2009;30:79–84.
- Talon MD, Woodson LC, Sherwood E, et al. Nasal dexmedetomidine is comparable to midazolam as a preoperative sedative for children. *Anesthesiology*. 2007;107:A1398.
- Tobbin J. Paradoxical effects of midazolam in the very young. *Anesthesiology*. 2008; 108:6–7.
- Tolksdorf W, Bremer H, Tokic B. Gibt es einen atemdepressiven Effekt von Flumazenil. In: Anexate (Flumazenil). Der erste spezifische Benzodiazepin-Antagonist. Editiones Roche, 1989:173–184.
- Uhlířová L, Šustková-Fišerová M, Kršiak M. Behavioral effects of flumazenil in the social conflict in mice. *Psychopharmacology*. 2004;171:259–269.
- Unseld E, Klotz U. Benzodiazepines: Are they of natural origin? *Pharm Res*. 1989;6:1–3
- Vaculín Š, Franěk M, Hess L, et al. Ovlivnění nocicepcí medetomidinem a jeho kombinací s ketaminem u laboratorního potkana. *Psychiatrie*. 2000;6(Suppl. 4):14–17.
- Vartiainen J, MacDonald E, Urtti A, et al. Dexmedetomidine-induced ocular hypotension in rabbits with normal or elevated intraocular pressure. *Invest Ophthalmol Vis Sci*. 1992;33:2019–2023.
- Velez L, Shepherd G, Mills LD, Rivera W. Systemic toxicity after an ocular exposure to xylazine hydrochloride. *J Emerg Med*. 2006;30:407–410.
- Vercauteren M, Boeckx E, Hanegreefs G, et al. Intranasal sufentanil for pre-operative sedation. *Anaesthesia*. 1988;43:270–273.
- Verghese ST, Hannallah RS, Brennan M, et al. The effect of intranasal administration of remifentanil on intubating conditions and airway response after sevoflurane induction of anesthesia in children. *Anesth Analg*. 2008;107:1176–1181.
- Vetter W, Wildmann J, Binder A, et al. Ist der Wirkstoff von „Valium Roche“ ein naturidentisches Produkt? *Roche Magazin*. 1988;31:6–15.
- Vickery RG, Sheridan BC, Segal IS, et al. Anesthetic and hemodynamic effects of the stereoisomers of medetomidine, an alpha₂-adrenergic agonist, in halothaneranessthetized dogs. *Anesth Anal*. 1988;67:611–615.
- Virtanen R. Pharmacological profiles of medetomidine and its antagonists, atipamezole. *Acta Vet Scand*. 1989;85:29–37.

- Votava M, Hess L, Kršiak M. Selective antiaggressive effect of an alpha-2 adrenoceptor agonist naphthylmedetomidine in mice. *Aggressive Behavior*. 2008;34:394–403.
- Votava M, Hess L, Slíva J, et al. Dexmedetomidine selectively suppresses dominant behaviour in aggressive and sociable mice. *Eur J Pharmacol*. 2005;523:79–85.
- Vrána M, Fejfar Z, Netušil M, et al. Stimulation threshold studies and effect of antiarrhythmic drugs. *Basic Res Cardiol*. 1978;73:618.
- Walzer C, Huber CH. Comparison of two benzodiazepine antagonists: flumazenil and sarmazenil in cheetah (*Acinonyx jubatus*). *Erkrankungen der Zootiere, VISEZ39 Wien*. Berlin: IZW, 1999:377–382.
- Watson D. Hyaluronidase. *Brit J Anaesthesia*. 1993;71:422–425.
- Weinbroum A, Geller E. Flumazenil improves cognitive and neuromotor emergence and attenuates shivering after halothane, enflurane and isoflurane based anesthesia. *Can J Anaesth*. 2001;48:963–72.
- Weinbroum A, Weisenberg M, Rudick V, et al. Flumazenil potentiation of postoperative morphine analgesia. *Clin J Pain*. 2000;16:193–199.
- Weinger MB, Partridge BL, Henry AF. Dexmedetomidine does not modify the neuromuscular blocking action of vecuronium in the anaesthetized rat. *Br J Anaesth*. 1995;74:455–457.
- Weiske A. Untersuchungen einer Injectionsnarkose bei der Taube (*Columbia livia*, var. Dom. Gmel. 1789) mit Medetomidin, Midazolam und Fentanyl und ihrer vollständigen Antagonisierung mit Atipamezol, Sarmazenil und Naloxon. *Vet Med München*: Dissertation, 2001.
- Weksler N, Ovadia L, Muati G, et al. Nasal ketamine for paediatric premedication. *Can J Anaesth*. 1993;40:119–121.
- Whittington RA, Virag L, Vulliemot Y, et al. Dexmedetomidine increases the cocaine seizure threshold in rats. *Anesthesiology*. 2002;97:693–700.
- Whitwam, JG. Drugs for sedation. In: *Quality control in endoscopy*. Ed. McCloy R. Springer Verlag, 1991:21–28.
- Whitwam JG. Midazolam-Flumazenil. Der neueste Stand Minim. Invas Ther. 1995(Suppl. 4):31–38.
- Wiener-Kronish JP, Gropper MA. Conscious sedation. Philadelphia: Hanley and Belfus, 2001.
- Wiesner H, Hegel G. Von Praktische Hinweise zur Immobilisation von Wild-und Zoo-tieren. *Tierärztl Prax*. 1985;13:113–127.
- Worsley MH, Macleod AD, Brodie MJ, et al. Inhaled fentanyl as a method of analgesia. *Anaesthesia*. 1990;45:449–445.
- Zehentbauer J. Körpereigene Drogen. Die ungenutzten Fähigkeiten unseres Gehirns. 1. Auflage. Patmos Verlag, 1992.
- Zhang J, Maland L, Hague B, et al. Buccal absorption of etomidate from a solid formulation in dogs. *Anesth Analg*. 1998;86:1116–1122.
- Ziegler WH, Schalch E. Antagonism of benzodiazepine-induced sedation in man. In: Koella W, ed. *Sleep* 1982. Basel: Karger, 1983:427–429.
- Zöllner Ch, Mousa S, Klinger A, et al. Topical fentanyl in a randomized, double-blind study in patients with corneal damage. *Clin J Pain*. 2008;24:690–696.