

## Literatura

- [1] ADUEN, JF., et al. An alternative echocardiographic method to estimate mean pulmonary artery pressure. *J. Am. Soc. Echo.*, 2009, 22, p. 814–819.
- [2] American Society of Echocardiography (ASE): *ASE/EACVI-Guidelines*, 2016.
- [3] American Society of Echocardiography Committee on Standards, Subcommittee on Quantitation of Two-Dimensional Echocardiograms, SCHILLER, NB., et al. Recommendations for Quantitation of the left Ventricle by Two-Dimensional Echocardiography. *J. Am. Soc. Echocardiogr.*, 1989, 2, p. 358–367.
- [4] BANGALORE, S., et al. Role of right ventricular wall motion abnormalities in risk stratification and prognosis of patients referred for stress echocardiography. *JACC*, 2007, 50, p. 1981–1989.
- [5] BIAMINO, G., LANGE, L. *Echokardiographie*. Frankfurt: Hochst, 1983.
- [6] BOEKSTEGERS, P., et al. Interventionelle Behandlung der Mitralklappeninsuffizienz mit dem MitraClip-Verfahren. *Kardiologe*, 2013, 7, p. 91–104.
- [7] BONOW, RO., et al. ACC/AHA 2006 Guidelines for the management of patients with valvular heart disease. *Circulation*, 2006, 114, p. 84–231.
- [8] CERQUEIRA, MD., et al. Standardized myocardial segmentation and tomographic imaging of the heart. *Circulation*, 2002, 105, p. 539–542.
- [9] DALEN, H., et al. Segmental and global longitudinal strain and strain rate based on echocardiography of 1266 healthy individuals: the HUNT study in Norway. *Eur. J. Echo.*, 2010, 11, p. 176–183.
- [10] DANIEL, W., et al. Comparison of transthoracic and transesophageal echocardiography for detection and abnormalities of prosthetic and bioprosthetic valve in the mitral and aortic positions. *Am. J. Cardiol.*, 1993, 71, p. 210–215.
- [11] DENNIG, K., et al. Doppler-echokardiographische Bestimmung des Schweregrades der Trikuspidalstenose. *Herz*, 1986, 6, p. 332.
- [12] DITTMANN, H., et al. Vergleich doppler-echokardiographischer Methoden zur Herzminutenvolumenbestimmung. *Z. Kardiol.*, 1987, 76, p. 433.

- [13] DUMESNIL, JG., et al. Validation and applications of indexed aortic prosthetic valve areas calculated by doppler echocardiography. *J. Am. Cali. Cardiol.*, 1990, 16, p. 637–643.
- [14] DUMESNIL, JG., et al. Validation and applications of mitral prosthetic valvular areas calculated by doppler echocardiography. *J. Am. Cali. Cardiol.*, 1990, 65, p. 443–448.
- [15] DURACK, DT., et al. New criteria for diagnosis of infective endocarditis: utilization of specific echocardiographic findings. *Am. J. Med.*, 1994, 96, p. 200–209.
- [16] FELDMAN, T., et al. EVEREST I. *JACC*, 2009, 54, p. 686.
- [17] FELDMAN, T., et al. Percutaneous Repair or Surgery for Mitral Regurgitation. *N. Engl. J. Med.*, 2011, 641, p. 395–406.
- [18] GALIE, N., et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur. Heart J.*, 2015, 30, p. 2493–2537.
- [19] GARDIN, J., et al. Evaluation of blood flow velocity in the ascending aorta and main pulmonary artery of normal subjects by doppler echocardiography. *Am. Heart J.*, 1984, 107, p. 310.
- [20] GILMAN, G., et al. Sonographic approach to the essential echocardiographic measurements of left ventricular diastolic function. *J. Am. Soc. Echo.*, 2007, 20, p. 199–209.
- [21] HABIB, G., et al. 2015 Guidelines for the management of infective endocarditis. *Eur. Heart J.*, 2015, 36, p. 3075–3123.
- [22] HATLE, L., et al. Non-invasive assessment of atrioventricular pressure half-time by doppler ultrasound. *Circulation*, 1979, 60, p. 1096.
- [23] HATLE, L., et al. Noninvasive assessment of pressure drop in mitral stenosis by doppler ultrasound. *B. Heart J.*, 1978, 40, p. 131–140.
- [24] HATLE, L., ANGELSON, B. *Doppler ultrasound in cardiology. Physical principles and clinical applications*. Philadelphia: Lea&Febiger, 1985.
- [25] HATLE, L. *Doppler Ultrasound in Cardiology*. Philadelphia: Lea&Febiger, 1985.
- [26] HELMCKE, F., et al. Color doppler assessment of mitral regurgitation with orthogonal planes. *Circulation*, 1987, 75, p. 175.
- [27] KHOURI, SJ., et al. A practical approach to the echocardiographic evaluation of diastolic function. *J. Am. Soc. Echo.*, 2004, 17, p. 290–297.
- [28] KISSLO, J., et al. *Basic Doppler Echocardiography*. New York: Churchill Livingstone, 1988.

- [29] KITABATAKE, A., et al. Non-invasive evaluation of pulmonary hypertension by pulsed doppler technique. *Circulation*, 1983, 68, p. 302.
- [30] KLEIN, A., COHEN, G. Doppler echocardiographic assessment of constrictive pericarditis, cardiac amyloidosis, and cardiac tamponade. *Cleve. Clin. J. Med.*, 1992, 59, p. 281.
- [31] KONSTANTINIDES, SV., TORBICKI, A., AGNELLI, G., et al. 2014 ESC Guidelines on the diagnosis and management of acute pulmonary embolism. *Eur. J. Heart*, 2014, 35, p. 3033–3069.
- [32] LABOVITZ, A. *Doppler Echocardiography*. Philadelphia: Lea&Febiger, 1988.
- [33] LANCELLOTTI, P., et al. *The EACVI Echo Handbook*. Oxford: Oxford University Press, 2016.
- [34] LANG, R., et al. Recommendations for chamber quantification by echocardiography in adults. *J. Am. Soc. Echo.*, 2015, 28, p. 1–39.
- [35] LANG, R., et al. Recommendations for chamber quantification. *J. Am. Soc. Echo.*, 2005, 18, p. 1440–1463.
- [36] LEE, K., et al. Echocardiographic assessment of right heart dynamic parameters. *J. Am. Soc. Echo.*, 2007, 20, p. 773–782.
- [37] MARCUS, FI., et al. Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the task force criteria. *Circulation*, 2010, 121, p. 1533.
- [38] MARON, BJ., et al. Contemporary Definitions and Classification of the Cardiomyopathies. *Circulation*, 2006, 113, p. 1807–1816.
- [39] MARWICK, TH. Recommendations on the use of echocardiography in adult hypertension. *EHJCI*, 2015, 16, p. 577–605.
- [40] MASUYAMA, T., et al. Continuous-wave doppler echocardiographic detection of pulmonary regurgitation and its application to noninvasive estimation of pulmonary artery pressure. *Circulation*, 1986, 74, p. 484–492.
- [41] MASUYAMA, T., et al. Pulmonary arterial end-diastolic pressure non-invasively estimated by continuous wave doppler echocardiography. *J. Cardiogr.*, 1986, 16, p. 669.
- [42] MILLER, G. *Invasive investigation of the heart. A guide to cardiac catheterization and related procedures*. Oxford: Blackwell, 1989, p. 271–274.
- [43] NADA, A. Pulmonary stenosis, aortic stenosis, ventricular septal defect: Clinical course and indirect assessment. *Circulation*, 1977, 55, p. 138.

- [44] NAGUEH, S., et al. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *Eur. J. Echo.*, 2009, 10, p. 165–193.
- [45] NAGUEH, S., et al. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *J. Am. Soc. Echo.*, 2016, 29, p. 277–314.
- [46] National Cancer Institute. Common Terminology Criteria for Adverse Events (CTCAE v. 2.0), 1999.
- [47] NISHIMURA, R., OTTO, CM., BONOW, RO., et al. 2014 AHA/ACC Guideline for the Management of Patients with Valvular Heart Disease. *J. Am. Cali. Cardiol.*, 2014, 63, p. e57–e185.
- [48] OH, JK., et al. *The Echo Manual*. Philadelphia: Lippincott, Williams&Wilkins, 1999.
- [49] OH, JI., et al. The noninvasive assessment of left ventricular diastolic function with two-dimensional and doppler echocardiography. *J. Am. Soc. Echo.*, 1997, 10, p. 246–270.
- [50] PIBAROT, P. Prosthesis patient mismatch: definition, clinical impact, and prevention. *Heart*, 2006, 92, p. 1022–1029.
- [51] RICHARDSON, P., et al. Report of the 1999 World Health Organisation/International Society and Federation of Cardiology Task Force on the definition and classification of cardiomyopathies. *Circulation*, 1996, 93, p. 841–842.
- [52] ROSENHEK, R. Normal values for doppler echocardiographic assessment of heart valve prostheses. *JASE*, 2003, 16, p. 1116–1127.
- [53] RUDSKI, LG., et al. Guidelines for the echocardiographic assessment of the right heart in adults. *J. Am. Soc. Echo.*, 2010, 23, p. 685–713.
- [54] SCHILLER, NB., et al. Recommendation for quantification of the left ventricle by 2-D echocardiography. *J. Am. Soc. Echo.*, 1989, 2, p. 358–367.
- [55] SHIVELY, BI., et al. Diagnostic value of transesophageal compared with transthoracic echocardiography in infective endocarditis. *J. Am. Cali. Cardiol.*, 1991, 18, p. 391–397.
- [56] SIEVERS, HH., et al. A classification system for the bicuspid aortic valve from 304 surgical specimen. *J. Thorac. Cardiovasc. Surg.*, 2007, 133, p. 1226–1233.
- [57] SPETHMANN, S., et al. Doppler haemodynamics and effective orifice areas of Edwards SAPIEN and CoreValve transcatheter aortic valves. *EHJCVI*, 2012, 13, p. 690–696.

- [58] TEI, C., et al. Non-invasive doppler-derived myocardial performance index: correlation with simultaneous measurements of cardiac catheterization measurements. *J. Am. Soc. Echo.*, 1997, 10, p. 169–178.
- [59] VAHANIAN, A., et al. ESC guidelines on the management of valvular heart disease. *Eur. Heart J.*, 2007, 28, p. 230–268.
- [60] VARGHESE, V., et al. EVEREST II. *JACCCI*, 2011, 4, p. 825.
- [61] WARNES, CA., WILLIAMS, RG., BASHORE, TM., et al. ACC-AHA 2008 Guidelines for the Management of Adults With Congenital Heart Disease. *J. Am. Cali. Cardiol.*, 2008, 52, p. e143–e263.
- [62] WEIDEMANN, F., et al. Verschiedene Formen der Linksventrikulären Hypertrophie. *Herz*, 2011, DOI: 10.1007/s00059-010-3416-1.
- [63] WILKENSHOFF, U., et al. Age dependent changes in regional diastolic function evaluated by color doppler myocardial imaging. A comparison with pulsed doppler indices of global function. *J. Am. Soc. Echo.*, 2001, 14, p. 959–969.
- [64] WILKENSHOFF, U., et al. Regional mean systolic velocity estimation by realtime color doppler myocardial imaging: a new technique for quantifying regional systolic function. *J. Am. Soc. Echo.*, 1998, 11, p. 683–692.
- [65] WILKENSHOFF, U., KRUCK, I., GAST, D. Validity of continuous wave doppler und colour doppler in the assessment of aortic regurgitation. *Eur. Heart J.*, 1994, 15, p. 1227.
- [66] WILKINS, GT., et al. Percutaneous balloon dilatation of the mitral valve: an analysis of echocardiographic variables related to outcome and mechanism of dilatation. *Br. Heart J.*, 1988, 60, p. 299.
- [67] XIE, GY., et al. Quantification of mitral regurgitant volume by the color doppler proximal isovelocity surface area method: A clinical study. *J. Am. Soc. Echo.*, 1995, 8, p. 48–54.
- [68] ZOGHBI, WA., et al. Recommendations for evaluation of prosthetic valves with echocardiography and doppler ultrasound. *J. Am. Soc. Echo.*, 2009, 229, p. 975–1013.
- [69] ZOGHBI, WA., et al. Recommendations for evaluation of the severity of native valvular regurgitation with two-dimensional and doppler echocardiography. *J. Am. Soc. Echo.*, 2003, 16, p. 777–802.