



**Fig. 11** Schematic diagram of the control volume that illustrates the fluid-solid mass-transfer during a chemical reaction in the washcoat with resistances using a series approach.

## REFERENCES

- [1] T.J. Mason, J.P. Lorimer, *Applied Sonochemistry: Uses of Power Ultrasound in Chemistry and Processing*, Wiley-VCH, Weinheim, Germany, 2002. 314 p.
- [2] J.-L. Capelo-Martínez (Ed.), *Ultrasound in Chemistry: Analytical Applications*, Wiley-VCH, Weinheim, Germany, 2009. 171 p.
- [3] K.S. Suslick, *Sonochemistry and sonoluminescence*, in: *Encyclopedia of Physical Science and Technology*, third ed., vol. 17, Academic Press, San Diego, CA, 2001, pp. 363–376.
- [4] W. Carruthers, I. Coldham, *Modern Methods of Organic Synthesis*, fourth ed., Cambridge University Press, England, U.K., 2004. 506 p.

- [5] J.N. Lalena, D.A. Cleary, E. Carpenter, N.F. Dean, Inorganic Materials: Synthesis and Fabrication, Wiley-Interscience, New York, 2008. 303 p.
- [6] K.S. Suslick, S.E. Skrabalak, Sonocatalysis, in: G. Ertl, H. Knözinger, F. Schüth, J. Weitkamp (Eds.), Handbook of Heterogeneous Catalysis, vol. 4, Wiley-VCH, Weinheim, Germany, 2008, pp. 2006–2017.
- [7] J.-M. Basset, R. Psaro, D. Roberto, R. Ugo (Eds.), Modern Surface Organometallic Chemistry, Wiley-VCH, Weinheim, Germany, 2009. 725 p.
- [8] K.S. Suslick, Applications of ultrasound to materials chemistry, MRS Bull. (1995) 29–34.
- [9] P.W. Cains, P.D. Martin, C.J. Price, The use of ultrasound in industrial chemical synthesis and crystallization. 1. Applications to synthetic chemistry, Org. Process. Res. Dev. 2 (1) (1998) 34–48.
- [10] G. Cravotto, P. Cintas, Power ultrasound in organic synthesis: moving cavitation chemistry from academia to innovative and large-scale applications, Chem. Soc. Rev. 35 (2) (2006) 180–196.
- [11] K.S. Suslick, D.J. Flannigan, Sonoluminescence, Annu. Rev. Phys. Chem. 59 (2008) 659–683.
- [12] J.H. Bang, K.S. Suslick, Applications of ultrasound to the synthesis of nanostructured materials, Adv. Mater. 22 (2010) 1039–1059.
- [13] SONITEK: Ultrasonic horns. <http://www.usedultrasonicwelders.com/Horn%20Brochure.pdf> (accessed 10.07.17).
- [14] G. Cravotto, E. Calcio Gaudino, P. Cintas, On the mechanochemical activation by ultrasound, Chem. Soc. Rev. 42 (2013) 7521–7534.
- [15] S.C. Ameta, P.B. Punjabi, H. Swarnkar, N. Chhabra, M. Jain, Sonochemistry: past, present and future, J. Indian Chem. Soc. 78 (10–12) (2001) 627–633.
- [16] Y.-J. Bian, J.-T. Li, T.-S. Li, Applications of ultrasound in organic synthesis using metal reagents, Youji Huaxue 22 (4) (2002) 227–232.
- [17] B.I. Kharisov, O.V. Kharissova, U. Ortiz-Méndez, Use of ultrasound in coordination and organometallic chemistry, in: D. Chen, S.K. Sharma, A. Mudhoo (Eds.), Handbook on Applications of Ultrasound: Sonochemistry for Sustainability, 2011, Taylor & Francis Group, LLC, Boca Raton, FL, USA, 2011, pp. 183–211.
- [18] M. Shi, X. Chen, Preparation of 1,4-bis(dimethylsilyl)benzene under ultrasonic irradiation, Fujian Fenxi Ceshi 16 (3) (2007) 70–73.
- [19] M.-H. Shi, X.-J. Chen, Preparation of 1,4-bis(dimethylsilyl)benzene under ultrasonic irradiation, Huagong Jishu Yu Kaifa 36 (9) (2007) 8–10.
- [20] X. Chen, Y. Cui, G. Yin, Z. Jia, Preparation and thermal property of poly(tetramethyl-p-silphenylene-dimethyl) siloxane, Huagong Jinzhan 26 (9) (2007) 1333–1337.
- [21] U. Kunz, U. Hoffmann, A. Rosenplaenter, C. Horst, Use of ultrasonics for solid/liquid reaction. Patent DE 19808239, 1999, 20 p.
- [22] A.S.-Y. Lee, S.-F. Chu, Y.-T. Chang, S.-H. Wang, Synthesis of homopropargyl alcohols via sonochemical Barbier-type reaction, Tetrahedron Lett. 45 (7) (2004) 1551–1553.
- [23] B.I. Kharisov, L.A. Garza-Rodríguez, H. Leija Gutiérrez, U. Ortiz Méndez, R. García Caballero, A.Y. Tsividze, Preparation of non-substituted metal phthalocyanines at low temperature using activated Rieke zinc and magnesium, Synth. React. Inorg. Metal-Org. NanoMetal Chem. 35 (10) (2005) 755–760.
- [24] B.I. Kharisov, A.D. Garnovskii, O.V. Kharissova, U. Ortiz Mendez, A.Y. Tsividze, Direct electrochemical synthesis of metal complexes of phthalocyanines and azomethines as model compounds: advantages and problems of this method versus traditional synthetic techniques, J. Coord. Chem. 60 (13) (2007) 1435–1455.
- [25] B.I. Kharisov, L.M. Blanco, L.M. Torres-Martínez, A. García-Luna, Electrosynthesis of metal phthalocyanines: influence of solvent, Ind. Eng. Chem. Res. 38 (8) (1999) 2880–2887.
- [26] B.I. Kharisov, L.M. Blanco, M.V. Salinas, A.D. Garnovskii, Direct electrochemical synthesis of copper dimethyldithiocarbamate: an optimization using ultrasonic treatment, J. Coord. Chem. 47 (1999) 135–143.
- [27] A.J. Hulkes, A.F. Hill, B.A. Nasir, A.J.P. White, D.J. Williams, Reactions of  $\mu$ -alkylidyne complexes with tellurium. Telluroacyl versus  $\mu$ -telluride formation, Organometallics 23 (4) (2004) 679–686.

- [28] R.D. Rieke, Chemical Synthesis Using Highly Reactive Metals, Wiley, 2016. 480 p.
- [29] L.A. Garza-Rodríguez, B.I. Kharisov, O.V. Kharissova, Overview on the synthesis of activated micro- and nanostructurized Rieke metals: history and present state, *Synth. React. Inorg. Met.-Org. Nano-Met. Chem.* 39 (5) (2009) 270–290.
- [30] R.D. Rieke, Use of activated metals in organic and organometallic synthesis, *Top. Curr. Chem.* 59 (1975) 1–31.
- [31] R.D. Rieke, Preparation of highly reactive metal powders and their use in organic and organometallic synthesis, *Acc. Chem. Res.* 10 (1977) 301–306.
- [32] L.C. Chao, R.D. Rieke, W.R.K. Jr, Activated metals: VI. Preparation and reactions of highly reactive indium metal, *J. Organomet. Chem.* 67 (3) (1974) C64–C66.
- [33] L.C. Chao, R.D. Rieke, Activated metals. VII. Improved method for the synthesis of trialkyl and triaryl indium compounds, *Synth. React. Inorg. Metal-Org. Chem.* 4 (4) (1974) 373–378.
- [34] L.C. Chao, R.D. Rieke, Activated metals. X. Direct synthesis of diphenylindium iodide and ditolylindium iodide from activated indium metal, *Synth. React. Inorg. Metal-Org. Chem.* 5 (3) (1975) 165–173.
- [35] L.C. Chao, R.D. Rieke, Activated metals. IX. New Reformatsky reagent involving activated indium for the preparation of beta-hydroxy esters, *J. Organomet. Chem.* 40 (1975) 2253–2255.
- [36] R.D. Rieke, Preparation of organometallic compounds from highly reactive metal powders, *Science* 246 (1989) 1260–1264.
- [37] L.C. Chao, R.D. Rieke, Activated metals. IV. Preparation of highly reactive aluminum metal and the direct synthesis of phenylaluminum halides, *Synth. React. Inorg. Metal-Org. Chem.* 4 (2) (1974) 101–105.
- [38] G.A. Olah, S. Morteza, F. Faria, M.R. Bruce, F.L. Clouet, P.R. Morton, G.K. Surya Prakash, R. C. Stevens, R. Bau, et al., Aluminum dichloride and dibromide. Preparation, spectroscopic (including matrix isolation) study, reactions, and role (together with alkyl(aryl)aluminum monohalides) in the preparation of organoaluminum compounds, *J. Am. Chem. Soc.* 110 (1988) 3231–3238.
- [39] P.T.-J. Li, Dissertation: Preparation and Reactions of Rieke Metals Activated Aluminum and Zinc, University of Nebraska, 1981. 236 p. AAT 8228151, <http://digitalcommons.unl.edu/dissertations/AAI8228151/>.
- [40] S.H. Pyo, B.H. Han, Reduction of nitroarenes by activated metals, *Bull. Kor. Chem. Soc.* 16 (1995) 181–183.
- [41] T.C. Wu, H. Xiong, R.D. Rieke, Organocalcium chemistry: preparation and reactions of highly reactive calcium, *J. Organomet. Chem.* 55 (1990) 5045–5051.
- [42] P. Cintas, Activated Metals in Organic Synthesis, CRC Press, Boca Raton, 1993. Ch. 3.
- [43] R.D. Rieke, T.A. Chen, Facile synthesis of poly(phenylcarbyne): a precursor for diamondlike carbon, *Chem. Mater.* 6 (1994) 576–577.
- [44] R.D. Rieke, T.-C. Wu, L.I. Rieke, Highly reactive calcium for the preparation of organocalcium reagents: 1-adamantyl calcium halides and their addition to ketones: 1-(1-adamantyl)cyclohexanol, *Org. Synth.* 72 (1995) 147–153.
- [45] R.D. Rieke, The preparation of highly reactive metals and the development of novel organometallic reagents, *Aldrichimica Acta*, 33 (2) (2000) 52–60.
- [46] L. Pyor, A. Kiessling, Synthesis and reactions of pyridinylcalcium bromides, *Am. J. Undergrad. Res.* 1 (1) (2002) 25–28.
- [47] L.D. Rhyne. I. Activated nickel, copper, and uranium: II. Molecular orbital investigations of cyclobutadienoid compounds: III. Attempted synthesis of a new electron acceptor. Ph.D. Thesis. The University of North Carolina, Chapel Hill, NC, 1980.
- [48] R.D. Rieke, L.D. Rhyne, Preparation of highly reactive metal powders. Activated copper and uranium. The Ullmann coupling and preparation of organometallic species, *J. Organomet. Chem.* 44 (1979) 3445–3446.
- [49] F.O. Ginah, T.A. Donovan, S.D. Suchan Jr., D.R. Pfening, G.W. Ebert, Homocoupling of alkyl halides and cyclization of  $\alpha,\omega$ -dihaloalkanes via activated copper, *J. Organomet. Chem.* 55 (2) (1990) 584–589.

- [50] R.M. Wehmeyer, R.D. Rieke, Novel functionalized organocopper compounds by direct oxidative addition of zerovalent copper to organic halides and some of their reactions with epoxides, *J. Organomet. Chem.* 52 (1987) 5057–5059.
- [51] R.A. O'Brien, R.D. Rieke, Direct metalation of *p*-bromopolystyrene using highly reactive copper and the preparation and reaction of highly reactive copper bound to an insoluble polymer, *J. Organomet. Chem.* 55 (1990) 788–790.
- [52] T.-C. Wu, R.M. Wehmeyer, R.D. Rieke, Direct formation of functionalized alkylcopper reagents from alkyl halides using activated copper. Conjugate addition reactions with 2-cyclohexen-1-one, *J. Organomet. Chem.* 52 (22) (1987) 5056–5057.
- [53] G.L. Rochfort, R.D. Rieke, Preparation, characterization, and chemistry of activated cobalt, *Inorg. Chem.* 25 (1986) 348–355.
- [54] G.L. Rochfort, R.D. Rieke, Preparation of activated cobalt and its use for the preparation of octacarbonyldicobalt, *Inorg. Chem.* 23 (6) (1984) 787–789.
- [55] P.I. Volkers, C.A. Boyke, J. Chen, T.B. Rauchfuss, C.M. Whaley, S.R. Wilson, H. Yao, Haijun. precursors to [FeFe]-hydrogenase models: syntheses of  $\text{Fe}_2(\text{SR})_2(\text{CO})_6$  from CO-free iron sources, *Inorg. Chem.* 47 (15) (2008) 7002–7008.
- [56] P. Boudjouk, D.P. Thompson, W.H. Ohrbom, B.-H. Han, Effects of ultrasonic waves on the generation and reactivities of some metal powders, *Organometallics* 5 (6) (1986) 1257–1260.
- [57] R.D. Rieke, Y.S. Suh, S.-H. Kim, Heteroaryl manganese reagents: direct preparation and reactivity studies, *Tetrahedron Lett.* 46 (35) (2005) 5961–5964.
- [58] Y.S. Suh, J.-S. Lee, S.-H. Kim, R.D. Rieke, Direct preparation of benzylic manganese reagents from benzyl halides, sulfonates, and phosphates and their reactions: applications in organic synthesis, *J. Organomet. Chem.* 684 (1–2) (2003) 20–36.
- [59] Y.S. Suha, R.D. Rieke, Synthesis of  $\beta$ -hydroxy esters using highly active manganese, *Tetrahedron Lett.* 45 (8) (2004) 1807–1809.
- [60] S.-H. Kim, R.D. Rieke, A new synthetic protocol for the direct preparation of organomanganese reagents; organomanganese tosylates and mesylates, *Tetrahedron Lett.* 40 (27) (1999) 4931–4934.
- [61] J.M. Concellón, H. Rodríguez-Solla, P. Díaz, Stereoselective synthesis of (Z)-alpha-haloacrylic acid derivatives, and (Z)-haloallylic alcohols from aldehydes and trihaloesters or amides promoted by Rieke manganese, *Org. Biomol. Chem.* 6 (16) (2008) 2934–2940.
- [62] R.D. Rieke, A. Kavaliunas, Preparation of highly reactive metal powders. Preparation and reactions of highly reactive palladium and platinum metal slurries, *J. Organomet. Chem.* 44 (17) (1979) 3069–3072.
- [63] R.D. Rieke. Method of storing active zero valent zinc metal and applications in organic synthesis. US5964919, 1999, 33 p.
- [64] R.D. Rieke. Water-soluble polythiophenes. 22 pp. WO 2005066243, 2005, 22 p.
- [65] J. Kroemer, C. Kirkpatrick, B. Maricle, R. Gawrych, M.D. Mosher, D. Kaufman, *Rieke* zinc as a reducing agent for common organic functional groups, *Tetrahedron Lett.* 47 (36) (2006) 6339–6341.
- [66] A. Yanagisawa, K. Yase, H. Yamamoto, Regio- and stereoselective carboxylation of allylic barium reagents: (E)-4,8-dimethyl-3,7-nonadienoic acid, *Org. Synth. Coll. Vol.* 9 (1998) 317. 1997, 74, 178.
- [67] C.D. Stevenson, R.C. Reiter, R.D. Burton, T.D. Halvorsen, Electron transfer from rieke cadmium yielding  $\text{Cd}^{2+}$  coordinated and ion associated anion radical systems, *Inorg. Chem.* 34 (6) (1995) 1368–1372.
- [68] R.D. Rieke, W.J. Wolf, N. Kujundzic, A.V. Kavaliunas, Highly reactive transition metal powders. Oxidative insertion of nickel, palladium, and platinum metal powders into aryl-halide bonds, *J. Am. Chem. Soc.* 99 (12) (1977) 4159–4160.
- [69] R.D. Rieke, P.M. Hudnall, Activated metals. I. Preparation of highly reactive magnesium metal, *J. Am. Chem. Soc.* 94 (1972) 7178–7179.
- [70] J.-S. Lee, R. Velarde-Ortiz, A. Guijarro, J.R. Wurst, R.D. Rieke, Low-temperature formation of functionalized Grignard reagents from direct oxidative addition of active magnesium to aryl bromides, *J. Organomet. Chem.* 65 (17) (2000) 5428–5430.
- [71] J. Kvicala, J. Stambasky, M. Skalicky, O. Paleta, Preparation of fluorohalomethylmagnesium halides using highly active magnesium metal and their reactions, *J. Fluor. Chem.* 126 (9–10) (2005) 1390–1395.

- [72] H. Hazimeh, F. Kanoufi, C. Combellas, J.-M. Mattalia, C. Marchi-Delapierre, M. Chanon, Radical clocks, solvated electrons, and magnesium. Heterogeneous versus homogeneous electron transfer. Selectivity at interfaces, *J. Phys. Chem. C* 112 (7) (2008) 2545–2557.
- [73] B.I. Kharisov, U. Ortiz Méndez, L.A. Garza-Rodríguez, H.M. Leija Gutiérrez, A. Medina Medina, S.S. Berdonosov, Use of various activated forms of elemental nickel and copper for the synthesis of phthalocyanine at low temperature, *J. Coord. Chem.* 59 (15) (2006) 1657–1666.
- [74] E.A. Koerner von Gustorf, O. Jaenicke, O. Wolfbeis, C.R. Eady, The laser-evaporation of metals and its application to organometallic synthesis, *Angew. Chem.* 14 (5) (1975) 278–286.
- [75] T.M. Ayers, B.C. Westlake, D.V. Preda, L.T. Scott, M.A. Duncan, Laser plasma production of metal-corannulene ion-molecule complexes, *Organometallics* 24 (2005) 4573–4578.
- [76] S. Fedrigo, T.L. Haslett, M. Moskovits, Direct synthesis of metal cluster complexes by deposition of mass-selected clusters with ligand: iron with CO, *J. Am. Chem. Soc.* 118 (21) (1996) 5084–5085.
- [77] A. Krzton-Maziopa, E.V. Pomjakushina, V.Y. Pomjakushin, F. von Rohr, A. Schilling, K. Conder, Synthesis of a new alkali metal-organic solvent intercalated iron selenide superconductor with Tc 45 K, *J. Phys. Condens. Matter* 24 (38) (2012).
- [78] J. Dye, Electrides and alkalides—comparison with metal solutions, *J. Phys. IV Coll.* 01 (1991) C5. C5-259–C5-282.
- [79] T.M.M. Richter, R. Niewa, Chemistry of ammonothermal synthesis, *Inorganics* 2 (2014) 29–78.
- [80] Preparation method of anhydrous  $(\text{Ba}(\text{tmhd})_2)$ . Patent of China CN 201110061384, 2011.
- [81] J. Garcia, M.J. Allen, Developments in the coordination chemistry of Europium(II), *Eur. J. Inorg. Chem.* 2012 (29) (2012) 4550–4563.
- [82] A. Sawczenko, B. Miroslaw, T. Lis, A.E. Koziol, Direct synthesis of novel polymeric bimetallic phthalimide complex: poly{catena-bis( $\mu_3$ -phthalimido-N,O,O')-bis( $\mu_2$ -phthalimido-N,O)-aqua-dipropanol-copper(II)disodium}, *Pure Appl. Chem. Sci.* 2 (2) (2014) 73–86.
- [83] B.I. Kharisov, J. Rivera de la Rosa, O.V. Kharissova, U. Ortiz Mendez, M. Jose Yacaman, Nonsubstituted phthalocyanine nanostructures obtained using activated metals and unstable complexes at ambient temperature, *J. Compos. Mater.* 44 (17) (2010) 2145–2155.
- [84] F. Sánchez-De la Torre, J. Rivera De la Rosa, B.I. Kharisov, C.J. Lucio-Ortiz, Preparation and characterization of Cu and Ni on alumina supports and their use in the synthesis of low-temperature metal-phthalocyanine using a parallel-plate reactor, *Materials* 6 (2013) 4324–4344.

## FURTHER READING

- [1] E.V. Skorb, D. Fix, D.G. Shchukin, H. Mohwald, D.V. Sviridov, R. Mousa, N. Wanderka, J. Schaferhans, N. Pazos-Pérez, A. Fery, D.V. Andreeva, Sonochemical formation of metal sponges, *Nanoscale* 3 (2011) 985–993.