

## 8 Zoznam literatúry

- AFZELIUS A. Verhandlungen der dermatologischen gesellschaft zu Stockholm sitzung vom 28. Oktober 1909. Archiv fur Dermatologie and Syphilis 1910, 101: 404.
- AGUERO-ROSENFELD ME, WANG G, SCHWARTZ I, WORMSER GP. Diagnosis of lyme borreliosis. Clin Microbiol Rev. 2005;18(3):484-509. doi: 10.1128/CMR.18.3.484-509.2005.
- ANDERSON C, BRISSETTE CA. The brilliance of *Borrelia*: Mechanisms of host immune evasion by Lyme disease-causing spirochetes. Pathogens 2021, 2, 10 (3): 281. Doi: 10.3390/pathogens10030281.
- ANTOLOVÁ D, AVDIČOVÁ M, BEDRIOVÁ M, BELAY G, et al. Správa o zoonózach, alimentárnych nákazách a nákazách z vody v Slovenskej republike za rok 2019. Report on zoonosis, alimentary diseases and diseases from water in Slovak Republic in 2019. 1. vyd. Ministerstvo pôdohospodárstva a rozvoja vidieka Slovenskej republiky: Bratislava, 2020, 136 s. ISBN 978-80-972963-8-4
- ANTOLOVÁ D, AVDIČOVÁ M, BAKO J, et al. Správa o zoonózach, alimentárnych nákazách a nákazách z vody v Slovenskej republike za rok 2021. 1. vyd. Ministerstvo pôdohospodárstva a rozvoja vidieka SR: Bratislava 2022, 123s. ISBN 978-80-973917-5-1
- ANTONARA S, CHAFEL RM, LAFRANCE M, COBURN J. *Borrelia burgdorferi* adhesins identified using in vivo phage display. Mol Microbiol 2007, 66 (1): 262-276. Doi: 10.1111/j.1365-2958.2007.05924.x.
- ÅSBRINK E, HOVMARK A, HEDERSTEDT B. The spirochetal etiology of acrodermatitis chronica atropicans Herxheimer. Acta Dermato-Venereologica 1984, 64: 506- 512.
- BAKER PJ. A review of antibiotic-tolerant persisters and their relevance to posttreatment Lyme Disease symptoms. AJM 2020, 133 (4): 429-431. Doi: 10.1016/j.amjmed.2019.12.007
- BARANTON G, POSTIC D, SAINT GIRONS I, BOERLIN P, et al. Delineation of *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* sp. nov., and group VS461 associated with Lyme borreliosis. Int J Syst Bacteriol 1992, 42: 378-383. <https://doi.org/10.1099/00207713-42-3-378>
- BARBOUR AG. *Borreliaceae*. In: WHITMAN WB, RAINEY F, KÄMPFER P, TRUJILLO M, CHUN J, et al. (eds). Bergey's Manual of Systematics of Archaea and Bacteria. 2018.
- BARTŮŇEK P, GORIČAN K, KOCÍK M, MRÁZEK V, et al. Lyme carditis. Karolinum: Praha 2010. ISBN 978-80-246-1761-9.
- BARTŮŇEK P, BOJAR M, CALDA P, DIBLÍK P, et al. Lymeská borelióza. 4., přepracované a doplněné vydání. Grada Publ.: Praha 2013. ISBN 978-80-247-4355-4.
- BAUERFEIND R, SCHIEFER HG, SCHWARZ T, SLENCZKA W, et al. Zoonoses: Infectious diseases transmissible between animals and humans. 4th ed. 2016. Doi: 10.1128/9781555819262
- BECK G, HABICHT GS, BENACH JL, COLEMAN JL. Chemical and biological characterization of a lipopolysaccharide extracted from the Lyme disease spirochete, (*Borrelia burgdorferi*). J Infect Dis 1985, 152: 108-117. <https://www.jstor.org/stable/30104646>
- BHIDE MR, TRAVNICEK M, LEVKUTOVA M, CURLIK J, et al. Sensitivity of *Borrelia* genospecies to serum complement from different animals and human: a host-pathogen relationship. FEMS Immunology and Medical Microbiology 2005, 43, 165-172.
- BOCKENSTEDT LK, WOOTEN RM, BAUMGARTH N. Immune response to *Borrelia*: lessons from Lyme disease spirochetes. Curr Issues Mol Biol 2021, 42: 145-190. <https://doi.org/10.21775/cimb.042.145>
- BRANDA J, BODY BA, BOYLE J, BRANSON BM, DATTWYLER RJ, et al. Advances in serodiagnostic testing for Lyme disease are at hand. Clinical Infectious Diseases 2018, 66 (7): 1133-1139. Doi: 10.1093/cid/cix943
- BRANDA JA, Steere a kol.AC. Laboratory Diagnosis of Lyme Borreliosis. Clin Microbiol Rev 2021, 34 (2): e00018-19. Doi: 10.1128/CMR.00018-19.
- BRORSON O, BRORSON SH. An in vitro study of the activity of telithromycin against mobile and cystic forms of *Borrelia afzelii*. Infection 2006, 34: 26-28. Doi: 10.1007/s15010-006-4121-0



- CAMERON DJ, JOHNSON LB, MALONEY EL. Evidence assessments and guideline recommendations in Lyme disease: the clinical management of known tick bites, erythema migrans rashes and persistent disease. *Expert Review of Anti-infective Therapy* 2014, 12 (9): 1103-1135. Doi: 10.1586/14787210.2014.940900
- CARREIRO MM, LAUX DC, NELSON DR. Characterization of the heat shock response and identification of heat shock protein antigens of *Borrelia burgdorferi*. *Infect Immun* 1990, 58 (7): 2186-2191. Doi: 10.1128/iai.58.7.2186-2191.1990.
- CASJENS S, PALMER N, VAN VUGT R, HUANG WM, STEVENSON B, et al. A bacterial genome in flux: the twelve linear and nine circular extrachromosomal DNAs in an infectious isolate of the Lyme disease spirochete *Borrelia burgdorferi*. *Mol Microbiol* 2000, 35 (3): 490-516. Doi: 10.1046/j.1365-2958.2000.01698.x. PMID: 10672174.
- CASKEY JR, HASENKAMPF NR, MARTIN DS, CHOULJENKO VN, SUBRAMANIAN R, et al. The functional and molecular effects of doxycycline treatment on *Borrelia burgdorferi* phenotype. *Front Microbiol* 2019, 10: 690. Doi: 10.3389/fmicb.2019.00690
- COMSTEDT P, SCHÜLER W, MEINKE A, LUNDBERG U. The novel Lyme borreliosis vaccine VLA15 shows broad protection against *Borrelia* species expressing six different OspA serotypes. *PLoS One* 2017, 12 (9): e0184357. Doi: 10.1371/journal.pone.0184357.
- DEMAERSCHALCK I, BENMESSAOUD A, DE KESEL M, HOYOIS B, et al. Simultaneous presence of different *Borrelia burgdorferi* genospecies in biological fluids of Lyme disease patients. *J Clin Microbiol* 1995, 33: 602-608.
- DOU TLÍK S. In: BARTŮNĚK P: Lymeská borelióza. Grada, Avicenum: Praha 1996, s. 17-19.
- ECDC Comment: European Commission Updates Communicable Disease Surveillance List. Lyme Neuroborreliosis, EU/EEA Surveillance. 2018. <https://www.ecdc.europa.eu/en/news-events/ecdc-comment-european-commission-updates-communicable-disease-surveillance-list-lyme> [prístupné z r. 2022].
- EMBERS ME, NARASIMHAN S. Vaccination against Lyme disease: past, present, and future. *Front Cell Infect Microbiol* 2013, 12, 3: 6. Doi: 10.3389/fcimb.2013.00006.
- EPIS. Analýza epidemiologickej situácie v Slovenskej republike vrátane ochorenia COVID-19 za január 2023. 31. 1. 2023, dostupné online <https://www.epis.sk>
- FOGAÇA AC, SOUSA G, PAVANELO DB, ESTEVES E, et al. Tick immune system: what is known, the interconnections, the gaps, and the challenges. *Front Immunol* 2021, 12: 628054. Doi: 10.3389/fimmu.2021.628054
- FINNIAN R, CAUSLAND MC, BIJOL SV, RENNKE HG, et al. Lyme disease-associated glomerulonephritis. *Nephrology Dialysis Transplantation* 2011, 26 (9): 3054-3056. <https://doi.org/10.1093/ndt/gfr335>
- FISHER MA, GRIMM D, HENION AH, ELIAS AF, et al. *Borrelia burgdorferi*  $\sigma 54$  is required for mammalian infection and vector transmission but not for tick colonization. *Proc Natl Acad Sci USA* 2005, 102: 5162-5167. <https://doi.org/10.1073/pnas.0408536102>
- GEEBELEN L, LERNOUT T, DEVLEESSCHAUWER B, KABAMBA-MUKADI B, SAEGEMAN V, et al. Non-specific symptoms and post-treatment Lyme disease syndrome in patients with Lyme borreliosis: a prospective cohort study in Belgium (2016-2020). *BMC Infect Dis* 2022, 22 (1): 756. Doi: 10.1186/s12879-022-07686-8.
- GERN L, HUMAIR PF. Ecology of *Borrelia burgdorferi* sensu lato in Europe. In: Gray JS, Kahl O, Lane RS, Stanek G, (eds.). *Lyme Borreliosis: Biology, Epidemiology and Control*. CAB International: Wallingford 2002, s.141-174.
- GOMES-SOLECKI MJ, DUNN JJ, LUFT BJ, CASTILLO J, et al. Recombinant chimeric *Borrelia* proteins for diagnosis of Lyme disease. *J Clin Microbiol* 2000, 38: 2530-2535.
- GRUNTAR I, MALOVRH T, MURGIA R, CINCO M. Conversion of *Borrelia garinii* cystic forms to motile spirochetes in vivo. *APMIS* 2001, 109: 383-388. Doi: 10.1034/j.16000463.2001.090507.x.
- GÜRTLER L, BLÜMEL J, BURGER R, DROSTEN C, GRÖNER A, et al. Arbobacteria - pathogens transmittable by arthropods. *Transfus Med Hemother* 2008, 36 (1): 62-78. <https://doi.org/10.1159/000112812>



- HALOUZKA J, POSTIC D, HUBALEK Z. Isolation of the spirochaete *Borrelia afzelii* from the mosquito *Aedes vexans* in the Czech Republic. *Medical and Veterinary Entomology* 1998, 12: 103–105. Doi: 10.1046/j.1365-2915.1998.00086.x
- HANINCOVÁ K, TARAGELOVÁ V, KOCI J, SCHÄFER MR, et al. Association of *Borrelia garinii* and *B. valaisiana* with songbirds in Slovakia. *Appl Environ Microbiol* 2003, 69: 2825–2830.
- HOLEČKOVÁ K, KOŠČÁLOVÁ A, SCHRÉTER I. Lymeská borelióza. 2021, 16 s. Dostupné na: [https://health.gov.sk/30-6-2021/18\\_komisia\\_Lymska-borelioza-OP](https://health.gov.sk/30-6-2021/18_komisia_Lymska-borelioza-OP)
- HOVIS KM, TRAN E, SUNDY CM, BUCKLES E, et al. Selective binding of *Borrelia burgdorferi* OspE paralogs to factor H and serum proteins from diverse animals: possible expansion of the role of OspE in Lyme disease pathogenesis. *Infect Immun* 2006, 74 (3): 1967-1972. Doi: 10.1128/IAI.74.3.1967-1972.2006.
- HUBÁLEK Z, HALOUZKA J. Prevalence rates of *Borrelia burgdorferi* sensu lato in host-seeking *Ixodes ricinus* ticks in Europe. *Parasitol Res* 1998, 84 (3): 167-172. Doi: 10.1007/s004360050378.
- HULÍNSKÁ D. Mikrobiologie. S. 14-22. In: Bartůněk P. Lymeská borelióza. 3. vyd. Grada Publ.: Praha 2006.
- HUNFELD KP, BRADE V. Antimicrobial susceptibility of *Borrelia burgdorferi* sensu lato: what we know, what we don't know, and what we need to know. *Wien Klin Wochenschr* 2006, 118 (21-22): 659-668. Doi: 10.1007/s00508-006-0693-z.
- CHACONAS G, CASTELLANOS M, VERHEY TB. Changing of the guard: How the Lyme disease spirochete subverts the host immune response. *J Biol Chem* 2020, 295 (2): 301–313. Doi 10.1074/jbc.REV119.008583
- JAENSON TGT. The epidemiology of Lyme borreliosis. *Parasitol Today* 1991, 7: 39-45.
- JOVANOVIĆ D, ATANASIEVSKA S, PROTIC-DJOKIĆ V, RAKIĆ U, et al. Seroprevalence of *Borrelia burgdorferi* in occupationally exposed persons in the Belgrade area, Serbia. *Braz J Microbiol* 2015, 1, 46 (3): 807-814. Doi: 10.1590/S1517-838246320140698.
- JOVČIĆ VL, GREGO EM, LAKO BL, RISTOVIĆ BM, et al. Improved serodiagnosis of early Lyme borreliosis: Immunoblot with local *Borrelia afzelii* strain. *APMIS* 2003, 111: 1053-1059.
- KAMP HD, SWANSON KA, WEI RR, DHAL PK, et al. Design of a broadly reactive Lyme disease vaccine. *NPJ Vaccines* 2020, 1, 5 (1): 33. Doi: 10.1038/s41541-020-0183-8.
- KAPLAN RF, TREVINO RP, JOHNSON GM, LEVY L, et al. Cognitive function in post-treatment Lyme disease: do additional antibiotics help? *Neurology* 2003, 24, 60 (12): 1916-1922. Doi: 10.1212/01.wnl.0000068030.26992.25.
- KELLER A, GRAEFEN A, BALL M, et al. New insights into the Tyrolean Iceman's origin and phenotype as inferred by whole-genome sequencing. *Nat Commun* 2012, 3: 698. <https://doi.org/10.1038/ncomms1701>
- KENEDY MR, LENHART TR, AKINS DR. The role of *Borrelia burgdorferi* outer surface proteins. *FEMS Immunology & Medical Microbiology* 2012, 66 (1): 1–19. <https://doi.org/10.1111/j.1574-695X.2012.00980.x>
- KIM CM, YUN NR, KIM DM. Case Report: The First *Borrelia yangtzensis* infection in a human in Korea. *Am J Trop Med Hyg* 2022, 106 (1): 45-46. Doi: 10.4269/ajtmh.21-0052.
- KIŠOVÁ-VARGOVÁ L, ČERŇANSKÁ D, BĚHIDE M. Comparative study of binding of ovine complement factor H with different *Borrelia* genospecies. *Folia Microbiol* 2012, 57 (2): 123-128. Doi: 10.1007/s12223-012-0104-y.
- KMETY E, REHÁČEK J, VÝROSTEKOVÁ V. Investigations of ticks for the presence of *Borrelia* in Czechoslovakia. *Zentralbl Bakteriol Mikrobiol Hyg A* 1987, 263 (3): 468-470. Doi: 10.1016/s0176-6724(87)80111-6.
- KOŠŤANOVÁ Z, KOCIANOVÁ E, ŠPITALSKÁ E, SMETANOVÁ K, SCHWARZOVÁ K. Epidemiologické a epizootologické sledovanie nákaz prenášaných kliešťami v okrese Žiar nad Hronom v roku 2003. *Bulletin Československé spoločnosti mikrobiologickej* 2004, 45: 197. ISSN 0009-0646
- KRUPP LB, HYMAN LG, GRIMSON R, COYLE PK, et al. Study and treatment of post Lyme disease (STOP-LD): a randomized double masked clinical trial. *Neurology* 2003, 24, 60 (12): 1923-1930. Doi: 10.1212/01.wnl.0000071227.23769.9e.



- KUNG F, ANGUITA J, PAL U. *Borrelia burgdorferi* and tick proteins supporting pathogen persistence in the vector. *Future Microbiol* 2013, 8 (1): 41-56. Doi: 10.2217/fmb.12.121.
- LANTOS PM, RUMBAUGH J, BOCKENSTEDT LK, FALCK-YTTER IT, et al. Clinical Practice Guidelines by the Infectious Diseases Society of America (IDSA), American Academy of Neurology (AAN), and American College of Rheumatology (ACR): 2020 Guidelines for the Prevention, Diagnosis and Treatment of Lyme Disease, *Clinical Infectious Diseases* 2021, 72 (1): e1–e48, <https://doi.org/10.1093/cid/ciaa1215>
- LANTOS PM, AUWAERTER PG, WORMSER GP. A systematic review of *Borrelia burgdorferi* morphologic variants does not support a role in chronic Lyme Disease. *Clin Infect Dis* 2014, 58 (5): 663–671. <https://doi.org/10.1093/cid/cit810>
- LANTOS PM. Chronic Lyme disease. *Infect Dis Clin North Am* 2015, 29 (2): 325-340. Doi: 10.1016/j.idc.2015.02.006.
- LI X, NEELAKANTA G, LIU X, BECK DS, KANTOR FS, et al. Role of outer surface protein D in the *Borrelia burgdorferi* life cycle. *Infect Immun* 2007, 75 (9): 4237-4244. Doi: 10.1128/IAI.00632-07.
- LIMA CM, ZEIDNER NS, BEARD CB, SOARES CA, DOLAN MC, DIETRICH G, PIESMAN J. Differential infectivity of the Lyme disease spirochete *Borrelia burgdorferi* derived from *Ixodes scapularis* salivary glands and midgut. *J Med Entomol* 2005, 42 (3): 506–510. Doi: 10.1093/jmedent/42.3.506.
- LIU D, (ed.). *Molecular detection of human bacterial pathogens* (1st ed.). CRC Press 2011. <https://doi.org/10.1201/b10848>
- LJØSTAD U, MYGLAND Í. Chronic Lyme; diagnostic and therapeutic challenges. *Acta Neurol Scand Suppl* 2013, 196: 38-47. Doi: 10.1111/ane.12048.
- MAJLÁTHOVÁ V, MAJLÁTH I, DERDÁKOVÁ M, VÍCHOVÁ B, et al. *Borrelia lusitaniae* and green lizards (*Lacerta viridis*), Karst Region, Slovakia. *Emerg Infect Dis* 2006, 12 (12):1895-1901. Doi: 10.3201/eid1212.060784.
- MATLAHOVÁ K. Využitie biofyzikálnych a biochemických analýz v mikrobiologickej diagnostike Lymeskej boreliózy. Diplomová práca. Fakulta matematiky, fyziky a informatiky UK v Bratislave, 2015.
- MAKSIMYAN S, SYED MS, SOTI V. Post-treatment Lyme Disease syndrome: Need for diagnosis and treatment. *Cureus* 2021, 12, 13 (10): e18703. Doi: 10.7759/cureus.18703.
- MARGOS G, GOFTON A, WIBBERG D, DANGEL A, et al. The genus *Borrelia* reloaded. *PloS one* 2018, 13 (12): e0208432. <https://doi.org/10.1371/journal.pone.0208432>
- MEAD P, PETERSEN J, HINCKLEY A. Updated CDC Recommendation for Serologic Diagnosis of Lyme Disease. *MMWR Morb Mortal Wkly Rep* 2019, 68: 703. Doi: <http://dx.doi.org/10.15585/mmwr.mm6832a4external icon>
- MIDDELVEEN MJ, STRICKER RB. Morgellons disease: a filamentous borrelial dermatitis. *Int J Gen Med* 2016, 9: 349-354. Doi: 10.2147/IJGM.S116608.
- MIKLOSSY J, KASAS S, ZURN AD, MCCALL S, et al. Persisting atypical and cystic forms of *Borrelia burgdorferi* and local inflammation in Lyme neuroborreliosis. *J Neuroinflammation* 2008, 5: 40. Doi:10.1186/1742-2094-5-40
- MIKLOSSY J, KHALILI K, GERN L, ERICSON RL, et al. *Borrelia burgdorferi* persists in the brain in chronic lyme neuroborreliosis and may be associated with Alzheimer disease. *J Alzheimers Dis* 2004, 6 (6): 639-649. Discussion 673-681. Doi: 10.3233/jad-2004-6608.
- MONTGOMERY RR, LUSITANI D, DE BOISFLEURY CA, MALAWISTA SE. Tick saliva reduces adherence and area of human neutrophils. *Infect Immun* 2004, 72: 2989-2994.
- MURFIN KE, KLEINBARD R, AYDIN M, SALAZAR SA, FIKRIG E. *Borrelia burgdorferi* chemotaxis toward tick protein Salp12 contributes to acquisition. *Ticks Tick Borne Dis* 2019, 10 (5): 1124-1134. Doi: 10.1016/j.ttbdis.2019.06.002.
- NEELAKANTA G, LI X, PAL U, LIU X, BECK DS, et al. Outer surface protein B is critical for *Borrelia burgdorferi* adherence and survival within *Ixodes* ticks. *PLoS Pathog* 2007, 3 (3): e33. Doi: 10.1371/journal.ppat.0030033.
- NORDSTRAND A, BARBOUR AG, BERGSTRÖM S. *Borrelia* pathogenesis research in the post-genomic and post-vaccine era. *Curr Opin Microbiol* 2000, 3 (1): 86-92. Doi: 10.1016/S1369-5274(99)00056-9.



- NOVAK P, FELSENSTEIN D, MAO C, OCTAVIEN NR, et al. Association of small fiber neuropathy and post treatment Lyme disease syndrome. *PLoS One* 2019, 12, 14 (2): e0212222. Doi: 10.1371/journal.pone.0212222.
- NUTTALL P.A. Tick saliva and its role in pathogen transmission. *Wien Klin Wochenschr* 2023, 135: 165–176. <https://doi.org/10.1007/s00508-019-1500-y>
- OHNISHI J, PIESMAN J, DE SILVA AM. Antigenic and genetic heterogeneity of *Borrelia burgdorferi* populations transmitted by ticks. *Proc Natl Acad Sci USA* 2001, 16, 98 (2): 670-675. Doi: 10.1073/pnas.98.2.670.
- OLIVEIRA GMB, MUÑOZ-LEAL S, SANTODOMINGO A, WECK BC, et al. A Novel relapsing fever group *Borrelia* Isolated from *Ornithodoros* Ticks of the Brazilian Caatinga. *Microorganisms* 2023, 11 (2): 370. Doi: 10.3390/microorganisms11020370.
- OLSÉN B, JAENSON TG, NOPPA L, BUNIKIS J, BERGSTRÖM SA. Lyme borreliosis cycle in sea-birds and *Ixodes uriae* ticks. *Nature* 1993, 362 (6418): 340-342. Doi: 10.1038/362340a0.
- PAL U, FIKRIG E. Adaptation of *Borrelia burgdorferi* in the vector and vertebrate host. *Microbes Infect* 2003, 5 (7): 659-666. Doi: 10.1016/s1286-4579(03)00097
- PEŤKO B, ŠTEFANČÍKOVÁ A, TRESOVÁ G, PETERKOVÁ J, et al. Kliešť obyčajný (*Ixodes ricinus*) ako zdroj infekcie ľudí a psov pôvodcom lymeskej boreliózy. *Slovenský veterinársky časopis* 1996, 21: 313–319.
- PERRONNE C. Critical review of studies trying to evaluate the treatment of chronic Lyme disease. *Presse Med* 2015, 44 (7-8): 828-831. Doi: 10.1016/j.lpm.2015.06.002.
- POSTIC D, ASSOUS M, BELFAIZA J, BARANTON G. Genetic diversity of *Borrelia* of Lyme borreliosis. *Wien Klin Wochenschr* 1996, 108 (23): 748-751.
- POUPON MA, LOMMANO E, HUMAIR PF, DOUET V, et al. Prevalence of *Borrelia burgdorferi* sensu lato in ticks collected from migratory birds in Switzerland. *Appl Environ Microbiol* 2006, 72: 976-979. Doi: 10.1128/AEM.72.1.976-979.2006
- PURI BK, MONRO JA. Detection of 41-kDa bacterial flagellin protein by the lymphocyte transformation test-memory lymphocyte immunostimulation assay. *Am J Clin Exp Immunol* 2022, 15, 11 (4): 72-77.
- RAMESH G, ALVAREZ AL, ROBERTS ED, DENNIS VA, et al. Pathogenesis of Lyme neuroborreliosis: *Borrelia burgdorferi* lipoproteins induce both proliferation and apoptosis in rhesus monkey astrocytes. *Eur J Immunol* 2003, 33: 2539-2550.
- RUDENKO a kol., N, GOLOVCHENKO M, VANCOVA M, CLARK K, et al. Isolation of live *Borrelia burgdorferi* sensu lato spirochaetes from patients with undefined disorders and symptoms not typical for Lyme borreliosis. *Clin Microbiol Infect* 2016, 22 (3): 267. e9-15. Doi: 10.1016/j.cmi.2015.11.009.
- RUDENKO a kol., N, GOLOVCHENKO M, GRUBHOFFER L, OLIVER JH, Jr. Updates on *Borrelia burgdorferi* sensu lato complex with respect to public health. *Ticks Tick Borne Dis* 2011, 2 (3):123-128. Doi: 10.1016/j.ttbdis.2011.04.002.
- ROBERTS DM, THEISEN M, MARCONI RT. Analysis of the cellular localization of Bdr paralogs in *Borrelia burgdorferi*, a causative agent of lyme disease: evidence for functional diversity. *J Bacteriol* 2000, 182 (15): 4222-4226. Doi: 10.1128/JB.182.15.4222-4226.2000.
- SAPI E, BASTIAN SL, MPOY CM, SCOTT S, et al. Characterization of biofilm formation by *Borrelia burgdorferi* in vitro. *PLoS One* 2012, 7: e48277. Doi: 10.1371/journal.pone.0048277
- SHAPIRO ED. Clinical practice. Lyme disease. *N Engl J Med* 2014, 1, 370 (18): 1724-1731. Doi: 10.1056/NEJMcp1314325.
- SINGH SK, GIRSCHICK HJ. Lyme borreliosis: from infection to autoimmunity. *Clin Microbiol Infect* 2004, 10: 598-614. <https://doi.org/10.1111/j.1469-0691.2004.00895.x>
- SINGH SK, GIRSCHICK HJ. Toll-like receptors in *Borrelia burgdorferi*-induced inflammation. *Clin Microbiol Infect* 2006, 12 (8): 705-717. Doi: 10.1111/j.1469-0691.2006.01440.x.
- SOOD SK, SALZMAN MB, JOHNSON BJ, HAPP CM, et al. Duration of tick attachment as a predictor of the risk of Lyme disease in an area in which Lyme disease is endemic. *J Infect Dis* 1997, 175 (4): 996-999. Doi: 10.1086/514009.
- SCHWARZOVÁ K, SOBOŇOVÁ K. Lymeská borelióza. Diferenciálna diagnostika pacientov s nešpecifickými príznakmi. *Acta chemotherapeut* 2020, 29 (2): 17–25.



- SCHWARZOVA K, CIZNAR I, SVIHROVA V, HUDECKOVA H. Initial attachment of *Borrelia burgdorferi* spirochetes to Vero cells. Bratisl Lek Listy 2019, 120 (11): 872-875. Doi: 10.4149/BLL\_2019\_146.
- SCHWARZOVÁ K. Arbobaktérie, leptospiry, kapnocytofágy, streptobacily - baktérie s patogénnym potenciálom prenosné na človeka. Preveda: Bratislava 2019, 54 s. ISBN 978-80-972360-3-8.
- SCHWARZOVÁ K, KOZUB P, SZEP Z, GOLOVCHENKO M, Rudenko a kol., N. Detection of *Borrelia burgdorferi* sensu stricto and *Borrelia garinii* DNAs in patient with Hyperkeratosis lenticularis perstans (Flegel disease). Folia Microbiol 2016a, 61 (5): 359-363. Doi: 10.1007/s12223-016-0444-0.
- SCHWARZOVÁ K, WACZULÍKOVÁ I. Early localized and disseminated Lyme borreliosis in Western Slovakia. In: 5th Annual Scientific Congress on Zoonoses, Foodborne and Waterborne Diseases. Protection of Public and Animal Health [elektronický zdroj]. MPRV SR: Bratislava 2016b, 318-323 [CD-ROM]. ISBN 978-80-89738-09-0
- SCHWARZOVÁ K, BETÁKOVÁ T, NEMÉTH J, MIŽÁKOVÁ A. Detection of *Borrelia burgdorferi* sensu lato and *Chlamydomphila psittaci* in throat and cloacal swabs from birds migrating through Slovakia. Folia Microbiol 2006, 51 (6): 653-658. Doi: 10.1007/BF02931634.
- SCHWARZOVÁ K, KOŠŤANOVÁ Z, HOLEČKOVÁ K, ŠPITALSKÁ E, BOLDIŠ V. Direct detection of *Borrelia burgdorferi* spirochetes in patients with early disseminated Lyme borreliosis. Centr Eur J Public Health 2009, 17 (4): 179-182. Doi: 10.21101/cejph.b0015
- SCHWARZOVÁ K, ČIŽNÁR I. Immunochemical analysis of lipopolysaccharide-like component extracted from *Borrelia burgdorferi* sensu lato. Folia Microbiol 2004, 49 (5): 625-629. Doi: 10.1007/BF02931545
- SKAAR EP. The battle for iron between bacterial pathogens and their vertebrate hosts. PLoS Pathogens 2010, 6: e1000949. Doi: 10.1371/journal.ppat.1000949.
- SMETANOVÁ K, SCHWARZOVÁ K, KOCIANOVÁ E. Detection of *Anaplasma phagocytophilum*, *Coxiella burnetii*, *Rickettsia* spp., and *Borrelia burgdorferi* s.l. in ticks, and wild-living animals in western and middle Slovakia. Ann NY Acad Sci 2006, 1078: 312-315. Doi: 10.1196/annals.1374.058
- SMITH K. Lyme borreliosis research: An overview of Lyme and direction for further research required in Australia. 2012, p. 10 (www.lymeaustralia.com). ISBN: 978-0-646-90428-3.
- SPARAGANO OAE. Review of „*Borrelia*: Molecular Biology, Host interaction and Pathogenesis“, by DS Samuels and JD Radolf. Parasit Vectors 2010, 17, 3: 52. Doi: 10.1186/1756-3305-3-52.
- STANEK G, FINGERLE V, HUNFELD KP, JAULHAC B, et al. Lyme borreliosis: clinical case definitions for diagnosis and management in Europe. Clin Microbiol Infect 2011, 17 (1): 69-79. Doi: 10.1111/j.1469-0691.2010.03175.x.
- STEINBRINK A, BRUGGER K, MARGOS G, et al. The evolving story of *Borrelia burgdorferi* sensu lato transmission in Europe. Parasitol Res 2022, 121: 781-803. <https://doi.org/10.1007/s00436-022-07445-3>
- Steere a kol.AC, STRLE F, WORMSER GP, HU LT, et al. Lyme borreliosis. Nat Rev Dis Primers 2016, 2, 16090. <https://doi.org/10.1038/nrdp.2016.90>
- STRLE F, RUŽIC-SABLJIC E, CIMPERMAN J, LOTRIČ-FURLAN S, MARASPIN V. Comparison of findings for patients with *Borrelia garinii* and *Borrelia afzelii* isolated from cerebrospinal fluid. Clin Infect Dis 2006, 15, 43 (6): 704-710. Doi: 10.1086/506936
- ŠPITÁLSKA E, SPARAGANO O, STANKO M, SCHWARZOVÁ K, et al. Diversity of *Coxiella*-like and *Francisella*-like endosymbionts, and *Rickettsia* spp., *Coxiella burnetii* as pathogens in the tick populations of Slovakia, Central Europe. Ticks Tick-Borne Dis 2018, 9: 1207-1211. <https://doi.org/10.1016/j.ttbdis.2018.05.002>
- TAKAYAMA K, ROTHENBERG RJ, BARBOUR AG. Absence of lipopolysaccharide in the Lyme disease spirochete, *Borrelia burgdorferi*. Infect Immun 1987, 55: 2311-2313. Doi: 10.1128/iai.55.9.2311-2313.1987
- TÄLLEKLINT L, JAENSON TG. Transmission of *Borrelia burgdorferi* s.l. from mammal reservoirs to the primary vector of Lyme borreliosis, *Ixodes ricinus* (Acari: Ixodidae), in Sweden. Journal of Medical Entomology 1994, 31: 880-886.



- TAN X, CASTELLANOS M, CHACONAS G. Choreography of Lyme disease spirochete adhesins to promote vascular escape. *Microbiol Spectr* 2023: e0125423. Doi: 10.1128/spectrum.01254-23.
- TAN X, LIN YP, PEREIRA MJ, CASTELLANOS M, HAHN BL, et al. VlsE, the nexus for antigenic variation of the Lyme disease spirochete, also mediates early bacterial attachment to the host microvasculature under shear force. *PLoS Pathog* 2022, 23, 18 (5): e1010511. Doi: 10.1371/journal.ppat.1010511.
- TARAGEĽOVÁ V, KOČI J, HANINCOVÁ K, OLEKŠÁK M. Songbirds as hosts of ticks (*Acari: Ixodidae*) in Slovakia. *Biologia* 2005, 60: 529-537.
- THOMAS V, ANGUITA J, SAMANTA S, ROSA PA, et al. Dissociation of infectivity and pathogenicity in *Borrelia burgdorferi*. *Infect Immun* 2001, 69: 3507-3509. Doi: 10.1128/IAI.69.5.3507-3509.2001.
- TILLY K, ROSA PA, STEWART PE. Biology of infection with *Borrelia burgdorferi*. *Infect Dis Clin North Am* 2008, 22 (2): 217-234. <https://doi.org/10.1016/j.idc.2007.12.013>.
- TILLY K, KRUM JG, BESTOR A, JEWETT MW, et al. *Borrelia burgdorferi* OspC protein required exclusively in a crucial early stage of mammalian infection. *Infect Immun* 2006, 74 (6): 3554-3564. Doi: 10.1128/IAI.01950-05.
- TICHA L, GOLOVCHENKO M, OLIVER, JH, Jr, GRUBHOFFER L, Rudenko a kol., N. Sensitivity of Lyme borreliosis spirochetes to serum complement of regular Zoo animals: Potential reservoir competence of some exotic vertebrates. *Vector Borne Zoonotic Dis* 2016, 16 (1): 13-19. Doi: 10.1089/vbz.2015.1847.
- TREVISAN G, BONIN S, RUSCIO MA. Practical approach to the diagnosis of Lyme borreliosis: from clinical heterogeneity to laboratory methods. *Front Med* 2020, 7: 265. Doi: 10.3389/fmed.2020.00265
- TREVISAN G, CINCO M, TREVISINI S, DI MEO N, CHERSI K, et al. *Borreliae* Part 1: *Borrelia* Lyme Group and Echidna-Reptile Group. *Biology* 2021, 10 (10): 1036. <https://doi.org/10.3390/biology10101036>
- TREVISAN G, CINCO M, TREVISINI S, DI MEO N, RUSCIO M, et al. *Borreliae* Part 2: *Borrelia* Relapsing Fever Group and Unclassified *Borrelia*. *Biology (Basel)* 2021, 29, 10 (11): 1117. Doi: 10.3390/biology10111117.
- TREVISAN G, RUSCIO M, DI MEO N, NAN K, et al. Case Report: Lyme Borreliosis and Pregnancy - Our Experience. *Front Med* 2022, 28, 9: 816868. Doi: 10.3389/fmed.2022.816868.
- VERHAEGH D, JOOSTEN LAB, OOSTING M. The role of host immune cells and *Borrelia burgdorferi* antigens in the etiology of Lyme disease. *Eur Cytokine Netw* 2017, 28 (2): 70-84. Doi:10.1684/ecn.2017.0396.
- VERHEY TB, CASTELLANOS M, CHACONAS Ch. Antigenic variation in the Lyme spirochete: insights into recombinational switching with a suggested role for error-prone repair. *Cell Reports* 2018, 23: 2595-2605. <https://doi.org/10.1016/j.celrep.2018.04.117>
- VESTNÍK MZ SR. Odborné usmernenie MZ SR na diagnostiku a liečbu lymfkej choroby v SR. Výnos č. Z52700-2013-OZS. Čiastka 50 - 60. 2013, s. 297 - 314.
- VON LACKUM K, STEVENSON B. Carbohydrate utilization by the Lyme borreliosis spirochete, *Borrelia burgdorferi*. *FEMS Microbiology Letters* 2005, 243: 173-179. Doi: 10.1016/j.femsle.2004.12.002.
- WACZULÍKOVÁ I, MIŠENKO P, MATLAHOVÁ K, SCHWARZOVÁ K. Pros and cons of Lyme disease tests: an agreement analysis. 5th Annual Scientific Congress on Zoonoses, Foodborne and Waterborne Diseases - Protection of Public and Animal Health [elektronický zdroj]. MPRV SR: Bratislava 2016, s. 318-323 [CD-ROM]. ISBN 978-80-89738-09-0
- WALLICH R, MOTER SE, SIMON MM, EBNET K, et al. The *Borrelia burgdorferi* flagellum-associated 41-kilodalton antigen (flagellin): molecular cloning, expression, and amplification of the gene. *Infect Immun* 1990, 58 (6): 1711-1719. Doi: 10.1128/iai.58.6.1711-1719.1990.
- WILSKE B, FINGERLE V, SCHULTE-SPECHTEL U. Microbiological and serological diagnosis of Lyme borreliosis. *FEMS Immunology and Medical Microbiology* 2007, 49 (1): 13-21. <https://doi.org/10.1111/j.1574-695X.2006.00139.x>



WORLD HEALTH ORGANIZATION. Regional office for South-East Asia. 2014. A brief guide to emerging infectious diseases and zoonoses. <https://apps.who.int/iris/handle/10665/204722>

WORLD HEALTH ORGANIZATION. Annex: Regional Classifications 2022. <https://www.who.int/observatories/global-observatory-on-health-research-and-development/classifications-and-standards/country-groupings>.

WORLD HEALTH STATISTICS. Monitoring health for the SDGs. Geneva: WHO 2023. <https://www.who.int/publications/i/item/9789240074323>. Z