
References

- Aaronson NK, Ahmedzai S, Bergman B, et al. (1993) The European Organization for Research and Treatment of Cancer QLQ-30: A quality-of-life instrument for use in international clinical trials in oncology. *Journal of the National Cancer Institute*, 85: 365-376.
- Albert PS. (1999) Tutorial in biostatistics: Longitudinal data analysis (repeated measures) in clinical trials. *Statistics in Medicine*, 18: 1707-1732.
- Bacik J, Mazumdar M, Murphy BA, Fairclough DL, Eremenco S, Mariani T, Motzer RJ, Cella D. (2004) The functional assessment of cancer therapy-BRM (FACT-BRM): A new tool for the assessment of quality of life in patients treated with biologic response modifiers. *Quality of Life Research*, 13: 137-154.
- Barnard J, Meng XL. (1999) Applications of multiple imputation in medical studies: From AIDS to NHANES. *Statistical Methods in Medical Research*, 8: 17-36.
- Barnard J, Rubin DB. (1999) Small-sample degrees of freedom with multiple imputation. *Biometrika*, 86: 948-955.
- Baron RM, Kenny DA. (1986) The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistic considerations. *Journal Personality and Social Psychology*, 51: 1173-1182.
- Bernhard J, Peterson HF, Coates AS, Gusset H, Isley M, Hinkle R, Gelber RD, Castiglione-Gertsch M, Hürny C. (1998) Quality of life assessment in International Breast Cancer Study Group (IBCSG) trials: Practical issues and factors associated with missing data. *Statistics in Medicine*, 17: 587-602.
- Bernhard J, Cella DF, Coates AS, et al. (1998) Missing quality of life data in clinical trials: Serious problems and challenges. *Statistics in Medicine*, 17: 517-532.
- Bonomi P, Kim KM, Fairclough D, Cella D, Kugler J, Rowinsky E, Jiroutek M, Johnson D. (2000) Comparison of survival and quality of life in advanced non-small-cell lung cancer patients treated with two dose levels of Paclitaxel combined with Cisplatin versus Etoposide-Cisplatin: Results of an Eastern Cooperative Group Trial. *Journal of Clinical Oncology*, 18: 623-31.

- Bouchet C, Guillemin F, Paul-Dauphin A, Brianon S. (2000) Selection of quality-of-life measures for a prevention trial: A psychometric analysis. *Controlled Clinical Trials*, 21: 30-43.
- Box GEP, Draper NR. (1987). *Empirical Model-Building and Response Surfaces*, p. 424, Wiley, New York.
- Brazier JE, Roberts J, Deverill M. (2002) The estimation of a preference-based measure of health status from the SF-36. *Journal of Health Economics*, 21: 271-292.
- Brooks R. (1996) EuroQoL: The current state of play. *Health Policy (Amsterdam)*, 37: 53-72.
- Brooks MM, Jenkins LS, Schron EB, Steinberg JS, Cross JA, Paeth DS. (1998) Quality of life at baseline: Is assessment after randomization valid? *Medical Care*, 36: 1515-1519.
- Brown CH. (1990) Protecting against nonrandomly missing data in longitudinal studies. *Biometrics*, 46: 143-155.
- Buck SF. (1960) A method of estimation of missing values in multivariate data suitable for use with a electronic computer. *Journal Royal Statistical Society, Series B*, 22: 302-306.
- Busch P, Schwendener P, Leu RE, VonDach B, Castiglione M. (1994) Life quality assessment of breast cancer patients receiving adjuvant therapy using incomplete data. *Health Economics*, 3: 203-220.
- Cella DF, Skeel RT, Bonomi AE. (1993) Policies and Procedures Manual. Eastern Cooperative Oncology Group Quality of Life Subcommittee (unpublished).
- Cella DF, Tulsky DS, Gray G, et al. (1993) The Functional Assessment of Cancer Therapy (FACT) scales: Development and validation of the general measure. *Journal of Clinical Oncology*, 11: 570-579.
- Cella DF, Bonomi AE, Lloyd SR, Tulsky DS, Kaplan E, Bonomi P. (1995) Reliability and validity of the Functional Assessment of Cancer therapy - Lung (FACT-L) quality of life instrument. *Lung Cancer*, 12: 199-220, 1995.
- Cella DF, Bonomi AE. (1995) Measuring quality of life: 1995 update. *Oncology*, 9 11(supplement) 47-60.
- Cella DF. (2001) Personal communication.
- Center for Drug Evaluation and Reserach, Food and Drug Administration. (1998) E9 Statistical principles for clinical trials. *Federal Register*, 63: 49584-49598.
- Chi Y, Ibrahim JG. (2006) Joint models for multivariate longitudinal and

- multivariate survival data. *Biometrics*, 62, 432-445.
- Chuang-Stein C, Tong DM. (1996) The impact of parameterization on the interpretation of the main-effect term in the presence of an interaction. *Drug Information Journal*, 30: 421-424.
- Cleeland CS, Mendoza TR, Wang XS, Chou C, Harle MT, Morrissey M, Engstrom MC. (2000) Assessing symptoms distress in cancer patients: The MD Anderson Symptom Inventory. *Cancer*, 89: 1634-1646.
- Cohen J. (1988) *Statistical Power Analysis for the Behavioral Sciences*, 2nd edition. Lawrence Erlbaum Associates, Hillsdale, New Jersey.
- Cole BF, Gelber RD, Goldhirsch A. (1993) Cox regression models for quality adjusted survival analysis. *Statistics in Medicine* 12: 975-987.
- Converse JM, Presser S. (1986) *Survey Questions: Handcrafting the Standardized Questionnaire*. *Survey Research Methods*. Sage Publications, Newbury Park, CA.
- Cook NR. (1997) An imputation method for non-ignorable missing data in studies of blood pressure. *Statistics in Medicine*, 16: 2713-2728.
- Cox DR, Fitzpatrick R, Fletcher AI, Gore SM, Spiegelhalter DJ, Jones DR. (1992) Quality-of-life assessment: Can we keep it simple? (with discussion). *Journal of the Royal Statistical Society A*, 155: 353-393.
- Crawford SL, Tennstedt SL, McKinlay JB. (1995) A comparison of analytic methods for non-random missingness of outcome data. *Journal of Clinical Epidemiology*, 48: 209-219.
- Curren D. (2000) *Analysis of Incomplete Longitudinal Quality of Life Data*. Doctoral dissertation, Linburgs Universitair Centrum.
- Curren D, Aaronson N, Standaert B, Molenberghs G, Therasse P, Ramirez A, Keepmanschap M, Erder H, Piccart M. (2000) Summary measures and statistics in the analysis of quality of life data: an example from an EORTC-NCIC-SAKK locally advanced breast cancer study. *European Journal of Cancer*, 36: 834-844.
- Daniels MJ, Hogan JW. (2000) Reparameterizing the pattern mixture model for sensitivity analyses under informative dropout. *Biometrics*, 5: 1241-1248.
- Dawson JD, Lagakos SW. (1991) Analyzing laboratory marker changes in AIDS clinical trials. *Journal of Acquired Immune Deficiency Syndromes*, 4: 667-676.
- Dawson JD, Lagakos SW. (1993) Size and power of two-sample tests of repeated measures data. *Biometrics*, 49: 1022-1032.

- Dawson JD. (1994) Stratification of summary statistic tests according to missing data patterns. *Statistics in Medicine*, 13: 1853-1863.
- DeGruttola V, Tu XM. (1994) Modeling progression of CD-4 lymphocyte count and its relationship to survival time. *Biometrics*, 50: 1003-1014.
- DeKlerk NH. (1986) Repeated warnings re repeated measures. *Australian New Zealand Journal of Medicine*, 16: 637-638.
- Dempster AP, Laird NM, Rubin DB. (1977) Maximum likelihood estimation from incomplete data via the EM algorithm (with discussion). *Journal of the Royal Statistical Society, Series B*, 39: 1-38.
- Devellis RF. (1991) *Scale Development: Theory and Application*. Sage Publications, Newbury Park, CA.
- Diehr P, Patrick DL, Hedrick S, Rothaman M, Grembowski D, Raghunathan TI, Beresford S. (1995) Including deaths when measuring health status over time. *Medical Care*, 33(suppl): AS164-172.
- Diggle PJ, Kenward MG. (1994) Informative dropout in longitudinal data analysis (with discussion). *Applied Statistics*, 43: 49-93.
- Diggle PJ, Liang KY, Zeger SL. (1994) *Analysis of Longitudinal Data*. Oxford Science Publications. Clarendon Press, Oxford.
- Dmitrienko A, Offen W, Westfall PH. (2003) Gatekeeping strategies for clinical trials that do not require all primary effects to be significant. *Statistics in Medicine*, 22: 2387-2400.
- Donaldson GW, Nakamura Y, Moinpour C. (2009) Mediators, moderators, and modulators of causal effects in clinical trials - Dynamically Modified Outcomes (DYNAMO) in health-related quality of life. *Quality of Life Research*, 18: 137-145.
- Drummond M. (2001) Introducing economic and quality of life measurements into clinical studies. *Annals of Medicine*, 33: 344-349.
- Elashoff RM, Li G, Li N. (2007) An approach to joint analysis of longitudinal measurements and competing risks failure time data. *Statistics in Medicine*, 26: 2813-35.
- Fairclough DL, Cella DF. (1996) A cooperative group report on quality of life research: Lessons learned. Eastern Cooperative Oncology Group (ECOG). *Journal of the National Cancer Institute*, 40: 73-75.
- Fairclough DL, Cella DF. (1996) Functional Assessment of Cancer Therapy (FACT-G): Non-response to individual questions. *Quality of Life Research*, 5: 321-329.
- Fairclough DL. (1997) Summary measures and statistics for comparison of

- quality of life in a clinical trial of cancer therapy, *Statistics in Medicine*, 16: 1197-1209.
- Fairclough DL. (1998) Method of analysis for longitudinal studies of health-related quality of life, Chapter 13. *Quality of Life Assessment in Clinical Trials: Methods and Practice*. Staquet MJ, Hayes RD and Fayers PM, eds. Oxford University Press, New York.
- Fairclough DL, Peterson H, Chang V. (1998) Why is missing quality of life data a problem in clinical trials of cancer therapy? *Statistics in Medicine*, 17: 667-678.
- Fairclough DL, Peterson H, Bonomi P, Cella DF. (1998) Comparison of model based methods dependent of the missing data mechanism in two clinical trials of cancer therapy. *Statistics in Medicine*, 17: 781-796.
- Fairclough DL, Fetting JH, Cella D, Wonson W, Moinpour C for the Eastern Cooperative Oncology Group. (1999) Quality of life and quality adjusted survival for breast cancer patients receiving adjuvant therapy. *Quality of Life Research*, 8: 723-731.
- Farivar SS, Cunningham WE, Hayes RD. (2007) Correlated physical and mental health summary scores for the SF-36 and SF-12 Health Survey, V.1. *Health and Quality of Life Outcomes*, 5: 54-61.
- Fayers PM, Hand DJ. (1997) Factor analysis, causal indicators and quality of life. *Quality of Life Research*, 6: 139-150.
- Fayers PM, Machin D. (2000) *Quality of Life: Assessment, Analysis and Interpretation*. John Wiley and Sons, England.
- Feeny D, Furlong W, Barr RD, et al. (1992) A comprehensive multi-attribute system for classifying the health status of survivors of childhood cancer. *Journal of Clinical Oncology*, 10: 923-928.
- Feeny D. (2005) Preference-based measures: Utility and quality-adjusted life years. In *Assessing quality of life in Clinical Trials, 2nd edition*. pp 405-429. Fayers P and Hays R, eds. Oxford University Press.
- Feeny D, Farris K, Côté I, Johnson JA, Tsuyuki RT, Eng K. (2005) A cohort study found the RAND-12 and Health Utilities Index Mark 3 demonstrated construct validity in high-risk primary care patients. *Journal of Clinical Epidemiology*, 58: 138-41.
- Fetting J, Gray R, Abeloff M, et al. (1995) CAF vs. a 16-week multidrug regimen as adjuvant therapy for receptor-negative, node positive breast cancer: An intergroup study. *Proceedings of the American Society of Clinical Oncology*, 114: #83.
- Fetting JJ, Gray R, Abeloff MD, Fairclough DL, et al. (1998) A 16-week

- multidrug regimen versus cyclophosphamide, doxorubicin and 5-fluorouracil as adjuvant therapy for node-positive, receptor negative breast cancer: An intergroup study. *Journal of Clinical Oncology*, 16: 2382-91.
- Fitzmaurice GM, Laird NM, Shneyer L. (2001) An alternative parameterization of the general linear mixture model for longitudinal data with non-ignorable drop-outs. *Statistics in Medicine*, 20: 1009-1021.
- Fleiss JL. (1986) *The Design and Analysis of Clinical Experiments*. John Wiley and Sons, New York.
- Floyd J, Fowler JR. (1988) *Survey Research Methods*. Sage Publications, Newbury Park, CA.
- Follman D, Wu M. (1995) An approximate generalized linear model with random effects for informative missing data, *Biometrics*, 51: 151-168.
- Frank-Stromborg M, Olsen S. (1997) *Instruments for Clinical Health Care Research, Second Edition* Jones and Bartlett Publishers, Boston.
- Franks P, Lubetkin EI, Gold MR, Tancredi DJ. (2003) Mapping the SF-12 to preference-based instruments: Convergent validity in a low-income, minority population. *Medical Care*, 41: 1277-83.
- Freemantle N, Calvert M, Wood J, Eastangh J, Griffin C. (2003) Composite outcomes in randomized trials: Great precision but with greater uncertainty? *Journal of the American Medical Association*, 239: 2554-59.
- Friedman LM, Furberg CD, Demets D. (1985) *Fundamentals of Clinical Trials, 2nd edition*. John Wright PSG, Boston.
- Frison L, Pocock SJ. (1992) Repeated measures in clinical trials: Analysis using mean summary statistics and its implications for design. *Statistics in Medicine* 11: 1685-1704.
- Gelber RD, Gelman RS, Goldhirsh A. (1998) A quality of life oriented endpoint for comparing therapies. *Biometrics* 45: 781-95.
- Glasziou PP, Simes RJ, Gelber RD. (1990) Quality adjusted survival analysis. *Statistics in Medicine* 9: 1259-1276.
- Glynn RJ, Laird NM, Rubin DB. (1993) Multiple imputation in mixture models for non-ignorable nonresponse with follow-ups. *Journal of the American Statistical Association*, 88: 984-993.
- Goldhirsch A, Gelber RD, Simes RJ, et al. (1989) Costs and benefits of adjuvant therapy in breast cancer: A quality adjusted survival analysis. *Journal of Clinical Oncology* 7: 36-44, 1989.
- Gotay CC, Korn EL, McCabe MS, Moore TD, Cheson BD. (1992) Building quality of life assessment into cancer treatment studies. *Oncology*, 6: 25-28.

- Gould AL. (1980) A new approach to the analysis of clinical drug trails with withdrawals. *Biometrics*, 36: 721-727.
- Guo X, Carlin BP. (2004) Separate and joint modeling of longitudinal and event time data using standard computer packages. *The American Statistician*, 16-24.
- Guyatt GH, Townsend M, Merman LB, Keller JL. (1987) A comparison of Likert and visual analogue scales for measuring change in function. *Journal of Chronic Disease*, 40: 1129-1133.
- Guyatt G et al. (1991) Glossary. *Controlled Clinical Trials*, 12: 274S-280S.
- Heitjan DA, Landis JR. (1994) Assessing secular trends in blood pressure: A multiple imputation approach. *Journal of the American Statistical Association*, 89: 750-759.
- Heyting A, Tolbomm TBM, Essers JGA. (1992) Statistical handling of dropouts in longitudinal clinical trials. *Statistics in Medicine*, 11: 2043-2061.
- Hicks JE, Lampert MH, Gerber LH, Glastein E, Danoff J. (1985) Functional outcome update in patients with soft tissue sarcoma undergoing wide local excision and radiation (Abstract). *Archives of Physical Medicine and Rehabilitation*, 66: 542-543.
- Hochberg Y. (1988) A sharper Bonferroni procedure for multiple significance testing. *Biometrika*, 75: 800-803.
- Hogan JW, Laird NM. (1997) Mixture models for the joint distribution of repeated measures and event times. *Statistics in Medicine*, 16: 239-257.
- Hogan JW, Laird NM. (1997) Model-based approaches to analysing incomplete longitudinal and failure time data. *Statistics in Medicine*, 16: 259-272.
- Hogan JW, Roy J, Korkontzelou C. (2004) Tutorial in biostatistics: Handling drop-out in longitudinal studies. *Statistics in Medicine*, 23: 1455-1497.
- Hogan JW, Lin X, Herman G. (2004) Mixtures of varying coefficient models for longitudinal data with discrete or continuous non-ignorable dropout. *Biometrics*, 60: 854-864.
- Hollen PJ, Gralla RJ, Cox C, Eberly SW, Kris M. (1997) A dilemma in analysis: Issues in serial measurement of quality of life in patients with advanced lung cancer. *Lung Cancer*, 18: 119-136.
- Holm S. (1979) A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6: 65-70.
- Holmbeck GN. (1997) Toward terminological, conceptual and statistical clarity in the study of mediators and moderators: Examples from the child-

- clinical and pediatric psychology literatures. *Journal of Consulting and Clinical Psychology*, 65: 599-610.
- Holmbeck GN. (2002) Post-hoc probing of significant moderational and mediational effects in studies of pediatric populations. *Journal of Pediatric Psychology*, 27: 87-96.
- Hommel G. (1983) Tests of overall hypothesis for arbitrary dependence structures. *Biometric Journal*, 25: 423-30.
- Hommel G, Bretz F, Maurer W. (2007) Powerful sort-cuts for multiple testing procedures with special reference to gatekeeping strategies. *Statistics in Medicine*, 26: 4063-4073.
- Hopwood P, Harvey A, Davies J, Stephens RJ, et al. (1997) Survey of the administration of quality of life questionnaires in three multicentre randomised trials in cancer. *European Journal of Cancer*, 90: 49-57.
- Hürny C, Bernhard J, Gelber RD, Coates A, Castiglione M, Isley M, Dreher D, Peterson H, Goldhirsch A, Senn HJ. (1992) Quality of life measures for patients receiving adjuvant therapy for breast cancer: An international trial. *European Journal of Cancer*, 28: 118-124.
- Hürny C, Bernhard J, Bacchi M, et al. (1993) The Perceived Adjustment to Chronic Illness Scale (PACIS): A global indicator of coping for operable breast cancer patients in clinical trials. *Supportive Care in Cancer*, 1: 200-208.
- Hürny C, Bernhard J, Coates AS, et al. (1996) Impact of adjuvant therapy on quality of life in node-positive patients with operable breast cancer. *Lancet*, 347: 1279-1284.
- International Committee of Medical Journal Editors (2008) Uniform requirement for manuscripts submitted to biomedical journals. Retrieved from: www.jcmje.org
- Jaros, M. (2008) *A Joint Model for Longitudinal Data and Competing Risks*. Doctoral dissertation, University of Colorado at Denver.
- Jennrich R, Schluchter M. (1986) Unbalanced repeated-measures models with structured covariance matrices. *Biometrics*, 42: 805-820.
- Jhingran P, Osterhaus JT, Miller DW, Lee JT, Kirchdoerfer L. (1998) Development and validation of the migraine-specific quality of life questionnaire. *Headache*, 38: 295-302.
- Jones RH. (1993) *Longitudinal Data with Serial Correlation: A State-Space Approach*. Chapman and Hall, London.
- Juniper EF, Guyatt GH, Jaeschke R. (1996) How to develop and validate a new health-related quality of life instrument. in *Quality of Life and Pharma-*

- coeconomics in Clinical Trials*, Chapter 6. Spilker B, ed. Lippincott-Raven Publishers, Philadelphia.
- Kaplan RM, Bush JW. (1982) Health-related quality of life measurement for evaluation research and policy analysis. *Health Psychology*, 1: 61-80.
- Kenward MG, Roger JH. (1997) Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics*, 53: 983-997.
- Kittleson JM, Sharples K, Emerson SS. (2005) Group sequential clinical trials for longitudinal data with analyses using summary statistics. *Statistics in Medicine*, 24: 2457-2475.
- Korn EL, O'Fallon J. (1990) *Statistical Considerations, Statistics Working Group. Quality of Life Assessment in Cancer Clinical Trials*, Report on Workshop on Quality of Life Research in Cancer Clinical Trials. Division of Cancer Prevention and Control, National Cancer Institute.
- Korn DL. (1993) On estimating the distribution function for quality of life in cancer clinical trials. *Biometrika*, 80: 535-42.
- Kosinski M, Bayliss M, Bjorner JB, Ware JE. (2000) Improving estimates of SF-36 Health Survey scores for respondents with missing data. *Medical Outcomes Trust Monitor*, 5: 8-10.
- Kraemer HC, Wilson GT, Fairburn CG, Agras WS. (2002) Mediators and moderators of treatment effects in randomized clinical trials. *Archives of General Psychiatry*, 59: 877-883.
- Laird NM, Ware JH. (1982) Random-effects models for longitudinal data. *Biometrics*, 38: 963-974.
- Lauter J. (1996) Exact t and F tests for analyzing studies with multiple endpoints. *Biometrics*, 52: 964-970.
- Lavori PW, Dawson R, Shera D. (1995) A multiple imputation strategy for clinical trials with truncation of patient data. *Statistics in Medicine*, 14: 1912-1925.
- Law NJ, Taylor JMG, Sandler HM. (2002) The joint modeling of a longitudinal disease progression marker and the failure time process in the presence of cure. *Biostatistics*, 3: 547-563.
- Lechmaher W, Wassmer G, Reitmeir P. (1991) Procedures for two-sample comparisons with multiple endpoints controlling the experimentwise error rate. *Biometrics*, 47: 511-522.
- Lee CW, Chi KN. (2000) The standard of report of health-related quality of life in clinical cancer trials. *Journal of Clinical Epidemiology*, 53: 451-458.
- Levine M, Guyatt G, Gent M, et al. (1988) Quality of life in stage II breast

- cancer: An instrument for clinical trials. *Journal of Clinical Oncology*, 6: 1798-1810.
- Li L, Hu B, Greene T (2007) Semiparametric joint modeling of longitudinal and survival data. *Joint Statistical Meetings Proceedings, 2007*: 293-300.
- Listing J, Schlitten R. (1998) Tests if dropouts are missed at random. *Biometrical Journal*, 40: 929-935.
- Listing J, Schlitten R. (2003) A nonparametric test for random dropouts. *Biometrical Journal*, 45: 113-127.
- Littel RC, Miliken GA, Stroup WW, Wolfinger RD. (1996) *SAS System for Mixed Models*, SAS Institute Inc., Cary, NC.
- Little RJ, Rubin DB. (1987) *Statistical Analysis with Missing Data*. John Wiley and Sons, New York.
- Little RJA. (1988) A test of missing completely at random for multivariate data with missing values. *Journal of the American Statistical Association*, 83: 1198-1202.
- Little RJA. (1993) Pattern-mixture models for multivariate incomplete data. *Journal of the American Statistical Association*, 88: 125-134.
- Little RJA. (1994) A class of pattern-mixture models for normal incomplete data. *Biometrika*, 81: 471-483.
- Little RJA. (1995) Modeling the dropout mechanism in repeated-measures studies. *Journal of the American Statistical Association*, 90: 1112-1121.
- Little RJA, Wang Y. (1996) Pattern-mixture models for multivariate incomplete data with covariates. *Biometrics*, 52: 98-111.
- Little R, Yau L. (1996) Intent-to-treat analysis for longitudinal studies with dropouts. *Biometrics*, 52: 1324-1333.
- MacKinnon DP, Warsi G, Dwyer JH. (1995) A simulation study of mediated effect measures. *Multivariate Behavioral Research*, 30: 4162.
- MacKinnon DP, Fairchild AJ, and Fritz MS. (2007) Mediation analysis. *Annual Reviews Psychology*, 58: 593-614.
- MacKinnon DP. (2008) *Introduction to Statistical Mediation Analysis*. Taylor & Francis Group, New York, NY.
- Marcus R, Peritz E, Gabriel KR. (1976) On closed testing procedures with special reference to ordered analysis of variance. *Biometrika*, 63: 655-660.
- Martin BC, Pathak DS, Sharfman MI, Adelman JU, Taylor F, Kwong WJ, Jhingran P. (2000) Validity and reliability of the migraine specific quality of life questionnaire (MSQ Version 2.1). *Headache*, 40: 204-215.

- Matthews JNS, Altman DG, Campbell MJ, Royston P. (1990) Analysis of serial measurements in medical research. *British Medical Journal*, 300: 230-235.
- McDowell I, Newell C. (1996) *Measuring Health: A Guide to Rating Scales and Questionnaires*, 2nd edition. Oxford University Press, New York.
- McNeil BJ, Weichselbaum R, Pauker SG. (1981) Tradeoffs between quality and quantity of life in laryngeal cancer. *New England Journal of Medicine*, 305: 982-987.
- Meinert CL. (1986) *Clinical Trials: Design, Conduct and Analysis*. Oxford University Press, England.
- Michiels B, Molenberghs G, Lipsitz SR. (1999) Selection models and pattern-mixture models for incomplete data with covariates. *Biometrics*, 55: 978-983.
- Miller GA. (1956) The magic number seven plus or minus two: some limits on our capacity for information processing. *Psychological Bulletin*, 63: 81-97.
- Moinpour C, Feigl P, Metch B, et al. (1989) Quality of life endpoints in cancer clinical trials: Review and recommendations. *Journal of the National Cancer Institute*, 81: 485-495.
- Mori M, Woodworth GG, Woolson RF. (1992) Application of empirical Bayes inference to estimation of rate of change in the presence of informative right censoring. *Statistics in Medicine*, 11: 621-631.
- Motzer RJ, Murphy BA, Bacik J, et al. (2000) Phase III trial of Interferon Alfa-2a with or without 13-*cis*-retinoic acid for patients with advanced renal cell carcinoma. *Journal of Clinical Oncology*, 18: 2972-2980.
- Murray GD, Findlay JG. (1988) Correcting for the bias caused by dropouts in hypertension trials. *Statistics in Medicine*, 7: 941-946.
- Naughton MJ, Shumaker SA, Anderson RT, Czajkowski SM. (1996) Psychological aspects of health-related quality of life measurement: Tests and scales, in *Quality of Life and Pharmacoeconomics in Clinical Trials*, Chapter 15. Spilker B, ed. Lippincott-Raven Publishers, Philadelphia.
- Neter J, Wasserman W. (1974) *Applied Linear Statistical Models*, pp 313-317, Richard Irwin, Homewood, Illinois.
- Noll RB, Fairclough DL. (2004) Health-related quality of life: Developmental and psychometric issues (Editorial). *The Journal of Pediatrics*, 145: 8-9.
- Norton NJ, Lipsitz SR. (2001) Multiple imputation in practice: Comparison of software packages for regression models with missing variables. *The American Statistician*, 55: 244-254.

- O'Brien PC. (1984) Procedures for comparing samples with multiple endpoints. *Biometrics*, 40: 1079-1087.
- Omar PZ, Wright EM, Turner RM, Thompon SG. (1999) Analyzing repeated measurements data: A practical comparison of methods. *Statistics in Medicine*, 18: 1587-1608.
- Paik, SSS (1997) The generalized estimating approach when data are not missing completely at random. *Journal of the American Statistical Association*, 92: 1320-1329.
- Pater J, Osoba D, Zee B, Lofters W, Gore M, Dempsey E, Palmer M, Chin C. (1998) Effects of altering the time of administration and the time frame of quality of life assessments in clinical trials: An example using the EORTC QLQ-C30 in a large anti-emetic trial. *Quality of Life Research*, 7: 273-778.
- Patrick DL, Bush JW, Chen MM. (1973) Methods for measuring levels of well-being for a health status index. *Health Services Research*, 8: 228-245.
- Patrick D, Erickson P. (1993) *Health Status and Health Policy: Allocating Resources to Health Care*. Oxford University Press, New York.
- Pauler DK, McCoy S, Moinpour C. (2003) Pattern mixture models for longitudinal quality of life studies in advanced stage disease. *Statistics in Medicine*, 22: 795-809.
- Piantadosi S. (1997) *Clinical Trials: A Methodological Perspective*. John Wiley and Sons, New York.
- Pinheiro JC, Bates DM. (2000) *Mixed-Effects Models in S and S-PLUS*. Springer Verlag, New York, NY.
- Pledger G, Hall D. (1982) Withdrawals from drug trials (letter to editor). *Biometrics*, 38: 276-278.
- Pocock SJ, Geller NL, Tsiatis AA. (1987a) The analysis of multiple endpoints in clinical trials. *Biometrics*, 43: 487-498.
- Pocock SJ, Hughes MD, Lee RJ. (1987b) Statistical problems in the reporting of clinical trials. *New England Journal of Medicine* 317: 426-432.
- Proschan MA, Waclawiw MA. (2000) Practical guidelines for multiplicity adjustment in clinical trials. *Controlled Clinical Trials*, 21: 527-539.
- Raboud JM, Singer J, Thorne A, Schechter MT, Shafran SD. (1998) Estimating the effect of treatment on quality of life in the presence of missing data due to dropout and death. *Quality of Life Research*, 7: 487-494.
- Reitmeir J, Wassmer G. (1999) Resampling-based methods for the analysis of multiple endpoints in clinical trials. *Statistics in Medicine*, 18: 3455-3462.
- Revicki DA, Gold K, Buckman D, Chan K, Kallich JD, Woodley M. (2001)

- Imputing physical function scores missing owing to mortality: Results of a simulation comparing multiple techniques. *Medical Care*, 39: 61-71
- Ribaudo HJ, Thompson SG, Allen-Mersh TG (2000) A joint analysis of quality of life and survival using a random-effect selection model. *Statistics in Medicine*, 19: 3237-3250.
- Ridout MS. (1991) Testing for random dropouts in repeated measurement data. *Biometrics*, 47: 1617-1621.
- Rizopoulos D, Verbeke G, Lesaffre E. (2009) Fully exponential Laplace approximation for the joint modelling of survival and longitudinal data. *Journal of the Royal Statistical Society, Series B*, 71: 637-654.
- Rosenbaum PR, Rubin DB. (1984) Reducing bias in observational studies using subclassification in the propensity score. *Journal of the American Statistical Association*, 79: 516-524.
- Rubin DB, Schenker N. (1986) Multiple imputation for interval estimation from simple random samples with ignorable nonresponse. *Journal of the American Statistical Association*, 81: 366-374.
- Rubin DB. (1987), *Multiple Imputation for Nonresponse in Surveys*, Wiley, New York.
- Rubin DB, Schenker N. (1991) Multiple imputation in health-care data bases: An overview and some applications. *Statistics in Medicine*, 10: 585-598.
- Rubin DB. (1996) Multiple imputation after 18+ years. *Journal of the American Statistical Association*, 91: 473-489.
- Russell IJ, Crofford LJ, Leon T, Cappelleri JC, Bushmakin AG, Whalen E, Barrett JA, Sadosky A. (2009) The effects of pregabalin on sleep disturbance symptoms among individuals with fibromyalgia syndrome. *Sleep Medicine*, 10: 604-610.
- SAS Institute (1996) Chapter 18: The MIXED procedure. *SAS/STAT Software. Changes and Enhancements (through Release 6.11)*, SAS Institute, Cary, NC.
- Schafer J. (1997) *Analysis of Incomplete Multivariate Data*. Chapman and Hall, London.
- Schipper H, Clinich J, McMurray A, Levitt M. (1984) Measuring the quality of life of cancer patients. The Functional Living Index - Cancer: Development and validation. *Journal of Clinical Oncology*, 2: 472-482.
- Schipper H. (1990) Guidelines and caveats for quality of life measurement in clinical practice and research. *Oncology*, 4: 51-57.
- Schluchter MD. (1990) 5V: Unbalanced repeated measures models with struc-

- tured covariance matrices, pp 1207-1244, 1322-1327 in *BMDP Statistical Software Manual*, 2 W.J. Dixon, ed. University of California Press, Berkeley CA.
- Schluchter MD. (1992) Methods for the analysis of informatively censored longitudinal data. *Statistics in Medicine*, 11: 1861-1870.
- Schluchter DM, Green T, Beck GJ. (2001) Analysis of change in the presence of informative censoring: application to a longitudinal clinical trial of progressive renal disease. *Statistics in Medicine*, 20: 989-1007.
- Schluchter MD, Konstan MW, Davis PB. (2002) Jointly modelling the relationship between survival and pulmonary function in cystic fibrosis patients. *Statistics in Medicine*, 21: 1271-1287.
- Schumacher M, Olschewski M, Schulgen G. (1991) Assessment of quality of life in clinical trials. *Statistics in Medicine*, 10: 1915-1930.
- Schuman H, Presser S. (1981) *Questions and Answers in Attitude Surveys: Experiments on Question Form, Wording and Context*. Academic Press, London.
- Schwartz D, Flamant R, Lellouch J. (1980) *Clinical Trials*. Academic Press, London.
- Searle SR. (1971) *Linear Models* pp 46-47, John Wiley and Sons, New York.
- Simes RJ. (1986) An improved Bonferroni procedure for multiple tests of significance. *Biometrika*, 73: 751-754.
- Simon GE, Revicki DA, Grothaus L, Vonkorff M. (1998) SF-summary scores: Are physical and mental health truly distinct? *Medical Care*, 36: 567-572.
- Smith, DM. (1996) Oswald: Object-oriented software for the analysis of longitudinal data in S. <http://www.maths.lancs.ac.uk/Software/Oswald>.
- Smith DM, Schwarz N, Roberts TR, Ubel PA. (2006) Why are you calling me? How study introductions change response patterns. *Quality of Life Research*, 15: 621-630.
- Statistical Solutions (1997) *Solas for Missing Data Analysis 1.0 User Reference*. Statistical Solutions Ltd., Cork, Ireland.
- Spilker B., ed. (1996) *Quality of Life and Pharmacoeconomics in Clinical Trials*. Lippincott-Raven Publishers, Philadelphia.
- Spilker B. (1991) *Guide to Clinical Trials*. Lippincott Williams & Wilkins.
- Staquet M, Aaronson NK, Ahmedzai S, et al. (1992) Editorial: Health-related quality of life research. *Quality of Life Research*, 1: 3.
- Staquet M, Berson R, Osoba D, Machin D. (1996) Guidelines for reporting re-

- sults of quality of life assessments in clinical trials. *Quality of Life Research*, 5: 496-502.
- Staquet MJ, Hayes RD, Fayers PM. (1998) *Quality of life assessment in clinical trials: Methods and practice*. Oxford University Press, New York.
- Streiner DL, Norman GR. (1995) *Health Measurement Scales - A Practical Guide to their Development and Use - Second Edition*. Oxford Medical Publications, Oxford University Press, New York.
- Sugarbaker PH, Barofsky I, Rosenberg SA, Gianola FJ. (1982) Quality of life assessment of patients in extremity sarcoma clinical trials. *Surgery*, 91: 17-23.
- Tandon PK. (1990) Applications of global statistics in analyzing quality of life data. *Statistics in Medicine* 9: 819-827.
- Tang ST, McCorkle R. (2002) Appropriate time frames for data collection in quality of life research among cancer patients at the end of life. *Quality of Life Research*, 11: 145-155.
- Thiebaut R, Jacqmin-Gabba H, Babiker A, Commenges D. (2005) Joint modeling of bivariate longitudinal data with informative dropout and left-censoring, with application to the evolution of CD4+cell count and HIV RNA viral load in response to treatment of HIV infection. *Statistics in Medicine*, 24: 65-82.
- Torrance GW, Thomas WH, Sackett DL. (1971) A utility maximizing model for evaluation of health care programs. *Health Services Research*, 7: 118-133.
- Touloumi G, Pocock SJ, Babiker AG, Daryshire JH. (1999) Estimation and comparison of rates of change in longitudinal studies with informative dropouts. *Statistics in Medicine*, 18: 1215-1233.
- Troxel AB, Lipsitz SR, Harrington DP. (1994) Marginal models for the analysis of longitudinal measurements with nonignorable non-monotone missing data. *Biometrika* 94: 1096-1120.
- Troxel A, Fairclough DL, Curren D, Hahn EA. (1998) Statistical analysis of quality of life data in cancer clinical trials. *Statistics in Medicine* 17: 653-666.
- VanBuuren S, Boshuizen HC, Knook DL. (1999) Multiple imputation of missing blood pressure covariates in survival analysis. *Statistics in Medicine*, 18: 681-694.
- Verbeke G, Molenberghs G. (ed.) (1997) *Linear Mixed Models in Practice: A SAS-Oriented Approach*. Springer, New York.
- Verbeke G, Molenberghs G. (2000) *Linear Mixed Models for Longitudinal Data*. Springer, New York.

- Vonesh EF, Greene T, Schluchter MD. (2006) Shared parameter models for the joint analysis of longitudinal data and event times. *Statistics in Medicine*, 25: 143-163.
- von Hippel PT. (2004) Biases in SPSS 12.0 missing value analysis. *The American Statistician*, 58: 160-164.
- Wang XS, Fairclough DL, Liao Z, Komaki R, Chang JY, Mobley GM, Cleeland CS. (2006) Longitudinal study of the relationship between chemoradiation therapy for non-small-cell lung cancer and patient symptoms. *Journal of Clinical Oncology*, 24: 4485-4491.
- Ware JE, Borrk RH, Davies AR, Lohr KN. (1981) Choosing measures of health status for individuals in general populations. *American Journal of Public Health*, 71: 620-625.
- Ware JE, Snow KK, Kosinski M, Gandek B. (1993) *SF-36 Health Survey: Manual and Interpretation Guide*. The Health Institute, New England Medical Center, Boston, MA.
- Ware J, Kosinski M, Keller SD. (1994) *SF-36 Physical and Mental Component Summary Scales: A User's Manual*. The Health Institute, New England Medical Center, Boston, MA.
- Weeks J. (1992) Quality-of-life assessment: Performance status upstaged? *Journal of Clinical Oncology*, 10: 1827-1829.
- Wei LJ, Johnson WE. (1985) Combining dependent tests with incomplete repeated measurements. *Biometrika* 72: 359-364.
- Westfall PH, Young SS. (1989) p-value adjustment for multiple testing in multivariate binomial models. *Journal of the American Statistical Association*, 84: 780-786.
- Wiklund I, Dimenas E, Wahl M. (1990) Factor of importance when evaluating quality of life in clinical trials. *Controlled Clinical Trials*, 11: 169-179.
- Wilson IB, Cleary PD. (1995) Linking clinical variables with health-related quality of life - a conceptual model of patient outcomes. *Journal of the American Medical Association*, 273: 59-65.
- World Health Organization (1948) Constitution of the World Health Organization. *Basic Documents*, WHO, Geneva.
- World Health Organization (1958) The First Ten Years of the World Health Organization, WHO, Geneva.
- Wu MC, Bailey KR. (1988) Analyzing changes in the presence of informative right censoring caused by death and withdrawal. *Statistics in Medicine*, 7: 337-346.

- Wu MC, Bailey KR. (1989) Estimation and comparison of changes in the presence of informative right censoring: Conditional linear model. *Biometrics*, 45: 939-955.
- Wu MC, Carroll RJ. (1988) Estimation and comparison of changes in the presence of informative right censoring by modeling the censoring process. *Biometrics*, 44: 175-188.
- Wulfsohn M, Tsiatis A. (1997) A joint model for survival and longitudinal data measured with error. *Biometrics*, 53: 330-339.
- Yabroff KR, Linas BP, Schulman K. (1996) Evaluation of quality of life for diverse patient populations. *Breast Cancer Research and Treatment*, 40: 87-104.
- Yao Q, Wei LJ, Hogan JW. (1998) Analysis of incomplete repeated measurements with dependent censoring times. *Biometrika*, 85: 139-149.
- Young T, Maher J. (1999) Collecting quality of life data in EORTC clinical trials - what happens in practice? *Psycho-oncology*, 8: 260-263.
- Yu M, Law NJ, Taylor JMG, Sandler HM. (2004) Joint longitudinal-survival-cure models and their applications to prostate cancer. *Statistica Sinica*, 14: 835-862.
- Zeger SL, Liang K-Y. (1992) An overview of methods for the analysis of longitudinal data. *Statistics in Medicine*, 11: 1825-1839.
- Zhang J., Quan H, Ng J, Stepanavage ME. (1997) Some statistical methods for multiple endpoints in clinical trials. *Controlled Clinical Trials*, 18: 204-221.
- Zwinderman AH. (1990) The measurement of change of quality of life in clinical trials. *Statistics in Medicine*, 9: 931-42.