

Bibliografie

- Abram, K. M. and Teplin, L. A. (1991), 'Co-occurring disorders among mentally ill jail detainees. Implications for public policy', *The American Psychologist* 46(10), 1036-45.
- Adamson, S. J. and Sellman, J. D. (2003), 'A prototype screening instrument for cannabis use disorder: The Cannabis Use Disorders Identification Test (CUDIT) in an alcohol-dependent clinical sample', *Drug and Alcohol Review* 22(3), 309-15.
- Aharonovich, E., Garawi, F., Bisaga, A., Brooks, D., Raby, W. N., Rubin, E., Nunes, E. V. and Levin, F. R. (2006), 'Concurrent cannabis use during treatment for comorbid ADHD and cocaine dependence: effects on outcome', *The American Journal of Drug and Alcohol Abuse* 32(4), 629-35.
- Aichhorn, W., Santeler, S., Stelzig-Scholer, R., Kemmler, G., Steinmayr-Gensluckner, M. and Hinterhuber, H. (2008), '[Prevalence of psychiatric disorders among homeless adolescents]', *Neuropsychiatrie: Klinik, Diagnostik, Therapie Und Rehabilitation* 22(3), 180-88.
- Alexander, M. J., Haugland, G., Lin, S. P., Bertollo, D. N. and McCorry, F. A. (2008), 'Mental health screening in addiction, corrections and social service settings: Validating the MMS', *International Journal of Mental Health and Addiction* 6(1), 105-19.
- APA (1980), *Diagnostic and statistical manual of mental disorders, third edition*, American Psychiatric Association, Washington, D.C.
- APA (1987), *Diagnostic and statistical manual of mental disorders, revised third edition*, American Psychiatric Association, Washington, D.C.
- APA (1994), *Diagnostic and statistical manual of mental disorders, fourth edition*, American Psychiatric Association, Washington, D.C.
- APA (2000), *Diagnostic and statistical manual of mental disorders, fourth edition, text revision*, American Psychiatric Association, Washington, D.C.
- APA (2013), *Diagnostic and statistical manual of mental disorders, fifth edition*, American Psychiatric Association, Washington, D.C.
- Araos, P., Vergara-Moragues, E., Pedraz, M., Pavón, F. J., Campos Cloute, R., et al. (2014), '[Psychopathological comorbidity in cocaine users in outpatient treatment]', *Adicciones* 26(1), 15-26.
- Arendt, M., Sher, L., Fjordback, L., Brandholdt, J. and Munk-Jorgensen, P. (2007), 'Parental alcoholism predicts suicidal behavior in adolescents and young adults with cannabis dependence', *International Journal of Adolescent Medicine and Health* 19(1), 67-77.
- Arendt, M., Munk-Jorgensen, P., Sher, L. and Jensen, S. O. W. (2011), 'Mortality among individuals with cannabis, cocaine, amphetamine, MDMA, and opioid use disorders: A nationwide follow-up study of Danish substance users in treatment', *Drug and Alcohol Dependence* 114(2-3), 134-39.

- Astals, M., Domingo-Salvany, A., Buenaventura, C. C., Tato, J., Vazquez, J. M., Martín-Santos, R. and Torrens, M. (2008), 'Impact of substance dependence and dual diagnosis on the quality of life of heroin users seeking treatment', *Substance Use & Misuse* 43(5), 612-32.
- Astals, M., Díaz, L., Domingo-Salvany, A., Martín-Santos, R., Bulbena, A. and Torrens, M. (2009), 'Impact of co-occurring psychiatric disorders on retention in a methadone maintenance program: an 18-month follow-up study', *International Journal of Environmental Research and Public Health* 6(11), 2822-32.
- Australian Institute of Health and Welfare (2005), *National comorbidity initiative: A review of data collections relating to people with coexisting substance use and mental health disorders*, AIHW Cat. No. PHE 60. (Drug Statistics Series No. 14). Canberra: AIHW.
- Baldacchino, A. (2007), 'Co-morbid substance misuse and mental health problems: policy and practice in Scotland', *The American Journal on Addictions* 16(3), 147-59.
- Baldacchino, A. and Corkery, J. (editors) (2006), *Comorbidity: perspectives across Europe*, European Collaborating Centres in Addiction Studies, London.
- Baldacchino, A., Blair, H., Scherbaum, N., Grosse-Vehne, E., Riglietta, M., et al. (2009), 'Drugs and psychosis project: a multi-centre European study on comorbidity', *Drug and Alcohol Review* 28(4), 379-89.
- Ball, S. A., Rounsaville, B. J., Tennen, H. and Kranzler, H. R. (2001), 'Reliability of personality disorder symptoms and personality traits in substance-dependent inpatients', *Journal of Abnormal Psychology* 110(2), 341-52.
- Barkley, R. A., Fischer, M., Smallish, L. and Fletcher, K. (2003), 'Does the treatment of attention-deficit/hyperactivity disorder with stimulants contribute to drug use/abuse? A 13-year prospective study', *Pediatrics* 111(1), 97-109.
- Barkus, E. and Murray, R. M. (2010), 'Substance use in adolescence and psychosis: clarifying the relationship', *Annual Review of Clinical Psychology* 6, 365-89.
- Barnett, J. H., Werners, U., Secher, S. M., Hill, K. E., Brazil, R., et al. (2007), 'Substance use in a population-based clinic sample of people with first-episode psychosis', *British Journal of Psychiatry* 190, 515-20.
- Batalla, A., Garcia-Rizo, C., Castellví, P., Fernandez-Egea, E., Yücel, M., et al. (2013), 'Screening for substance use disorders in first-episode psychosis: Implications for readmission', *Schizophrenia Research* 146(1-3), 125-31.
- Beijer, U., Andréasson, A., Agren, G. and Fugelstad, A. (2007), 'Mortality, mental disorders and addiction: a 5-year follow-up of 82 homeless men in Stockholm', *Nordic Journal of Psychiatry* 61(5), 363-68.
- Benjamin, A. B., Mossman, D., Graves, N. S. and Sanders, R. D. (2006), 'Tests of a symptom checklist to screen for comorbid psychiatric disorders in alcoholism', *Comprehensive Psychiatry* 47(3), 227-33.
- Berkson, J. (1946), 'Limitations of the application of fourfold table analysis to hospital data', *Biometrics* 2(3), 47-53.

- Berman, A. H., Bergman, H., Palmstierna, T. and Schlyter, F. (2005), 'Evaluation of the Drug Use Disorders Identification Test (DUDIT) in criminal justice and detoxification settings and in a Swedish population sample', *European Addiction Research* 11(1), 22-31.
- Bernacer, J., Corlett, P. R., Ramachandra, P., McFarlane, B., Turner, D. C., et al. (2013), 'Methamphetamine-induced disruption of frontostriatal reward learning signals: relation to psychotic symptoms', *American Journal of Psychiatry* 170(11), 1326-34.
- Bizzarri, J. V., Rucci, P., Sbrana, A., Miniati, M., Raimondi, F., et al. (2009), 'Substance use in severe mental illness: self-medication and vulnerability factors', *Psychiatry Research* 165(1-2), 88-95.
- Błachut, M., Badura-Brzoza, K., Jarzab, M., Gorczyca, P. and Hese, R. T. (2013), '[Dual diagnosis in psychoactive substance abusing or dependent persons]', *Psychiatria Polska* 47(2), 335-52.
- Blanco, C., Alegría, A. A., Liu, S.-M., Secades-Villa, R., Sugaya, L., Davies, C. and Nunes, E. V. (2012), 'Differences among major depressive disorder with and without co-occurring substance use disorders and substance-induced depressive disorder: results from the National Epidemiologic Survey on Alcohol and Related Conditions', *The Journal of Clinical Psychiatry* 73(6), 865-73.
- Blinder, B. J., Cumella, E. J. and Sanathara, V. A. (2006), 'Psychiatric comorbidities of female inpatients with eating disorders', *Psychosomatic Medicine* 68(3), 454-62.
- Boden, M. T. and Moos, R. (2009), 'Dually diagnosed patients' responses to substance use disorder treatment', *Journal of Substance Abuse Treatment* 37(4), 335-45.
- Booth, B. M., Walton, M. A., Barry, K. L., Cunningham, R. M., Chermack, S. T. and Blow, F. C. (2011), 'Substance use depression and mental health functioning in patients seeking acute medical care in an inner-city ED', *Journal of Behavioral Health Services and Research* 38(3), 358-72.
- Brady, K. T. and Sinha, R. (2005), 'Co-occurring mental and substance use disorders: the neurobiological effects of chronic stress', *American Journal of Psychiatry* 162(8), 1483-93.
- Brady, K. T. and Verduin, M. L. (2005), 'Pharmacotherapy of comorbid mood, anxiety, and substance use disorders', *Substance Use & Misuse* 40(13-14), 2021-41, 2043-48.
- Brand, E. F. J. M., Lucker, T. P. C. and van den Hurk, A. A. (2009), '[Addiction and recidivism in forensic psychiatry]', *Tijdschrift Voor Psychiatrie* 51(11), 813-20.
- Broadhead, W. E., Leon, A. C., Weissman, M. M., Barrett, J. E., Blacklow, R. S., et al. (1995), 'Development and validation of the SDDS-PC screen for multiple mental disorders in primary care', *Archives of Family Medicine* 4(3), 211-19.
- Burnam, M. A. and Watkins, K. E. (2006), 'Substance abuse with mental disorders: specialized public systems and integrated care', *Health Affairs* 25(3), 648-58.
- Carey, M. P., Carey, K. B., Maisto, S. A., Gordon, C. M. and Venable, P. A. (2001), 'Prevalence and correlates of sexual activity and HIV-related risk behavior among psychiatric outpatients', *Journal of Consulting and Clinical Psychology* 69(5), 846-50.

- Carra, G. and Johnson, S. (2009), 'Variations in rates of comorbid substance use in psychosis between mental health settings and geographical areas in the UK. A systematic review', *Social Psychiatry and Psychiatric Epidemiology* 44(6), 429-47.
- Carra, G., Johnson, S., Bebbington, P., Angermeyer, M. C., Heider, D., Brugha, T., Azorin, J. M. and Toumi, M. (2012), 'The lifetime and past-year prevalence of dual diagnosis in people with schizophrenia across Europe: findings from the European Schizophrenia Cohort (EuroSC)', *European Archives of Psychiatry and Clinical Neuroscience* 262(7), 607-16.
- Carroll, J. F. X. and McGinley, J. J. (2001), 'A screening form for identifying mental health problems in alcohol/other drug dependent persons', *Alcoholism Treatment Quarterly* 19(4), 33-47.
- Casadio, P., Olivoni, D., Ferrari, B., Pintori, C., Speranza, E., et al. (2014), 'Personality disorders in addiction outpatients: prevalence and effects on psychosocial functioning', *Substance Abuse: Research and Treatment* 8, 17-24.
- Casares-López, M. J., González-Menéndez, A., Bobes-Bascarán, M. T., Secades, R., Martínez-Cordero, A. and Bobes, J. (2011), '[Need for the assessment of dual diagnosis in prisons]', *Adicciones* 23(1), 37-44.
- Caton, C. L. M., Shrout, P. E., Eagle, P. F., Opler, L. A., Felix, A. and Dominguez, B. (1994), 'Risk factors for homelessness among schizophrenic men: a case-control study', *American Journal of Public Health* 84(2), 265-70.
- Clark, C. and Young, M. S. (2009), 'Outcomes of mandated treatment for women with histories of abuse and co-occurring disorders', *Journal of Substance Abuse Treatment* 37(4), 346-52.
- Colins, O., Vermeiren, R., Vahl, P., Markus, M., Broekaert, E. and Doreleijers, T. (2011), 'Psychiatric disorder in detained male adolescents as risk factor for serious recidivism', *Canadian Journal of Psychiatry* 56(1), 44-50.
- Combaluzier, S., Gouvernet, B. and Bernoussi, A. (2009), '[Impact of personality disorders in a sample of 212 homeless drug users]', *L'Encéphale* 35(5), 448-53.
- Conner, K. R. (2011), 'Clarifying the relationship between alcohol and depression', *Addiction* 106(5), 915-16.
- Corte, C. and Stein, K. F. (2000), 'Eating disorders and substance use. An examination of behavioral associations', *Eating Behaviors* 1(2), 173-89.
- Coscas, S., Benyamina, A., Reynaud, M. and Karila, L. (2013), '[Psychiatric complications of cannabis use]', *La Revue Du Praticien* 63(10), 1426-28.
- Craig, T. K. J., Johnson, S., McCrone, P., Afuwape, S., Hughes, E., et al. (2008), 'Integrated care for co-occurring disorders: psychiatric symptoms, social functioning, and service costs at 18 months', *Psychiatric Services* 59(3), 276-82.
- Cuenca-Royo, A. M., Sánchez-Niubó, A., Forero, C. G., Torrens, M., Suelves, J. M. and Domingo-Salvany, A. (2012), 'Psychometric properties of the CAST and SDS scales in young adult cannabis users', *Addictive Behaviors* 37(6), 709-15.
- Cuenca-Royo, A. M., Torrens, M., Sánchez-Niubó, A., Suelves, J. M. and Domingo-Salvany, A. (2013), 'Psychiatric morbidity among young-adults cannabis users', *Adicciones* 25(1), 45-53.

- Cuffel, B. J., Shumway, M., Chouljian, T. L. and MacDonald, T. (1994), 'A longitudinal study of substance use and community violence in schizophrenia', *The Journal of Nervous and Mental Disease* 182(12), 704-8.
- Curran, G. M., Sullivan, G., Williams, K., Han, X., Allee, E. and Kotrla, K. J. (2008), 'The association of psychiatric comorbidity and use of the emergency department among persons with substance use disorders: an observational cohort study', *BMC Emergency Medicine* 8, 17.
- Curran, H. V. and Robjant, K. (2006), 'Eating attitudes, weight concerns and beliefs about drug effects in women who use ecstasy', *Journal of Psychopharmacology* 20(3), 425-31.
- De Beurs, E., Beekman, A., Geerlings, S., Deeg, D., Van Dyck, R. and Van Tilburg, W. (2001), 'On becoming depressed or anxious in late life: similar vulnerability factors but different effects of stressful life events', *British Journal of Psychiatry* 179, 426-31.
- De Wilde, J., Broekaert, E., Rosseel, Y., Delespaul, P. and Soyez, V. (2007), 'The role of gender differences and other client characteristics in the prevalence of DSM-IV affective disorders among a European therapeutic community population', *The Psychiatric Quarterly* 78(1), 39-51.
- Delgadoillo, J., Payne, S., Gilbody, S., Godfrey, C., Gore, S., Jessop, D. and Dale, V. (2011), 'How reliable is depression screening in alcohol and drug users? A validation of brief and ultra-brief questionnaires', *Journal of Affective Disorders* 134(1-3), 266-71.
- Dell'osso, L. and Pini, S. (2012), 'What did we learn from research on comorbidity in psychiatry? Advantages and limitations in the forthcoming DSM-V era', *Clinical Practice and Epidemiology in Mental Health* 8, 180-84.
- DeLorenze, G. N., Tsai, A.-L., Horberg, M. A. and Quesenberry, C. P. (2014), 'Cost of care for HIV-infected patients with co-occurring substance use disorder or psychiatric disease: report from a large, integrated health plan', *AIDS Research and Treatment* doi:10.1155/2014/570546.
- Demyttenaere, K., Bruffaerts, R., Posada-Villa, J., Gasquet, I., Kovess, V., et al. (2004), 'Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys', *JAMA* 291(21), 2581-90.
- Department of Health (2002), *Mental health policy implementation guide: Dual diagnosis goodpractice guide*, Department of Health, London.
- Department of Health and the devolved administrations (2007), *Drug misuse and dependence: UK guidelines on clinical management*, Department of Health and the devolved administrations, London (k dispozici na http://www.nta.nhs.uk/uploads/clinical_guidelines_2007.pdf).
- Derogatis, L. R. and Melisaratos, N. (1983), 'The Brief Symptom Inventory: an introductory report', *Psychological Medicine* 13(3), 595-605.
- Derogatis, L. R., Lipman, R. S. and Covi, L. (1973), 'SCL-90: an outpatient psychiatric rating scale. Preliminary report', *Psychopharmacology Bulletin* 9(1), 13-28.

- Di Furia, L., Pizza, M., Rampazzo, L. and Corti, A. (2006), 'The Italian experience: (B) Comorbidity in Padua', in *Co-morbidity: Perspectives across Europe* edited by J. Corkery and A. Baldacchino, European Collaborating Centres on Addiction Studies (ECCAS), London, 186-97.
- Drake, R. E., Mercer-McFadden, C., Mueser, K. T., McHugo, G. J., Bond, G. R. (1998), 'Review of integrated mental health and substance abuse treatment for patients with dual disorders', *Schizophrenia Bulletin* 24(4), 589-608.
- Drake, R. E., Goldman, H. H., Leff, H. S., Lehman, A. F., Dixon, L., Mueser, K. T. and Torrey, W. C. (2001), 'Implementing evidence-based practices in routine mental health service settings', *Psychiatric Services* 52(2), 179-82.
- Drake, R. E., Wallach, M. A. and McGovern, M. P. (2005), 'Future directions in preventing relapse to substance abuse among clients with severe mental illnesses', *Psychiatric Services* 56(10), 1297-1302.
- Durvasula, R. and Miller, T. R. (2014), 'Substance abuse treatment in persons with HIV/AIDS: challenges in managing triple diagnosis', *Behavioral Medicine* 40(2), 43-52.
- Einarsson, E., Sigurdsson, J. F., Gudjonsson, G. H., Newton, A. K. and Bragason, O. O. (2009), 'Screening for attention-deficit hyperactivity disorder and co-morbid mental disorders among prison inmates', *Nordic Journal of Psychiatry* 63(5), 361-67.
- Elonheimo, H., Niemelä, S., Parkkola, K., Multimäki, P., Helenius, H., Nuutila, A. M. and Sourander, A. (2007), 'Police-registered offenses and psychiatric disorders among young males: the Finnish "from a boy to a man" birth cohort study', *Social Psychiatry and Psychiatric Epidemiology* 42(6), 477-84.
- EMCDDA and Pempidou Group (2000), *Treatment demand indicator, standard protocol 2.0*, European Monitoring Centre for Drugs and Drug Addiction, Lisbon (k dispozici na <http://www.emcdda.europa.eu/html.cfm/index65315EN.html>).
- EMCDDA (2004), 'Co-morbidity', in *Annual Report 2004: The state of the drugs problem in the European Union and Norway*, Office for Official Publications of the European Communities, Luxembourg, 94-102 (k dispozici na <http://www.emcdda.europa.eu/html.cfm/index34901EN.html>).
- EMCDDA (2012), *Treatment demand indicator (TDI) standard protocol 3.0: Guidelines for reporting data on people entering drug treatment in European countries*, Publications Office of the European Union, Luxembourg (k dispozici na <http://www.emcdda.europa.eu/publications/manuals/tdi-protocol-3.0>).
- EMCDDA (2013a), *Co-morbid substance use and mental disorders in Europe: a review of the data*, EMCDDA Papers, Publications Office of the European Union, Luxembourg (k dispozici na <http://www.emcdda.europa.eu/publications/emcdda-papers/co-morbidity>).
- EMCDDA (2013b), *Models of addiction*, Publications Office of the European Union, Luxembourg (k dispozici na <http://www.emcdda.europa.eu/publications/insights/models-addiction>).
- EMCDDA (2015), *The misuse of benzodiazepines among high-risk opioid users in Europe* (<http://www.emcdda.europa.eu/topics/pods/benzodiazepines>), zobrazeno 7. srpna 2015.

- Enatescu, V. R. and Dehelean, L. (2006), 'Psychiatric comorbidity in patients with substance misuse in Romania', in *Co-morbidity: Perspectives across Europe* edited by J. Corkery and A. Baldacchino, European Collaborating Centres on Addiction Studies (ECCAS), London, 221-34.
- Endicott, J. and Spitzer, R. L. (1978), 'A diagnostic interview: the schedule for affective disorders and schizophrenia', *Archives of General Psychiatry* 35(7), 837-44.
- Evans, E. A. and Sullivan, M. A. (2014), 'Abuse and misuse of antidepressants', *Substance Abuse and Rehabilitation* 5, 107-20.
- Farrell, M. (2001), 'Nicotine, alcohol and drug dependence and psychiatric comorbidity. Results of a national household survey', *The British Journal of Psychiatry* 179(5), 432-37.
- Fatséas, M., Denis, C., Lavie, E. and Auriacombe, M. (2010), 'Relationship between anxiety disorders and opiate dependence. A systematic review of the literature. Implications for diagnosis and treatment', *Journal of Substance Abuse Treatment* 38(3), 220-30.
- Feighner, J. P., Robins, E., Guze, S. B., Woodruff, R. A., Winokur, G. and Munoz, R. (1972), 'Diagnostic criteria for use in psychiatric research', *Archives of General Psychiatry* 26(1), 57-63.
- Feinstein, A. R. (1970), 'The pre-therapeutic classification of co-morbidity in chronic disease', *Journal of Chronic Diseases* 23(7), 455-68.
- Fiorentini, A., Volonteri, L. S., Dragogna, F., Rovera, C., Maffini, M., Mauri, M. C. and Altamura, C. A. (2011), 'Substance-induced psychoses: a critical review of the literature', *Current Drug Abuse Reviews* 4(4), 228-40.
- First, M. B. (1997), *User's guide for the structured clinical interview for DSM-IV axis II personality disorders: SCID-II*, American Psychiatric Press, Washington, D.C.
- Fischer, S. and Le Grange, D. (2007), 'Comorbidity and high-risk behaviors in treatment-seeking adolescents with bulimia nervosa', *International Journal of Eating Disorders* 40(8), 751-53.
- Flynn, P. M. and Brown, B. S. (2008), 'Co-occurring disorders in substance abuse treatment: issues and prospects', *Journal of Substance Abuse Treatment* 34(1), 36-47.
- Franco, D. L., Dorer, D. J., Keel, P. K., Jackson, S., Manzo, M. P. and Herzog, D. B. (2005), 'How do eating disorders and alcohol use disorder influence each other?', *The International Journal of Eating Disorders* 38(3), 200-207.
- Funk, S. (2013), '[Pharmacological treatment in alcohol-, drug- and benzodiazepine-dependent patients: the significance of trazodone]', *Neuropsychopharmacologia Hungarica* 15(2), 85-93.
- Goldberg, D. (1978), *Manual of the general health questionnaire*, National Foundation for Educational Research, Windsor.
- Goldberg, D. (1986), 'Use of the general health questionnaire in clinical work', *British Medical Journal (Clinical Research Ed)* 293(6556), 1188-89.
- Grant, B. F., Harford, T. C., Dawson, D. A., Chou, P. S. and Pickering, R. P. (1995), 'The Alcohol Use Disorder and Associated Disabilities Interview Schedule (AUDADIS): reliability of alcohol and drug modules in a general population sample', *Drug and Alcohol Dependence* 39(1), 37-44.

- Grant, B. F., Dawson, D. A. and Hasin, D. S. (2001), *The Alcohol Use Disorder and Associated Disabilities Interview Schedule DSM-IV version*, National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD
- Grant, B. F., Dawson, D. A., Stinson, F. S., Chou, P. S., Kay, W. and Pickering, R. (2003), 'The Alcohol Use Disorder and Associated Disabilities Interview Schedule-IV (AUDADIS-IV): reliability of alcohol consumption, tobacco use, family history of depression and psychiatric diagnostic modules in a general population sample', *Drug and Alcohol Dependence* 71(1), 7-16.
- Green, A. I. (2005), 'Schizophrenia and comorbid substance use disorder: effects of antipsychotics', *The Journal of Clinical Psychiatry* 66 Suppl 6, 21-26.
- Green, A. I. (2006), 'Treatment of schizophrenia and comorbid substance abuse: pharmacologic approaches', *The Journal of Clinical Psychiatry* 67 Suppl 7, 31-35; quiz 36-37.
- Green, A. I. and Brown, E. S. (2006), 'Comorbid schizophrenia and substance abuse', *The Journal of Clinical Psychiatry* 67(9), e08.
- Green, A. I., Noordsy, D. L., Brunette, M. F. and O'Keefe, C. (2008), 'Substance abuse and schizophrenia: pharmacotherapeutic intervention', *Journal of Substance Abuse Treatment* 34(1), 61-71.
- Greenberg, G. A. and Rosenheck, R. A. (2014), 'Psychiatric correlates of past incarceration in the national co-morbidity study replication', *Criminal Behaviour and Mental Health* 24(1), 18-35.
- Gregg, L., Barrowclough, C. and Haddock, G. (2007), 'Reasons for increased substance use in psychosis', *Clinical Psychology Review* 27(4), 494-510.
- Gregorowski, C., Seedat, S. and Jordaan, G. P. (2013), 'A clinical approach to the assessment and management of co-morbid eating disorders and substance use disorders', *BMC Psychiatry* 13, 289.
- Gual, A. (2007), 'Dual diagnosis in Spain', *Drug and Alcohol Review* 26(1), 65-71.
- Halmoy, A., Fasmer, O. B., Gillberg, C. and Haavik, J. (2009), 'Occupational outcome in adult ADHD: impact of symptom profile, comorbid psychiatric problems, and treatment: a cross-sectional study of 414 clinically diagnosed adult ADHD patients', *Journal of Attention Disorders* 13(2), 175-87.
- Harrison, I., Joyce, E. M., Mutsatsa, S. H., Hutton, S. B., Huddy, V., Kapasi, M. and Barnes, T. R. E. (2008), 'Naturalistic follow-up of co-morbid substance use in schizophrenia: the West London first-episode study', *Psychological Medicine* 38(1), 79-88.
- Harsch, S., Bergk, J. E., Steinert, T., Keller, F. and Jockusch, U. (2006), 'Prevalence of mental disorders among sexual offenders in forensic psychiatry and prison', *International Journal of Law and Psychiatry* 29(5), 443-49.
- Hasin, D. S., Trautman, K. D., Miele, G. M., Samet, S., Smith, M. and Endicott, J. (1996), 'Psychiatric Research Interview for Substance and Mental Disorders (PRISM): reliability for substance abusers', *The American Journal of Psychiatry* 153(9), 1195-1201.

- Hasin, D., Carpenter, K. M., McCloud, S., Smith, M. and Grant, B. F. (1997), 'The alcohol use disorder and associated disabilities interview schedule (AUDADIS): reliability of alcohol and drug modules in a clinical sample', *Drug and Alcohol Dependence* 44(2-3), 133-41.
- Hasin, D. S., Samet, S., Meydan, J., Miele, G. and Endicott, J. (2001), *Psychiatric Research Interview for Substance and Mental Disorders (PRISM-IV)*, version 6.0.
- Hasin, D. S. and Grant, B. F. (2004), 'The co-occurrence of DSM-IV alcohol abuse in DSM-IV alcohol dependence: results of the National Epidemiologic Survey on Alcohol and Related Conditions on heterogeneity that differ by population subgroup', *Archives of General Psychiatry* 61(9), 891-96.
- Hasin, D., Samet, S., Nunes, E., Meydan, J., Matseoane, K. and Waxman, R. (2006), 'Diagnosis of comorbid psychiatric disorders in substance users assessed with the Psychiatric Research Interview for Substance and Mental Disorders for DSM-IV', *The American Journal of Psychiatry* 163(4), 689-96.
- Haug, N. A., Heinberg, L. J. and Guarda, A. S. (2001), 'Cigarette smoking and its relationship to other substance use among eating disordered inpatients', *Eating and Weight Disorders : EWD* 6(3), 130-39.
- Havens, J. R., Ompad, D. C., Latkin, C. A., Fuller, C. M., Arria, A. M., Vlahov, D. and Strathdee, S. A. (2005), 'Suicidal ideation among African-American non-injection drug users', *Ethnicity & Disease* 15(1), 110-15.
- Haver, B. (1997), 'Screening for psychiatric comorbidity among female alcoholics: the use of a questionnaire (SCL-90) among women early in their treatment programme', *Alcohol and Alcoholism* 32(6), 725-30.
- Haw, C. M. and Hawton, K. (2011), 'Problem drug use, drug misuse and deliberate self-harm: trends and patient characteristics, with a focus on young people, Oxford, 1993-2006', *Social Psychiatry and Psychiatric Epidemiology* 46(2), 85-93.
- Helzer, J. E. (1981), 'The use of a structured diagnostic interview for routine psychiatric evaluations', *Journal Nervous & Mental Disease* 169(1), 45-9.
- Helzer, J. E. and Canino, G. J. (editors) (1992), *Alcoholism in North America, Europe, and Asia*, Oxford University Press, Oxford.
- Hermle, L., Szlak-Rubin, R., Täschner, K. L., Peukert, P. and Batra, A. (2013), '[Substance use associated disorders: frequency in patients with schizophrenic and affective psychoses]', *Der Nervenarzt* 84(3), 315-25.
- Herrero, M. J., Domingo-Salvany, A., Torrens, M. and Brugal, M. T. (2008), 'Psychiatric comorbidity in young cocaine users: induced versus independent disorders', *Addiction* 103(2), 284-93.
- Herzog, D. B., Franko, D. L., Dorer, D. J., Keel, P. K., Jackson, S. and Manzo, M. P. (2006), 'Drug abuse in women with eating disorders', *The International Journal of Eating Disorders* 39(5), 364-68.
- Hesse, M. (2009), 'Integrated psychological treatment for substance use and co-morbid anxiety or depression vs. treatment for substance use alone. A systematic review of the published literature', *BMC Psychiatry* 9, 6.
- Huntley, Z., Maltezos, S., Williams, C., Morinan, A., Hammon, A., et al. (2012), 'Rates of undiagnosed attention deficit hyperactivity disorder in London drug and alcohol detoxification units', *BMC Psychiatry* 12, 223.

- Charzynska, K., Hyldager, E., Baldacchino, A., Greacen, T., Henderson, Z., et al. (2011), 'Comorbidity patterns in dual diagnosis across seven European sites', *The European Journal of Psychiatry* 25(4), 179-91.
- Illomäki, R., Riala, K., Hakko, H., Lappalainen, J., Ollinen, T., Räsänen, P. and Timonen, M. (2008), 'Temporal association of onset of daily smoking with adolescent substance use and psychiatric morbidity', *European Psychiatry* 23(2), 85-91.
- Janca, A., Ustün, T. B. and Sartorius, N. (1994), 'New versions of World Health Organization instruments for the assessment of mental disorders', *Acta Psychiatrica Scandinavica* 90(2), 73-83.
- Jaworski, F., Dubertret, C., Ades, J. and Gorwood, P. (2011), 'Presence of co-morbid substance use disorder in bipolar patients worsens their social functioning to the level observed in patients with schizophrenia', *Psychiatry Research* 185(1-2), 129-34.
- Katzman, M. A., Bleau, P., Blier, P., Chokka, P., Kjernisted, K., et al. (2014), 'Canadian clinical practice guidelines for the management of anxiety, posttraumatic stress and obsessive-compulsive disorders', *BMC Psychiatry* 14 Suppl 1, S1.
- Kelly, T. M. and Daley, D. C. (2013), 'Integrated treatment of substance use and psychiatric disorders', *Social Work in Public Health* 28(3-4), 388-406.
- Kelly, T. M., Daley, D. C. and Douaihy, A. B. (2012), 'Treatment of substance abusing patients with comorbid psychiatric disorders', *Addictive Behaviors* 37(1), 11-24.
- Khalsa, J. H., Treisman, G., McCance-Katz, E. and Tedaldi, E. (2008), 'Medical consequences of drug abuse and co-occurring infections: research at the National Institute on Drug Abuse', *Substance Abuse* 29(3), 5-16.
- Khantzian, E. J. (1985), 'The self-medication hypothesis of addictive disorders: focus on heroin and cocaine dependence', *The American Journal of Psychiatry* 142(11), 1259-64.
- King, V. L., Kidorf, M. S., Stoller, K. B. and Brooner, R. K. (2000), 'Influence of psychiatric comorbidity on HIV risk behaviors: changes during drug abuse treatment', *Journal of Addictive Diseases* 19(4), 65-83.
- Knight, J. R., Sherritt, L., Shrier, L. A., Harris, S. K. and Chang, G. (2002), 'Validity of the CRAFFT substance abuse screening test among adolescent clinic patients', *Archives of Pediatrics & Adolescent Medicine* 156(6), 607-14.
- Kollins, S. H. (2008), 'A qualitative review of issues arising in the use of psycho-stimulant medications in patients with ADHD and co-morbid substance use disorders', *Current Medical Research and Opinion* 24(5), 1345-57.
- Krausz, R. M., Clarkson, A. F., Strehlau, V., Torchalla, I., Li, K. and Schuetz, C. G. (2013), 'Mental disorder, service use, and barriers to care among 500 homeless people in 3 different urban settings', *Social Psychiatry and Psychiatric Epidemiology* 48(8), 1235-43.
- Krug, I., Treasure, J., Anderluh, M., Bellodi, L., Cellini, E., et al. (2008), 'Present and lifetime comorbidity of tobacco, alcohol and drug use in eating disorders: a European multicenter study', *Drug and Alcohol Dependence* 97(1-2), 169-79.
- Lacey, J. H. and Evans, C. D. (1986), 'The impulsivist: a multi-impulsive personality disorder', *British Journal of Addiction* 81(5), 641-49.

- Lambert, M. T., LePage, J. P. and Schmitt, A. L. (2003), 'Five-year outcomes following psychiatric consultation to a tertiary care emergency room', *The American Journal of Psychiatry* 160(7), 1350-53.
- Langas, A.-M., Malt, U. F. and Opjordsmoen, S. (2011), 'Comorbid mental disorders in substance users from a single catchment area – a clinical study', *BMC Psychiatry* 11, 25.
- Langas, A.-M., Malt, U. F. and Opjordsmoen, S. (2012a), 'In-depth study of personality disorders in first-admission patients with substance use disorders', *BMC Psychiatry* 12, 180.
- Langas, A.-M., Malt, U. F. and Opjordsmoen, S. (2012b), 'Substance use disorders and comorbid mental disorders in first-time admitted patients from a catchment area', *European Addiction Research* 18(1), 16-25.
- Lecrubier, Y., Sheehan, D., Weiller, E., Amorim, P., Bonora, I., Harnett Sheehan, K., Janavs, J. and Dunbar, G. (1997), 'The Mini International Neuropsychiatric Interview (MINI). A short diagnostic structured interview: reliability and validity according to the CIDI', *European Psychiatry* 12(5), 224-31.
- Leeies, M., Pagura, J., Sareen, J. and Bolton, J. M. (2010), 'The use of alcohol and drugs to self-medicate symptoms of posttraumatic stress disorder', *Depression and Anxiety* 27(8), 731-36.
- Legleye, S., Karila, L., Beck, F. and Reynaud, M. (2007), 'Validation of the CAST, a general population Cannabis Abuse Screening Test', *Journal of Substance Use* 12(4), 233-42.
- Leray, E., Camara, A., Drapier, D., Riou, F., Bougeant, N., et al. (2011), 'Prevalence, characteristics and comorbidities of anxiety disorders in France: results from the "Mental Health in General Population" survey (MHGP)', *European Psychiatry* 26(6), 339-45.
- Levin, F. R., Bisaga, A., Raby, W., Aharonovich, E., Rubin, E., et al. (2008), 'Effects of major depressive disorder and attention-deficit/hyperactivity disorder on the outcome of treatment for cocaine dependence', *Journal of Substance Abuse Treatment* 34(1), 80-89.
- Levin, F. R., Mariani, J., Brooks, D. J., Pavlicova, M., Nunes, E. V., et al. (2013), 'A randomized double-blind, placebo-controlled trial of venlafaxine-extended release for co-occurring cannabis dependence and depressive disorders', *Addiction* 108(6), 1084-94.
- Lingford-Hughes, A. R., Welch, S., Peters, L. and Nutt, D. J. (2012), 'BAP updated guidelines: evidence-based guidelines for the pharmacological management of substance abuse, harmful use, addiction and comorbidity: recommendations from BAP', *Journal of Psychopharmacology* 26(7), 899-952.
- Loftis, J. M., Matthews, A. M. and Hauser, P. (2006), 'Psychiatric and substance use disorders in individuals with hepatitis C: epidemiology and management', *Drugs* 66(2), 155-74.
- Lukasiewicz, M., Blecha, L., Falissard, B., Neveu, X., Benyamina, A., Reynaud, M. and Gasquet, I. (2009), 'Dual diagnosis: prevalence, risk factors, and relationship with suicide risk in a nationwide sample of French prisoners', *Alcoholism, Clinical and Experimental Research* 33(1), 160-68.

- Magidson, J. F., Wang, S., Lejuez, C. W., Iza, M. and Blanco, C. (2013), 'Prospective study of substance-induced and independent major depressive disorder among individuals with substance use disorders in a nationally representative sample', *Depression and Anxiety* 30(6), 538-45.
- Magura, S. (2008), 'Effectiveness of dual focus mutual aid for co-occurring substance use and mental health disorders: a review and synthesis of the "double trouble" in recovery evaluation', *Substance Use & Misuse* 43(12-13), 1904-26.
- Magura, S., Rosenblum, A. and Betzler, T. (2009), 'Substance use and mental health outcomes for comorbid patients in psychiatric day treatment', *Substance Abuse: Research and Treatment* 3, 71-78.
- Maj, M. (2005), '"Psychiatric comorbidity": an artefact of current diagnostic systems?', *The British Journal of Psychiatry* 186, 182-84.
- Maremmani, A. G. I., Dell'Osso, L., Pacini, M., Popovic, D., Rovai, L., Torrens, M., Perugi, G. and Maremmani, I. (2011), 'Dual diagnosis and chronology of illness in treatment-seeking Italian patients dependent on heroin', *Journal of Addictive Diseases* 30(2), 123-35.
- Maremmani, A. G. I., Rovai, L., Rugani, F., Bacciardi, S., Dell'Osso, L. and Maremmani, I. (2014), 'Substance abuse and psychosis. The strange case of opioids', *European Review for Medical and Pharmacological Sciences* 18(3), 287-302.
- Maremmani, A. G. I., Rugani, F., Bacciardi, S., Rovai, L., Massimetti, E., Gazzarrini, D., Dell'Osso, L. and Maremmani, I. (2015), 'Differentiating between the course of illness in bipolar 1 and chronic-psychotic heroin-dependent patients at their first agonist opioid treatment', *Journal of Addictive Diseases* 34(1), 43-54.
- Marmorstein, N. R. (2011), 'Associations between subtypes of major depressive episodes and substance use disorders', *Psychiatry Research* 186(2-3), 248-53.
- Marmorstein, N. R. (2012), 'Anxiety disorders and substance use disorders: different associations by anxiety disorder', *Journal of Anxiety Disorders* 26(1), 88-94.
- Martín-Santos, R., Fonseca, F., Domingo-Salvany, A., Ginés, J. M., Ímaz, M. L., Navinés, R., Pascual, J. C. and Torrens, M. (2006), 'Dual diagnosis in the psychiatric emergency room in Spain', *The European Journal of Psychiatry* 20(3), 147-56.
- Martín-Santos, R., Torrens, M., Poudevida, S., Langohr, K., Cuyás, E., et al. (2010), '5-HTTLPR polymorphism, mood disorders and MDMA use in a 3-year follow-up study', *Addiction Biology* 15(1), 15-22.
- McCrone, P., Menezes, P. R., Johnson, S., Scott, H., Thornicroft, G., Marshall, J., Bebbington, P. and Kuipers, E. (2000), 'Service use and costs of people with dual diagnosis in South London', *Acta Psychiatrica Scandinavica* 101(6), 464-72.
- McKenzie, M. S. and McFarland, B. H. (2007), 'Trends in antidepressant overdoses', *Pharmacoepidemiology and Drug Safety* 16(5), 513-23.
- Mestre-Pintó, J. I., Domingo-Salvany, A., Martín-Santos, R. and Torrens, M. (2014), 'Dual diagnosis screening interview to identify psychiatric comorbidity in substance users: development and validation of a brief instrument', *European Addiction Research* 20(1), 41-48.

- Mills, K. L., Deady, M., Proudfoot, H., Sannibale, C., Teesson, M., Mattick, R. and Burns, L. (2009), *Guidelines on the management of co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings*, National Drug & Alcohol Research Centre, Sydney (k dispozici na <http://www.healthinfonet.ecu.edu.au/key-resources/promotion-resources?lid=17398>).
- Moeller, F. G., Barratt, E. S., Dougherty, D. M., Schmitz, J. M. and Swann, A. C. (2001), 'Psychiatric aspects of impulsivity', *The American Journal of Psychiatry* 158(11), 1783-93.
- Moggi, F., Giovanoli, A., Buri, C., Moos, B. S. and Moos, R. H. (2010), 'Patients with substance use and personality disorders: a comparison of patient characteristics, treatment process, and outcomes in Swiss and U.S. substance use disorder programs', *The American Journal of Drug and Alcohol Abuse* 36(1), 66-72.
- Morgello, S., Holzer, C. E., Ryan, E., Young, C., Naseer, M., et al. (2006), 'Interrater reliability of the Psychiatric Research Interview for Substance and Mental Disorders in an HIV-infected cohort: experience of the National NeuroAIDS Tissue Consortium', *International Journal of Methods in Psychiatric Research* 15(3), 131-38.
- Mueller, T. I., Lavori, P. W., Keller, M. B., Swartz, A., Warshaw, M., et al. (1994), 'Prognostic effect of the variable course of alcoholism on the 10-year course of depression', *The American Journal of Psychiatry* 151(5), 701-6.
- Neale, J., Tompkins, C. and Sheard, L. (2008), 'Barriers to accessing generic health and social care services: a qualitative study of injecting drug users', *Health & Social Care in the Community* 16(2), 147-54.
- Nelson, C. B., Rehm, J., Ustün, T. B., Grant, B. and Chatterji, S. (1999), 'Factor structures for DSM-IV substance disorder criteria endorsed by alcohol, cannabis, cocaine and opiate users: results from the WHO reliability and validity study', *Addiction* 94(6), 843-55.
- Ness, O., Borg, M. and Davidson, L. (2014), 'Facilitators and barriers in dual recovery: a literature review of first-person perspectives', *Advances in Dual Diagnosis* 7(3), 107-17.
- NICE (2011), *Psychosis with coexisting substance misuse: assessment and management in adults and young people*, National Institute of Health and Clinical Excellence, Manchester (k dispozici na <http://www.nice.org.uk/guidance/CG120>).
- Nocon, A., Bergé, D., Astals, M., Martín-Santos, R. and Torrens, M. (2007), 'Dual diagnosis in an inpatient drug-abuse detoxification unit', *European Addiction Research* 13(4), 192-200.
- Nordentoft, M., Mortensen, P. B. and Pedersen, C. B. (2011), 'Absolute risk of suicide after first hospital contact in mental disorder', *Archives of General Psychiatry* 68(10), 1058-64.
- Nunes, E. V and Levin, F. R. (2004), 'Treatment of depression in patients with alcohol or other drug dependence: a meta-analysis', *JAMA* 291(15), 1887-96.
- Nunes, E. V and Levin, F. R. (2006), 'Treating depression in substance abusers', *Current Psychiatry Reports* 8(5), 363-70.

- Nurnberger, J. I., Blehar, M. C., Kaufmann, C. A., York-Cooler, C., Simpson, S. G., et al. (1994), 'Diagnostic interview for genetic studies. Rationale, unique features, and training. NIMH Genetics Initiative', *Archives of General Psychiatry* 51(11), 849-59; discussion 863-64.
- O'Brien, M. S., Wu, L.-T. and Anthony, J. C. (2005), 'Cocaine use and the occurrence of panic attacks in the community: a case-crossover approach', *Substance Use & Misuse* 40(3), 285-97.
- OASAS (2005), *Screening for co-occurring disorders using the Modified Mini Screen (MMS) Provider Implementation Plan*, New York State Office of Alcoholism and Substance Abuse Services, Albany (k dispozici na <https://www.oasas.ny.gov/treatment/COD/documents/ModMINImpl.pdf>).
- Palijan, T. Z., Muzinić, L. and Radeljak, S. (2009), 'Psychiatric comorbidity in forensic psychiatry', *Psychiatria Danubina* 21(3), 429-36.
- Pani, P. P., Maremmanni, I., Trogu, E., Gessa, G. L., Ruiz, P. and Akiskal, H. S. (2010), 'Delineating the psychic structure of substance abuse and addictions: should anxiety, mood and impulse-control dysregulation be included?', *Journal of Affective Disorders* 122(3), 185-97.
- Pedrero-Perez, E. J., Ruiz-Sanchez de Leon, J. M., Rojo-Mota, G., Llanero-Luque, M. and Puerta-Garcia, C. (2011), '[Prevalence of attention-deficit hyperactivity disorder in substance addiction: from screening to diagnosis]', *Revista de Neurologia* 52(6), 331-40.
- Pérez Gálvez, B., García Fernández, L., de Vicente Manzanaro, M. P. and Oliveras Valenzuela, M. A. (2010), '[Validation of the Psychiatric Diagnostic Screening Questionnaire (PDSQ) in a Spanish sample of alcoholic patients]', *Adicciones* 22(3), 199-205.
- Pettinati, H. M., O'Brien, C. P. and Dundon, W. D. (2013), 'Current status of co-occurring mood and substance use disorders: a new therapeutic target', *The American Journal of Psychiatry* 170(1), 23-30.
- Philpot, M., Pearson, N., Petratou, V., Dayanandan, R., Silverman, M. and Marshall, J. (2003), 'Screening for problem drinking in older people referred to a mental health service: a comparison of CAGE and AUDIT', *Aging & Mental Health* 7(3), 171-75.
- Pierucci-Lagha, A., Gelernter, J., Feinn, R., Cubells, J. F., Pearson, D., Pollastri, A., Farrer, L. and Kranzler, H. R. (2005), 'Diagnostic reliability of the Semi-structured Assessment for Drug Dependence and Alcoholism (SSADDA)', *Drug and Alcohol Dependence* 80(3), 303-12.
- Pierucci-Lagha, A., Gelernter, J., Chan, G., Arias, A., Cubells, J. F., Farrer, L. and Kranzler, H. R. (2007), 'Reliability of DSM-IV diagnostic criteria using the semi-structured assessment for drug dependence and alcoholism (SSADDA)', *Drug and Alcohol Dependence* 91(1), 85-90.
- Pincus, H. A., Tew, J. D. and First, M. B. (2004), 'Psychiatric comorbidity: is more less?', *World Psychiatry* 3(1), 18-23.
- Piran, N. and Gadalla, T. (2007), 'Eating disorders and substance abuse in Canadian women: a national study', *Addiction* 102(1), 105-13.

- Piselli, M., Elisei, S., Murgia, N., Quartesan, R. and Abram, K. M. (2009), 'Co-occurring psychiatric and substance use disorders among male detainees in Italy', *International Journal of Law and Psychiatry* 32(2), 101-7.
- Pompili, M., Innamorati, M., Lester, D., Akiskal, H. S., Rihmer, Z., et al. (2009), 'Substance abuse, temperament and suicide risk: evidence from a case-control study', *Journal of Addictive Diseases* 28(1), 13-20.
- Radhakrishnan, R., Wilkinson, S. T. and D'Souza, D. C. (2014), 'Gone to pot: a review of the association between cannabis and psychosis', *Frontiers in Psychiatry* 5, 54.
- Ramos-Quiroga, J. A., Díaz-Digon, L., Comín, M., Bosch, R., Palomar, G., et al. (2012), 'Criteria and concurrent validity of adult ADHD section of the Psychiatry Research Interview for Substance and Mental Disorders', *Journal of Attention Disorders* doi:10.1177/1087054712454191.
- Rasmussen, K. and Levander, S. (2009), 'Untreated ADHD in adults: are there sex differences in symptoms, comorbidity, and impairment?', *Journal of Attention Disorders* 12(4), 353-60.
- Rhee, S. H., Hewitt, J. K., Lessem, J. M., Stallings, M. C., Corley, R. P. and Neale, M. C. (2004), 'The validity of the Neale and Kendler model-fitting approach in examining the etiology of comorbidity', *Behavior Genetics* 34(3), 251-65.
- Riglietta, M., Beato, E., Colombo, L. and Tidone, L. (2006), 'The Italian experience (A). Drugs and psychosis in Italy: historical overview and recent perspectives', in *Co-morbidity: Perspectives across Europe* edited by J. Corkery and A. Baldacchino, European Collaborating Centres on Addiction Studies (ECCAS), London, 165-85.
- Riper, H., Andersson, G., Hunter, S. B., de Wit, J., Berking, M. and Cuijpers, P. (2014), 'Treatment of comorbid alcohol use disorders and depression with cognitive-behavioural therapy and motivational interviewing: a meta-analysis', *Addiction* 109(3), 394-406.
- Robins, L. N., Helzer, J. E., Ratcliff, K. S. and Seyfried, W. (1982), 'Validity of the diagnostic interview schedule, version II: DSM-III diagnoses', *Psychological Medicine* 12(4), 855-70.
- Robins, L. N., Wing, J., Wittchen, H. U., Helzer, J. E., Babor, T. F., et al. (1988), 'The Composite International Diagnostic Interview: An epidemiologic instrument suitable for use in conjunction with different diagnostic systems and in different cultures', *Archives of General Psychiatry* 45(12), 1069-77.
- Robins, L. N., Cottler, L. B., Bucholz, K. K., Compton, W. M., North, C. S. and Rourke, K. M. (2000), *Diagnostic interview schedule for the DSM-IV (DIS-IV)*, Washington University School of Medicine, St. Louis.
- Rodríguez-Jiménez, R., Aragüés, M., Jiménez-Arriero, M. A., Ponce, G., Muñoz, A., Bagny, A., Hoenicka, J. and Palomo, T. (2008), '[Dual diagnosis in psychiatric inpatients: prevalence and general characteristics]', *Investigación Clínica* 49(2), 195-205.
- Rodríguez-Llera, M. C., Domingo-Salvany, A., Brugal, M. T., Silva, T. C., Sánchez-Niubó, A. and Torrens, M. (2006), 'Psychiatric comorbidity in young heroin users', *Drug and Alcohol Dependence* 84(1), 48-55.

- Rogers, R. (2001), *Handbook of diagnostic and structured interviewing*, Guilford Press, New York.
- Roncero, C., Ros-Cucurull, E., Daigre, C. and Casas, M. (2012), 'Prevalence and risk factors of psychotic symptoms in cocaine-dependent patients', *Actas Espanolas de Psiquiatria* 40(4), 187-97.
- Roncero, C., Daigre, C., Grau-López, L., Rodríguez-Cintas, L., Barral, C., et al. (2013), 'Cocaine-induced psychosis and impulsivity in cocaine-dependent patients', *Journal of Addictive Diseases* 32(3), 263-73.
- Root, T. L., Pisetsky, E. M., Thornton, L., Lichtenstein, P., Pedersen, N. L. and Bulik, C. M. (2010a), 'Patterns of co-morbidity of eating disorders and substance use in Swedish females', *Psychological Medicine* 40(1), 105-15.
- Root, T. L., Pinheiro, A. P., Thornton, L., Strober, M., Fernandez-Aranda, F., et al. (2010b), 'Substance use disorders in women with anorexia nervosa', *The International Journal of Eating Disorders* 43(1), 14-21.
- Rosenberg, S. D., Goodman, L. A., Osher, F. C., Swartz, M. S., Essock, S. M., et al. (2001), 'Prevalence of HIV, hepatitis B, and hepatitis C in people with severe mental illness', *American Journal of Public Health* 91(1), 31-37.
- Ruiz, M. A., Peters, R. H., Sanchez, G. M. and Bates, J. P. (2009), 'Psychometric properties of the Mental Health Screening form iii within a metropolitan jail', *Criminal Justice and Behavior* 36(6), 607-19.
- Sacks, S., Melnick, G., Coen, C., Banks, S., Friedmann, P. D., Grella, C., Knight, K. and Zlotnick, C. (2007), 'CJDATS Co-Occurring Disorders Screening Instrument (CODSI) for Mental Disorders (MD): A validation study', *Criminal Justice and Behavior* 34(9), 1198-1216.
- Sacks, S., Chaple, M., Sirikantraporn, J., Sacks, J. Y., Knickman, J. and Martinez, J. (2013), 'Improving the capability to provide integrated mental health and substance abuse services in a state system of outpatient care', *Journal of Substance Abuse Treatment* 44(5), 488-93.
- Salbach-Andrae, H., Lenz, K., Simmendinger, N., Klinkowski, N., Lehmkuhl, U. and Pfeiffer, E. (2008), 'Psychiatric comorbidities among female adolescents with anorexia nervosa', *Child Psychiatry and Human Development* 39(3), 261-72.
- Samet, S., Fenton, M. C., Nunes, E., Greenstein, E., Aharonovich, E. and Hasin, D. (2013), 'Effects of independent and substance-induced major depressive disorder on remission and relapse of alcohol, cocaine and heroin dependence', *Addiction* 108(1), 115-23.
- San, L., Arranz, B. and Martinez-Raga, J. (2007a), 'Antipsychotic drug treatment of schizophrenic patients with substance abuse disorders', *European Addiction Research* 13(4), 230-43.
- San, L., Ciudad, A., Alvarez, E., Bobes, J. and Gilaberte, I. (2007b), 'Symptomatic remission and social/vocational functioning in outpatients with schizophrenia: prevalence and associations in a cross-sectional study', *European Psychiatry* 22(8), 490-98.
- Sansone, R. A. and Levitt, J. L. (2002), 'Self-harm behaviors among those with eating disorders: an overview', *Eating Disorders* 10(3), 205-13.

- Sansone, R. A. and Sansone, L. A. (2010), 'Psychiatric disorders: a global look at facts and figures', *Psychiatry* 7(12), 16-19.
- Schier, A. R., Ribeiro, N. P., Silva, A. C., Hallak, J. E., Crippa, J. A., Nardi, A. E. and Zuardi, A. W. (2012), 'Cannabidiol, a Cannabis sativa constituent, as an anxiolytic drug', *Revista Brasileira de Psiquiatria* 34 Suppl 1, S104-10.
- Schippers, G. M., Broekman, T. G., Buchholz, A., Koeter, M. W. J. and van den Brink, W. (2010), 'Measurements in the Addictions for Triage and Evaluation (MATE): an instrument based on the World Health Organization family of international classifications', *Addiction* 105(5), 862-71.
- Schmidt, L. M., Hesse, M. and Lykke, J. (2011), 'The impact of substance use disorders on the course of schizophrenia - a 15-year follow-up study: dual diagnosis over 15 years', *Schizophrenia Research* 130(1-3), 228-33.
- Schmoll, S., Boyer, L., Henry, J.-M. and Belzeaux, R. (2015), '[Frequent visitors to psychiatric emergency service: Demographical and clinical analysis]', *L'Encéphale* 41(2), 123-29.
- Schnell, T., Neisius, K., Daumann, J. and Gouzoulis-Mayfrank, E. (2010), '[Prevalence of psychosis/substance abuse comorbidity. Clinical-epidemiological findings from different treatment settings in a large German city]', *Der Nervenarzt* 81(3), 323-28.
- Shahriyarmolki, K. and Meynen, T. (2014), 'Needs assessment of dual diagnosis: A cross-sectional survey using routine clinical data', *Drugs: Education, Prevention & Policy* 21(1), 43-49.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., et al. (1998), 'The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10', *The Journal of Clinical Psychiatry* 59 Suppl 2, 22-33; quiz 34-57.
- Sizoo, B., van den Brink, W., Koeter, M., Gorissen van Eenige, M., van Wijngaarden-Cremers, P. and van der Gaag, R. J. (2010), 'Treatment seeking adults with autism or ADHD and co-morbid substance use disorder: prevalence, risk factors and functional disability', *Drug and Alcohol Dependence* 107(1), 44-50.
- Smith, J. P. and Randall, C. L. (2012), 'Anxiety and alcohol use disorders: comorbidity and treatment considerations', *Alcohol Research: Current Reviews* 34(4), 414-31.
- Sorland, T. O. and Kjelsberg, E. (2009), 'Mental health among teenage boys remanded to prisoner', *Tidsskrift for Den Norske Laegeforening* 129(23), 2472-75.
- Soyka, M. (2000), 'Substance misuse, psychiatric disorder and violent and disturbed behaviour', *British Journal of Psychiatry* 176, 345-50.
- Spalletta, G., Bria, P. and Caltagirone, C. (2007), 'Differences in temperament, character and psychopathology among subjects with different patterns of cannabis use', *Psychopathology* 40(1), 29-34.
- Spitzer, R. L., Endicott, J. and Robins, E. (1978), 'Research diagnostic criteria: rationale and reliability', *Archives of General Psychiatry* 35(6), 773-82.
- Spitzer, R. L., Williams, J. B., Gibbon, M. and First, M. B. (1992), 'The Structured Clinical Interview for DSM-III-R (SCID). I: History, rationale, and description', *Archives of General Psychiatry* 49(8), 624-29.

- Spitzer, R. L., Kroenke, K. and Williams, J. B. (1999), 'Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. Primary care evaluation of mental disorders. Patient health questionnaire', *JAMA* 282(18), 1737-44.
- Stahler, G. J., Mennis, J., Cotlar, R. and Baron, D. A. (2009), 'The influence of neighborhood environment on treatment continuity and rehospitalization in dually diagnosed patients discharged from acute inpatient care', *American Journal of Psychiatry* 166(11), 1258-68.
- Substance Abuse and Mental Health Services Administration (2011) 'Results from the 2010 National Survey on Drug Use and Health: Summary of National Findings', NSDUH Series H-41, HHS Publication No. (SMA) 11-4658, Substance Abuse and Mental Health Services Administration, Rockville, MD.
- Swendsen, J. and Le Moal, M. (2011), 'Individual vulnerability to addiction', *Annals of the New York Academy of Sciences* 1216(1), 73-85.
- Szerman, N., Arias, F., Vega, P., Babín Vich, F., Mesías Perez, B., et al. (2011), '[Pilot study on the prevalence of dual pathology in community mental health and substance misuse services in Madrid]', *Adicciones* 23(3), 249-55.
- Szerman, N., Lopez-Castroman, J., Arias, F., Morant, C., Babín, F., et al. (2012), 'Dual diagnosis and suicide risk in a Spanish outpatient sample', *Substance Use & Misuse* 47(4), 383-89.
- Szerman, N., Vega, P., Grau-López, L., Barral, C., Basurte-Villamor, I., et al. (2014), 'Dual diagnosis resource needs in Spain: a national survey of professionals', *Journal of Dual Diagnosis* 10(2), 84-90.
- Thundiyil, J. G., Kearney, T. E. and Olson, K. R. (2007), 'Evolving epidemiology of drug-induced seizures reported to a poison control center system', *Journal of Medical Toxicology* 3(1), 15-19.
- Torrens, M., Serrano, D., Astals, M., Pérez-Domínguez, G. and Martín-Santos, R. (2004), 'Diagnosing comorbid psychiatric disorders in substance abusers: validity of the Spanish versions of the Psychiatric Research Interview for Substance and Mental Disorders and the Structured Clinical Interview for DSM-IV', *The American Journal of Psychiatry* 161(7), 1231-37.
- Torrens, M., Fonseca, F., Mateu, G. and Farre, M. (2005), 'Efficacy of antidepressants in substance use disorders with and without comorbid depression: A systematic review and meta-analysis', *Drug and Alcohol Dependence* 78(1), 1-22.
- Torrens, M., Martín-Santos, R. and Samet, S. (2006), 'Importance of clinical diagnoses for comorbidity studies in substance use disorders', *Neurotoxicity Research* 10(3-4), 253-61.
- Torrens, M., Martínez-Sanvisens, D., Martínez-Riera, R., Bulbena, A., Szerman, N. and Ruiz, P. (2011a), 'Dual diagnosis: Focusing on depression and recommendations for treatment', *Addictive Disorders & Their Treatment* 10(2), 50-59.
- Torrens, M., Gilchrist, G. and Domingo-Salvany, A. (2011b), 'Psychiatric comorbidity in illicit drug users: Substance-induced versus independent disorders', *Drug and Alcohol Dependence* 113(2-3), 147-56.

- Torrens, M., Rossi, P. C., Martinez-Riera, R., Martinez-Sanvisens, D. and Bulbena, A. (2012), 'Psychiatric co-morbidity and substance use disorders: Treatment in parallel systems or in one integrated system?', *Substance Use & Misuse* 47, 1005-14.
- Tosato, S., Lasalvia, A., Bonetto, C., Mazzoncini, R., Cristofalo, D., et al. (2013), 'The impact of cannabis use on age of onset and clinical characteristics in first-episode psychotic patients. Data from the Psychosis Incident Cohort Outcome Study (PICOS)', *Journal of Psychiatric Research* 47(4), 438-44.
- Toteva, S., Rizov, A., Tenev, V., Stoyanov, D. S. and Stoychev, K. (2006), 'The comorbidity situation in Bulgaria: epidemiological, clinical and therapeutic aspects', in *Co-morbidity: Perspectives across Europe* edited by J. Corkery and A. Baldacchino, European Collaborating Centres on Addiction Studies (ECCAS), London, 133-44.
- Valderas, J. M., Starfield, B., Sibbald, B., Salisbury, C. and Roland, M. (2009), 'Defining comorbidity: implications for understanding health and health services', *Annals of Family Medicine* 7(4), 357-63.
- Vallecillo, G., Mojal, S., Roquer, A., Martinez, D., Rossi, P., Fonseca, F., Muga, R. and Torrens, M. (2013), 'Risk of QTc prolongation in a cohort of opioid-dependent HIV-infected patients on methadone maintenance therapy', *Clinical Infectious Diseases* 57(8), 1189-94.
- Van de Glind, G., Konstenius, M., Koeter, M. W. J., van Emmerik-van Oortmerssen, K., Carpentier, P.-J., et al. (2014), 'Variability in the prevalence of adult ADHD in treatment seeking substance use disorder patients: results from an international multi-center study exploring DSM-IV and DSM-5 criteria', *Drug and Alcohol Dependence* 134, 158-66.
- Van den Bosch, L. M. and Verheul, R. (2007), 'Patients with addiction and personality disorder: Treatment outcomes and clinical implications', *Current Opinion in Psychiatry* 20(1), 67-71.
- Van Emmerik-van Oortmerssen, K., van de Glind, G., Koeter, M. W. J., Allsop, S., Auriacombe, M., et al. (2014), 'Psychiatric comorbidity in treatment-seeking substance use disorder patients with and without attention deficit hyperactivity disorder: results of the IASP study', *Addiction* 109(2), 262-72.
- Van Horn, J. E., Eisenberg, M. J., van Kuik, S. and van Kinderen, G. M. (2012), '[Psychopathology and recidivism among violent offenders with a dual diagnosis. A comparison with other subgroups of violent offenders]', *Tijdschrift Voor Psychiatrie* 54(6), 497-507.
- Vaz Carneiro, S. and Borrego, M. (2007), '[Psychosis and substance abuse]', *Acta Médica Portuguesa* 20(5), 413-22.
- Vázquez, C., Muñoz, M. and Sanz, J. (1997), 'Lifetime and 12-month prevalence of DSM-III-R mental disorders among the homeless in Madrid: a European study using the CID', *Acta Psychiatrica Scandinavica* 95(6), 523-30.
- Vázquez, F. L. (2010), 'Psychoactive substance use and dependence among Spanish university students: prevalence, correlates, polyconsumption, and comorbidity with depression', *Psychological Reports* 106(1), 297-313.

- Vázquez, F. L., Torres, Á., Otero, P. and Díaz, O. (2011), 'Prevalence, comorbidity, and correlates of DSM-IV axis I mental disorders among female university students', *The Journal of Nervous and Mental Disease* 199(6), 379-83.
- Vergara-Moragues, E., González-Saiz, F., Lozano, O. M., Betanzos Espinosa, P., Fernández Calderón, F., Bilbao-Acebos, I., Pérez García, M. and Verdejo García, A. (2012), 'Psychiatric comorbidity in cocaine users treated in therapeutic community: substance-induced versus independent disorders', *Psychiatry Research* 200(2-3), 734-41.
- Vorspan, F., Brousse, G., Bloch, V., Bellais, L., Romo, L., Guillem, E., Coeuru, P. and Lépine, J.-P. (2012), 'Cocaine-induced psychotic symptoms in French cocaine addicts', *Psychiatry Research* 200(2-3), 1074-76.
- Výroční zprávy národních monitorovacích středisek (nedatováno) (k dispozici na www.emcdda.europa.eu/publications/national-reports).
- Weaver, T., Madden, P., Charles, V., Stimson, G., Renton, A. et al. (2003) 'Comorbidity of substance misuse and mental illness in community mental health and substance misuse services', *The British Journal of Psychiatry* 183 (4), 304-13.
- Weissman, M. M., Olfson, M., Leon, A. C., Broadhead, W. E., Gilbert, T. T., et al. (1995), 'Brief diagnostic interviews (SDDS-PC) for multiple mental disorders in primary care. A pilot study', *Archives of Family Medicine* 4(3), 220-27.
- Whiteford, H. A., Degenhardt, L., Rehm, J., Baxter, A. J., Ferrari, A. J., et al. (2013), 'Global burden of disease attributable to mental and substance use disorders: Findings from the Global Burden of Disease Study 2010', *The Lancet* 382(9904), 1575-86.
- WHO (1990), *Composite International Diagnostic Interview (CIDI, Version 1.0)*, World Health Organization, Geneva.
- WHO (1992), *International statistical classification of diseases and related health problems*, World Health Organization, Geneva.
- WHO (1998), *Composite International Diagnostic Interview (CIDI Core, Version 2.1)*, WHO, Geneva.
- WHO (2010), *Lexicon of alcohol and drug terms published by the World Health Organization* (http://www.who.int/substance_abuse/terminology/who_lexicon/en/), zobrazeno 5. srpna 2015.
- WHO ASSIST (2002), 'The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): development, reliability and feasibility', *Addiction* 97(9), 1183-94.
- Wiederman, M. W. and Pryor, T. (1996), 'Substance use and impulsive behaviors among adolescents with eating disorders', *Addictive Behaviors* 21(2), 269-72.
- Wilens, T. E. and Biederman, J. (2006), 'Alcohol, drugs, and attention-deficit/hyperactivity disorder: a model for the study of addictions in youth', *Journal of Psychopharmacology* 20(4), 580-88.
- Wilens, T. E., Adamson, J., Monuteaux, M. C., Faraone, S. V., Schilling, M., Westerberg, D. and Biederman, J. (2008), 'Effect of prior stimulant treatment for attention-deficit/hyperactivity disorder on subsequent risk for cigarette smoking and alcohol and drug use disorders in adolescents', *Archives of Pediatrics & Adolescent Medicine* 162(10), 916-21.

- Wing, J. (1996), 'SCAN and the PSE tradition', *Social Psychiatry and Psychiatric Epidemiology* 31(2), 50-54.
- Wing, J. K., Babor, T., Brugha, T., Burke, J., Cooper, J. E., et al. (1990), 'SCAN. Schedules for Clinical Assessment in Neuropsychiatry', *Archives of General Psychiatry* 47(6), 589-93.
- Wittchen, H.-U., Fröhlich, C., Behrendt, S., Günther, A., Rehm, J., Zimmermann, P., Lieb, R. and Perkonig, A. (2007), 'Cannabis use and cannabis use disorders and their relationship to mental disorders: a 10-year prospective-longitudinal community study in adolescents', *Drug and Alcohol Dependence* 88 Suppl 1, S60-70.
- Wobrock, T. and Soyka, M. (2009), 'Pharmacotherapy of patients with schizophrenia and substance abuse', *Expert Opinion on Pharmacotherapy* 10(3), 353-67.
- IWPA (2014), *Dual disorders/pathology* http://www.wpanet.org/detail.php?section_id=11&content_id=1206, zobrazeno 5. srpna 2015.
- Wüsthoff, L. E., Waal, H., Ruud, T. and Grawe, R. W. (2011), 'A cross-sectional study of patients with and without substance use disorders in community mental health centres', *BMC Psychiatry* 11, 93.
- Wüsthoff, L. E., Waal, H. and Grawe, R. W. (2014), 'The effectiveness of integrated treatment in patients with substance use disorders co-occurring with anxiety and/or depression-a group randomized trial', *BMC Psychiatry* 14, 67.
- Zimmerman, M. and Chelminski, I. (2006), 'A scale to screen for DSM-IV Axis I disorders in psychiatric out-patients: performance of the Psychiatric Diagnostic Screening Questionnaire', *Psychological Medicine* 36(11), 1601-11.
- Zimmerman, M. and Mattia, J. I. (2001), 'The Psychiatric Diagnostic Screening Questionnaire: development, reliability and validity', *Comprehensive Psychiatry* 42(3), 175-89.
- Zimmerman, M., Sheeran, T., Chelminski, I. and Young, D. (2004), 'Screening for psychiatric disorders in outpatients with DSM-IV substance use disorders', *Journal of Substance Abuse Treatment* 26(3), 181-88.