

## Literatura:

1. Aitken, A., I. M. Mackie, J. H. Merritt a M. L. Windsor. (1982). Fish Handling and Processing. Aberdeen: Ministry of Agriculture, Fisheries, and Foods, Torry Research Station, Edinburgh, 123.
2. Akahane T., Chihara S., Niki T. P., Sano T., Tsuchiya T., Noguchi S. F., Ookami H. a Matsumoto J.J. (1985). Differentiation scanning calorimetric studies on thermal behaviors of myofibrillar proteins. Bull. Jap. Soc. Sci. Fish., 51, 1841-1846.
3. Ando M. Toyohara H., Shimizu Y., Sakaguchy M. (1991a) Post mortem tenderization of rainbow trout (*Oncorhynchus mykiss*) muscle caused by gradual desintegration of the extracellular matrix structure. J. Sci. Food Agric., 55, 489-597.
4. Ando M. Toyohara H., Shimizu Y., Sakaguchy M. (1991b) Post mortem tenderization of fish muscle proceed independently of resolution of rigor mortis. Nippon Suisan Gakkaishi, 57, 1165-1169.
5. Araki H., Seki N. (1993). Comparison of reactivity of transglutaminase to various fish actomyosins. Nippon Suisan Gakkaishi., 59, 1361-1364.
6. Berka R., Chábera S., (1977). Anglicko-český slovník z oboru rybářství, Výzkumný ústav rybářský a hydrobiologický Vodňany, 139.
7. Braekkan, O. R. (1976). Den ernaeringsmessige betydning av fisk. Fiskets Gang, 35.
8. Bykowski P., Dutkiewicz D., (1996). Zpracování sladkovodních ryb a vybavení malých zpracoven, Rybářské sdružení České Budějovice, 63.
9. Clucas, I. J. and Ward, A. R. (1996). Post – harvest Fisheries Development: A Guide to Handling, Preservation, Processing and Quality. Chatham Maritime, Kent ME4 4TB, United Kingdom, 254.
10. El-Fiky N., Wieser W. (1988). Life styles and patterns of development of gills and muscles in larval cyprinids (Cyprinidae; Teleostei). J. Fish. Biol., 33, 135-145.
11. Faria S. (1984). The Northeast Seafood Book, A Manual of Seafood Products, Marketing and Utilization, Massachusetts Division of Marine Fisheries, Boston, 315.
12. Graham J., Johnston W. A. a Nicholson F. J., (1993). Ice in fisheries, FAO Fish. Tech. Pap. 331.
13. Hatae K., Yoshimatsu F. a Matsumoto J. J. (1990). Role of muscle fibers in contributinfirmness of cooked fish. J. Food Sci., 55, 693-696.
14. Hatae K., Yoshimatsu F. a Matsumoto J. J. (1991). Effect of the preparation conditions on the texture of „Arai“ made of carp muscle. Nippon Suisa Gakkaishi, 56, 1521-1525.
15. Huss A., A., (1995). Quality and quality changes in fresh fish, FAO Fish. Tech. Pap. 348.
16. Huss H., H., (1994). Assurance of seafood quality, FAO Fish. Tech. Pap. 334.
17. Huss, H. H. (1988). Fresh fish. Quality and Quality Changes. FAO Fisheries Series No. 29.
18. Hwang G.C., Ushio H., Watabe S., Iwamoto M., Hashimoto K., (1991). The effect of thermal acclimation on rigor mortis progress of carp stored at different temperature. Nippon Suisan Gakkaishi, 57, 541-548.
19. Ingr I., (1994). Hodnocení a zpracování ryb, Vysoká škola zemědělská v Brně, 106.
20. Iwamoto, M., H. Yamanaka, S. Watabe a K. Hashimoto (1987). Effect of storage temperature on rigor-mortis and ATP degradation in plaice (*Paralichthys olivaceus*) muscle. J. Food Sci. 52, 6.

21. Kavina J., J. Winter, (1975). Drůbež – zvěřina – ryby - konzervy, 2. doplněné vydání, Merkur Praha, 62.
22. Kishi H., Nozawa H. a Seki N. (1991). Reactivity of muscle transglutaminase on carp myofibrils and myosin B. *Nippon Suisan Gakkaishi*, 57, 1203-1210.
23. Korhonen, R. W., T. C. Lanier a F. Giesbrecht (1990). An evaluation of simple methods for following rigor development in fish. *J. Food Sci.* 55, 2.
24. Love R. M. (1970). *The Chemical Biology of Fishes*. New York, Academic Press, 198.
25. Lundstrom, R. C. (1980). Fish species identification by thin layer polyacrylamide gel isoelectric focusing: Collaborative study. *J. Assoc. Off. Anal. Chem.* 63, 69-73.
26. Manikodan., Toyohara H., Ikeda S. (1984). Comparison of muscle proteinase activity among fish species. *Comp. Biochem. Physiol.*, 79B, 129, 134.
27. Martin, R. E. (1982). *Proceedings of the First National Conference on Seafood Packaging and Shipping*. Washington: National Fisheries Institute, 98.
28. Matyáš Z., (1993). *Analýza nebezpečí a kritické kontrolní / ochranné body*, Státní zdravotní ústav, centrum hygieny potravinových řetězců, Brno, 85.
29. Moustgard, J. (1957). *Laerebog i Husdyrenes Fysiologi og Ernoeringsfysiologi*, A/S C.Fr. Morstensen, Copenhagen, 272.
30. Murray J. , J. R. Burt (1969). The composition of fish. *Torry Advis. Note* 38, Torry Research Station, Aberdeen, 38.
31. Nakayama T., Liu D. J., Ooi A. (1992). Tension changes of stressed and unstressed carp muscle in isometric rigor contraction and resolution. *Nippon Suisan Gakkaishi*, 58, 1517-1522.
32. Nakayama, T., D.-J. Liu a A. Ooi (1992). Tension change of stressed and unstressed carp muscles in isometric rigor contraction and resolution. *Nippon Suisan Gakkaishi*, 58, 8.
33. Pavlova L. A. (1992). Some specific features of the thermotropic gelation of carp actomyosin. *Nahrung*, 36, 551-558.
34. Pokorný J. (1993). *Metody senzorické analýzy potravin a stanovení senzorické jakosti*, Ústav zemědělských a potravinářských informací, 196.
35. Poulter, N. H. , Nicolaidis (1985). Studies of the iced storage characteristics and composition of a variety of Bolivian freshwater fish. *Altiplano fish. J. Food Technol.* 20, 437 – 449.
36. Regenstien J. M., Regenstien C. E., (1991). *Introduction to fish technology*, Van Nostrand Reinhold, 269.
37. Roerbaek, K., B. Jensen a K. Mathiasen (1993). Oxidation and aroma in fish oil. In: G. Lambertsen (ed.) *Proceedings of the 17<sup>th</sup> Nordic symposium on lipids*, Imatra, Sf. Lipidforum, Bergen, Norway, 175.
38. Romanovský A. a kol. (1985). *Obecná biologie*, Státní pedagogické nakladatelství Praha, 695.
39. Saito, T., K. Arai. A M. Matsuyoshi (1959). A new method for estimating the freshness of fish. *Bull. Jap. Soc. Sci. Fish.* 24, 749-50.
40. Sano T., Noguchi S. F., Matsumoto J. J. a Tsuchiya T. (1990a). Effect of ionic strength on dynamic viscoelastic behavior of myosin during thermal gelation. *J. Food Sci.*, 55, 51-54.
41. Sano T., Noguchi S. F., Matsumoto J. J. a Tsuchiya T. (1990b). Thermal gelation characteristics of myosin subfragments. *J. Food Sci.*, 55, 55-58.
42. Sczerbowski, J. A., (1995). *Inland fisheries in Poland*, Instytut rybactwa śródladowego Olsztyn, 544.

43. Seki N., Tsuchiya H. (1991). Extensive changes during storage in carp myofibrillar proteins in relation to fragmentation. *Nippon Suisan Gakkaishi*, 57, 927-933.
44. Shewan, J. M. (1974). The biodeterioration of certain proteinaceous foodstuffs at chill temperatures. In: B. Spencer (ed.), *Industrial aspects of biochemistry*, 475-490.
45. Stansby, M. E. (1962). Proximate composition of fish. In E. Heen, R. Kreuzer (ed.) *Fish in nutrition*. Fishing News Books Ltd., London, 55 – 60.
46. Stroem, A.R., J.A. Olafsen a H. Larsen (1979). Trimethylamine oxide: a terminal electron acceptor in aerobic respiration of bacteria. *J. Gen. Microbiol.*, 112, 315-20.
47. Tokiwa, T., Matsumiya, H. (1969). Fragmentation of fish myofibril. Effect of storage condition and muscle cathepsin. *Bull. Jap. Soc. Sci. Fish.*, 35, 1099-1109.
48. Trucco, R. E., H. M. Lupin, D. H. Gianini, M. Grupkin, R. L. Beori, a C.A. Barassi (1982). Study on the evolution of rigor mortis in batches of fish. *Lebensm. – Wiss. & Technol.* 15, 77-79.
49. Tsuchiya H., Kita S., Seki N. (1992). Postmortem changes in  $\alpha$ -actinin and connectin in carp and rainbow trout muscles. *Nippon Suisan Gakkaishi*, 58, 793-798.
50. Vácha F., Prošková A., Kučera J., (1998), Aminokyselinové složení vybraných druhů sladkovodních ryb, Sborník Jih. univ., ZF v Českých Budějovicích, zootechnická řada, ISBN 80-7040-316-0, XV, 2, 79-89.
51. Vácha F., Tvrzická E., (1997), Polyunsaturated fatty acids content in fat of tench (*Tinca tinca*) under different rearing conditions. Ve: II International Workshop on Biology and Culture of the Tench, Badajoz, (Spain), 12.