

References

1. Strader, C. D., Fong, T. M., Graziano, M. P., and Tota, M. R. (1995) The family of G-protein-coupled receptors. *FASEB J.* **9**, 745–754.
2. Rohrer, D. K. and Kobilka, B. K. (1998) Insights from in vivo modification of adrenergic receptor gene expression. *Annu. Rev. Pharmacol. Toxicol.* **38**, 351–373.
3. Aoki, C., Go, C. G., Venkatesan, C., and Kurose, H. (1994) Perikaryal and synaptic localization of α_{2A} -adrenergic receptor immunoreactivity in brain as revealed by light and electron microscopic immunocytochemistry. *Brain Res.* **650**, 181–204.
4. Kalsner, S. and Westfall, T. C. (1990) Presynaptic receptors and the question of autoregulation of neurotransmitter release, *Ann. NY Acad. Sci.* **604**, 652.
5. Venkatesan, C., Kurose, H., and Aoki, C. (1996) Cellular and subcellular distribution of α_{2A} - adrenergic receptor in the visual cortex of neonatal and adult rats. *J. Comp. Neurol.* **365**, 79–95.
6. Aoki, C. (1998) Cellular and subcellular sites for noradrenergic action in the monkey dorsolateral prefrontal cortex as revealed by the immunocytochemical localization of noradrenergic receptors and axons. *Cereb. Cortex* **8**, 269–277.
7. Milner, T. A., Lee, A., Aicher, S., and Rosin, D. L. (1998) Hippocampal α_{2C} adrenergic receptors are located predominantly presynaptically but are also found postsynaptically and in selective astrocytes. *J. Comp. Neurol.* **395**, 310–327.
8. McCormick, D. A., Pape, H. C., and Williamson, A. (1991) Actions of norepi-

- nephrine in the cerebral cortex and thalamus: implications for function of the central noradrenergic system. *Prog. Brain Res.* **88**, 293–305.
9. Strader, C. D., Picke, V. M., Joh, T. H., Strohsacker, M. W., Shorr, Lefkowitz, R. J., and et al. (1983) Antibodies to the beta-adrenergic receptor: attenuation of catecholamine-sensitive adenylate cyclase and demonstration of postsynaptic receptor localization in brain, *Proc. Natl. Acad. Sci USA* **80**, 1840–1844.
 10. Aoki, T. H., Joh, and Pickel, V. M. (1987) Ultrastructural localization of immunoreactivity for β -adrenergic receptors in the cortex and neostriatum of rat brain. *Brain Res.* **437**, 264–282.
 1. Aoki, C., Zemcik, Z. A., Strader, C. D., and Pickel, V. M. (1989) Cytoplasmic loop of β -adrenergic receptors: synaptic and intracellular localization and relation to catecholaminergic neurons in the nuclei of the solitary tracts. *Brain Res.* **493**, 331–347.
 2. Aoki, C. and Pickel, V. M. (1990) Ultrastructural immunocytochemical evidence for presynaptic localization of beta-adrenergic receptors in the striatum and cerebral cortex of rat brain. *Ann NY Acad Sci.* **604**, 582–585.
 3. Aoki, C. (1992). C-terminal fragment of β -adrenergic receptors: astrocytic localization in the visual cortex and their relation to catecholamine axon terminals, *J. Neurosci.* **12**, 781–792.
 4. Aoki, C. and Pickel, V. M. (1992) Ultrastructural relations between β -adrenergic receptors and catecholaminergic neurons, *Brain Res. Bull.* **29**, 257–264.
 5. Aoki, C. and Pickel, V. M. (1992) C-terminal tail of beta-adrenergic receptors: immunocytochemical localization within astrocytes and their relation to catecholaminergic neurons in the N. tractus solitarii and area postrema, *Brain Res.* **571**, 35–49.
 6. Aoki, C., Lubin, M., and Fenstermaker, F. (1994) Columnar activity regulates astrocytic β -adrenergic receptor-like immunoreactivity in V1 of adult monkeys. *Vis. Neurosci.* **11**, 179–187.
 7. Aoki, C. (1997) Differential timing for the appearance of neuronal and astrocytic beta-adrenergic receptors in the developing rat visual cortex as revealed by light and electron-microscopic immunocytochemistry. *Vis. Neurosci.* **14**, 1129–1142.
 8. Susulic, V. S., Frederich, R. C., Lawitts, J., Tozzo, E., Kahn, B. B., Harper, M.-E., et al. (1995) Targeted disruption of the β_3 -adrenergic receptor gene. *J. Biol. Chem.* **270**, 29,483–29,492.
 9. Rohrer, D. K., Desai, K. H., Jasper, J. R., Stevens, M. E., Regula, D. P., Jr., Barsh, G. S., et al. (1996) Targeted disruption of the mouse beta1-adrenergic receptor gene: developmental and cardiovascular effects. *Proc. Natl. Acad. Sci. USA* **93**, 7375–7380.
 10. Jurevics, J. and Fischmeister, R. (1996) cAMP compartmentation is responsible for a local activation of cardiac Ca^{2+} channel by β -adrenergic agonists. *Proc. Natl. Acad. Sci. USA* **93**, 295–299.
 1. von Zastrow, M., Link, R., Daunt, D., Barsh, G., and Kobilka, B. K. (1993) Subtype-specific differences in the intracellular sorting of G-protein-coupled receptors. *J. Biol. Chem.* **268**, 763–766.

22. Hirasawa, A., Sugawara, T., Awaji, T., Tsumaya, K., Ito, H., and Tsujimoto, G. (1998) Subtype-specific differences in subcellular localization and chlorethylclonidine (CEC) inactivation of α_1 -adrenoceptors (ARs): CEC alkylates only the accessible cell surface α_1 -ARs irrespective of the subtypes. *Mol. Pharmacol.* **52**, 764–770.
23. Fuxe, K. and Agnati, L. F. (eds.) *Volume Transmission in the Brain: Novel Mechanisms for Neural Transmission*. Raven, New York.
24. King, L. C., Lechan, R. M., Kugel, G., and Anthony, E. L. P. (1983) Acrolein: a fixative for immunocytochemical localization of peptides in the central nervous system, *J. Histochem. Cytochem.* **31**, 62–68.
25. Sitte, H. (1996) Advanced instrumentation and methodology related to cryoultramicrotomy: a review. *Scanning Microsc.* **10 (Suppl.)**, 387–463.
26. Liou, W., Geuze, H. J., and Slot, J. W. (1996) Improving structural integrity of cryosections for immunogold labeling. *Histochem. Cell Biol.* **106**, 41–58.
27. Eldred, W. D., Zucker, C., Karten, H. J., and Yazula, S. (1983) Comparison of fixation and penetration enhancement techniques for use in ultrastructural immunocytochemistry, *J. Histochem. Cytochem.* **31**, 285–292.
28. Aoki, C., Starr, A., Kaneko, T., and Pickel, V. M. (1991) Identification of mitochondrial and non-mitochondrial glutaminase within select neurons and glia of rat forebrain by electron microscopic immunocytochemistry. *J. Neurosci. Res.* **28**, 531–548.
29. Erisir, A. and Aoki, C. (1998) Combined use of biocytin with avidin–biotin peroxidase for dual pre-embedding electron microscopy. *J. Neurosci. Methods* **81**, 189–197.
30. Sesack, S. R., Aoki, C., and Pickel, V. M. (1994) Ultrastructural localization of D2 receptor-like immunoreactivity in midbrain dopamine neurons and their striatal targets. *J. Neurosci.* **14**, 88–106.
31. Wouterlood, F. G. and Jorritsma-Byham, B. (1993) The anterograde neuroanatomical tracer biotinylated dextran-amine: comparison with the tracer Phaseolus vulgaris-leucoagglutinin in preparations for electron microscopy. *J. Neurosci. Methods* **48**, 75–87.
32. Baude, A., Molnar, E., Latawiec, D., McIlhinney, R. A. J., and Somogyi, P. (1994) Synaptic and nonsynaptic localization of the GluR1 subunit of the AMPA-type excitatory amino acid receptor in the rat cerebellum. *J. Neurosci.* **14**, 2830–2843.
33. Bernard, V., Somogyi, P., and Bolam, J. P. (1997) Cellular, subcellular and subsynaptic distribution of AMPA-type glutamate receptor subunits in the neostriatum of the rat. *J. Neurosci.* **17**, 819–833.
34. Raman I. M., Tong, G., and Jahr, C. E. (1996) Beta-adrenergic regulation of synaptic NMDA receptors by cAMP-dependent protein kinase. *Neuron* **16**, 415–421.
35. He, Y., Janssen, W. G. M., Vissavajhala, P., and Morrison, J. H. (1998) Synaptic distribution of GluR2 in hippocampal GABAergic interneurons and pyramidal cells: a double-label immunogold analysis. *Exp. Neurol.* **150**, 1–13.

36. Kharazia, V. N., Phend, K. D., Rustioni, A., and Weinberg, R. J. (1996) EM localization of AMPA and NMDA receptor subunits at synapses in rat cerebral cortex. *Neurosci Lett.* **210**, 37–40.
37. Phend, K. D., Rustioni, A., and Weinberg, R. J. (1995) An osmium-free method of Epon embedding that preserves both ultrastructure and antigenicity for post-embedding immunocytochemistry. *J. Histochem. Cytochem.* **43**, 283–292.
38. Rubio, M. E. and Wenthold, R. J. (1997) Glutamate receptors are selectively targeted to postsynaptic sites in neurons. *Neuron* **18**, 939–950.
39. Wang, Y.-X., Wenthold, R. J., Ottersen, O. P., and Petralia, R. S. (1998) Endbulb synapses in the anteroventral cochlear nucleus express a specific subset of AMPA-type glutamate receptor subunits. *J. Neurosci.* **18**, 1148–1160.
40. Veznedaroglu, E. and Milner, T. A. (1992) Elimination of artifactual labeling of hippocampal mossy fibers seen following preembedding immunogold-silver technique by pretreatment with zinc chelator. *J. Microsc. Res. Tech.* **23**, 100, 101.
41. Yasukawa, T., Kanei-Ishii, C., Maekawa, T., Fujimoto, J., Yamamoto, T., and Ishii, S. (1995) Increase of solubility of foreign proteins in *Escherichia coli* by coproduction of the bacterial thioredoxin. *J. Biol. Chem.* **270**, 25,328–25,331.
42. LaVallie, E. R., DiBlasio, E. A., Kovacic, S., Grant, K. L., Schendel, P. F., and McCoy, J. M. (1993) A thioredoxin gene fusion expression system that circumvents inclusion body formation in the *E. coli* cytoplasm. *Biotechnology* **11**, 187–193.
43. Frangioni, J. V. and Neel, B. G. (1993) Solubilization and purification of enzymatically active glutathione *S*-transferase (pGEX) fusion proteins. *Anal. Biochem.* **210**, 179–187.
44. Pickel, V. M., Chan, J., and Aoki, C. (1993) Electron microscopic immunocytochemical labeling of endogenous and/or transported antigens in rat brain using silver-intensified one-nanometre colloidal gold, in *Immunohistochemistry II* (Cuello, A. C., ed.) John Wiley, New York, pp. 265–280.