

CONTENTS

The interaction of fish trypanosome culture forms with some lectins P. ZAJÍČEK, H. PECKOVÁ	1
<i>Entamoeba histolytica</i> — the virulence of imoorted strains M. GIBODA, O. DITRICH, N. VRCHOTOVÁ, J. GUTVIRTH, P. NOLL, J. ŠTĚRBA	9
Double alteration of the small intestine in conventional and gnotobiotic, piglets experimentally infected with the coccidium <i>Isospora suis</i> (Apicomplexa, Eimeriidae) J. VÍTOVEC, B. KOUDELA	21
Some remarks on the prevalence and species composition of hare coccidia M. PAKANDL	35
Additional records of nematode parasites from papua New Guinea amphibians with a list of recorded endohelminths by amphibian hosts F. MORAVEC	43
Influence of temperature on reproduction and survival of <i>Pseudodactylogyrus anguillae</i> (Monogenea) from the European eel K. BUCHMANN	59
Electrophoretic analysis of gene-enzyme systems in <i>Chabertia ovina</i> J. E. ORTEGA, M. SANCHEZ-MORENO, A. FATOU, A. VALERO	63
A new genus of capillariids form birds, <i>Tridenticapillaria</i> gen. n. (Nematoda: Capillariidae) V. BARUŠ, T. P. SERGEEVA	67
Visualization of chitin-protein layer formation on <i>Ascaris lumbricoides</i> egg-shells D. ŠROMOVÁ, H. LÝSEK	77
Three species of the hair follicle mites (Acari: Demodicidae) parasitizing the sheep, <i>Ovis aries</i> L. V. BUKVA	81
Notes	
Synonymy of <i>Paraquadracanthus</i> Ergens, 1988 and <i>Quadracanthoides</i> Kritsky et Kulo, 1988 (Monogenea, Dactylogyridae) and their type species D. C. KRITSKY	76
Occurence of the tick <i>Ixodes laguri</i> Ol. on the territory of Czechoslovakia V. ČERNÝ	92
First record of <i>Avioserpens</i> larvae (Nematoda) from the naturally infected intermediate host F. MORAVEC, T. SCHOLZ	93
First record of mite hypopi (Acaridae) on fleas from Czechoslovakia J. PREISLER, K. SAMŠIŇÁK, V. ČERNÝ	94
Book reviews	8, 8, 19, 20, 34, 42, 62, 80, 95, 96