

CONTENTS

Amoebae isolated from organs of farmed tilapias, <i>Oreochromis niloticus</i> I. Dyková, B. Macháčková, H. Pecková	81 - 90
Development of sporozoites of the piscine coccidium <i>Eimeria (sensu lato) vanasi</i> in gut intraepithelial lymphocyte-like cell M. Vilenkin, I. Paperna	91 - 98
<i>Vipera berus</i> and <i>V. ammodytes</i> (Serpentes: Viperidae) represent new hosts for <i>Caryospora simplex</i> (Apicomplexa: Eimeriidae) in Europe D. Modrý, B. Koudela, J. Volf, P. Nečas, T. Hudcovic	99 - 102
Isolation and characterization of cytosolic malate dehydrogenase from <i>Trichomonas vaginalis</i> T. Drmota, J. Tachezy, J. Kulda	103 - 108
Larval stages of trematodes from freshwater molluscs the Yucatan Peninsula, Mexico O. Ditrich, T. Scholz, L. Aguirre-Macedo, J. Vargas-Vázquez	109 - 127
Electrophoretic variations of peptidase E (PEPE) in characterizing clones and isolates of <i>Plasmodium falciparum</i> from different geographical areas H.H. Ang, K. Lam Chan, J. Wah Mak	128 - 130
<i>Khawia baltica</i> (Cestoidea: Caryophyllidea) from <i>Barbus barbus bocagei</i> from some rivers in Northern Portugal J.C. Chubb, J.C. Eiras, A. Saraiva	131 - 138
Ultrastructure of three types of scolex gland cells in adult <i>Bothriocephalus claviceps</i> (Cestoda: Pseudophyllidea) Z. Žďárská, J. Nebesářová	139 - 146
Estimation of genetic affinities between sympatric populations of <i>Rhipicephalus pusillus</i> ticks (Acari: Ixodidae) by analysis of cuticular hydrocarbons A. Estrada-Peña, J.J. Osácar, C. Calvete, R. Estrada-Peña	147 - 154
Role of grey squirrels and pheasants in the transmission of <i>Borrelia burgdorferi sensu lato</i> , the Lyme disease spirochaete, in the U.K. N.G. Craine, P.A. Nuttall, A.C. Marriott, S.E. Randolph	155 - 160