

CONTENTS

Lectin-like sequences in genome of <i>Borrelia burgdorferi</i> N. Rudenko, M. Golovchenko, L. Grubhoffer	81-90
Cytokine response to infection with the microsporidian, <i>Encephalitozoon cuniculi</i> P. Braunfuchsová, J. Kopecký, O. Ditrich, B. Koudela	91-95
<i>Caryospora maxima</i> sp. n. (Apicomplexa: Eimeriidae), new coccidium from Forskal's sand snake, <i>Psammophis schokari</i> (Serpentes: Colubridae) from lava desert of Jordan D. Modrý, B. Koudela, R.M. Al-Oran, Z.S. Amr	96-98
Extraintestinal stages of coccidia in liver of Schneider's skink <i>Eumeces schneideri</i> (Sauria: Scincidae) from Northern Egypt B. Koudela, D. Modrý	99-102
Identification of surface proteins of trophic and cyst forms of <i>Blastocystis hominis</i> by biotin labeling J. Villar, J.-A. Carbajal, M.-D. Lanuza, M.-R. Navarro, R. Borrás	103-106
Evidence of host blood feeding by the monogenean, <i>Ancyrocephalus mogurndae</i> (Monogenea: Ancyrocephalidae) from the gills of the mandarin fish, <i>Siniperca chuatsi</i> Q. Gao, P. Nie, W.J. Yao	107-110
Post-cyclic transmission in <i>Pomphorhynchus laevis</i> (Acanthocephala) C.R. Kennedy	111-116
Amphipod intermediate host of <i>Polymorphus minutus</i> (Acanthocephala), parasite of water birds, with notes on ultrastructure of host-parasite interface B.S. Dezfuli, L. Giari	117-122
Morphology and taxonomy of <i>Salvelinema</i> species (Nematoda: Cystidicolidae), swimbladder parasites of Pacific area salmonids F. Moravec, K. Nagasawa	123-131
Digenetic trematodes parasitic on <i>Engraulis anchoita</i> (Pisces: Engraulidae) from Argentina and Uruguay J.T. Timi, S.R. Martorelli, N.H. Sardella	132-138
First record of the genus <i>Syncuaria</i> (Nematoda: Acuariidae) in Argentina, with description of a new species M.C. Digiani	139-144
Survey of entomopathogenic nematodes from the families Steinernematidae and Heterorhabditidae (Nematoda: Rhabditida) in the Czech Republic Z. Mráček, S. Bečvář, P. Kindlmann	145-148
Parasitic copepods of the narrownose smooth-hound shark <i>Mustelus schmitti</i> (Chondrichthyes: Triakidae) from Argentina J.A. Etchegoin, V.A. Ivanov	149-153
Notes	
Exoerythrocytic stage of <i>Haemoproteus pyodactyli</i> in the spleen of its host, the gecko <i>Ptyodactylus hasselquistii</i> I. Paperna, S. Finkelman	154
Parasitophorous vacuole of intranuclear coccidia maintains membranous junctions with the host-cell cytoplasm I. Paperna, M. Vilenkin	155-156
Encystment sites of metacercariae of <i>Hypoderaeum conoideum</i> (Trematoda: Echinostomatidae) in freshwater gastropods R. Toledo, C. Muñoz-Antolí, M. Pérez, J.-G. Esteban	157-158
<i>Comephoronema oschmarini</i> (Nematoda: Cystidicolidae), a specific parasite of <i>Lota lota</i> (Pisces), occurs in western Europe F. Moravec, V. Hanzelová, T. Scholz, D. Gerdeaux	159-160