

CONTENTS

ARTICLES

- 545 Limited variation in mitochondrial DNA of maize-associated *Ostrinia nubilalis* (Lepidoptera: Crambidae) in Russia, Turkey and Slovenia — S. HOSHIZAKI, R. WASHOMORI, S. KUBOTA, A.N. FROLOV, D. KAGEYAMA, S. GOMBOC, S. OHNO, S. TATSUKI & Y. ISHIKAWA
- 553 The genetic architecture of extended life span in the seed beetle *Acanthoscelides obtectus* (Coleoptera: Bruchidae) — D. ŠEŠLIJA & N. TUCIĆ
- 561 Leucine-rich fibroin gene of the Japanese wild silkworm, *Rhodinia fugax* (Lepidoptera: Saturniidae) — H. SEZUTSU, T. TAMURA & K. YUKUHIRO
- 567 Glutathione S-transferases from the larval gut of the silkworm *Bombyx mori*: cDNA cloning, gene structure, expression and distribution — Z.Z. GUI, B.Y. KIM, K.S. LEE, Y.D. WEI, X. GUO, H.D. SOHN & B.R. JIN
- 575 Regulation of the development of flight muscles in long-winged adults of the flightless bug *Pyrrhocoris apterus* (Heteroptera: Pyrrhocoridae) — R. SOCHA & J. ŠULA
- 585 Light-induced behavioural effects on the locomotor activity rhythm of the blow fly, *Calliphora vicina* (Diptera: Calliphoridae) — D.S. SAUNDERS & B. CYMBOROWSKI
- 591 Endogenous 20-hydroxyecdysone levels in the haemolymph of non-diapause-destined and diapause-destined generations of tasar silkworm, *Antheraea mylitta* (Lepidoptera: Saturniidae) and associated developmental changes — D. KUMAR, B. SUBRAHMANYAM, S.K. SHARAN, P.K. MISHRA, B.M.K. SINGH & N. SURYANARAYANA
- 599 The effect of temperature on the diapause and cold hardiness of *Dendrolimus tabulaeformis* (Lepidoptera: Lasiocampidae) — J.-P. ZENG, F. GE, J.-W. SU & Y. WANG
- 607 Influence of pre-diapause temperature on intensity of summer and winter diapause in the cabbage butterfly *Pieris melete* (Lepidoptera: Pieridae) — H.-J. XIAO, H.-M. HE, F. LI & F.-S. XUE
- 613 Role of volatile and contact pheromones in the mating behaviour of *Bagrada hilaris* (Heteroptera: Pentatomidae) — S. GUARIANO, C. DE PASQUALE, E. PERI, G. ALONZO & S. COLAZZA
- 619 Role of excreta in predator avoidance by the Kanzawa spider mite, *Tetranychus kanzawai* (Acari: Tetranychidae) — K. OKU
- 625 Effect of larval crowding on the life history traits of *Sesamia nonagrioides* (Lepidoptera: Noctuidae) — A.A. FANTINOU, D.C. PERDIKIS & N. STAMOGIANNIS
- 631 Substrate particle size-preference of wormlion *Vermileo vermileo* (Diptera: Vermileonidae) larvae and their interaction with ant-lions — D. DEVETAK
- 637 *Myrmica sabuleti* (Hymenoptera: Formicidae) not necessary for the survival of the population of *Phengaris (Maculinea) arion* (Lepidoptera: Lycaenidae) in eastern Poland: Lower host-ant specificity or evidence for geographical variation of an endangered social parasite? — M. SIELEZNIEW & A.M. STANKIEWICZ
- 643 Experimental evidence for competitive exclusion of *Myzus persicae nicotianae* by *Myzus persicae* s.s. (Hemiptera: Aphididae) on sweet pepper, *Capsicum annuum* (Solanaceae) — D.H. TAPIA, A.J. TRONCOSO, R.R. VARGAS, R. OLIVARES-DONOSO & H.M. NIEMAYER
- 649 Resources for British butterflies (Lepidoptera: Hesperioida, Papilionoidea). The alien consumer component and its significance for butterfly habitats — P.B. HARDY & R.L.H. DENNIS
- 659 The classification of insect communities: Lessons from orthopteran assemblages of semi-dry calcareous grasslands in central Germany — D. PONIATOWSKI & T. FARTMANN
- 673 Orthopteran communities in the conifer-broadleaved woodland zone of the Russian Far East — T. FARTMANN, M. BEHRENS & H. LORITZ
- 681 Establishment of ant communities in forests growing on former agricultural fields: Colonisation and 25 years of management are not enough (Hymenoptera: Formicidae) — W. DEKONINCK, K. DESENDER & P. GROOTAERT
- 691 Insects on decomposing carcasses of small rodents in a secondary forest in Southeastern Brazil — T. DE C. MORETTI, O.B. RIBEIRO, P.J. THYSSEN & D.R. SOLIS
- 697 The developmental process during metamorphosis that results in wing reduction in females of three species of wingless-legged bagworm moths, *Taleporia trichopterella*, *Bacotia sakabei* and *Proutia* sp. (Lepidoptera: Psychidae) — S. NIITSU & Y. KOBAYASHI
- 707 Morphology and ultrastructure of the male reproductive system of the woolly beech aphid *Phyllaphis fagi* (Hemiptera: Aphididae: Phylloxerinae) — K. WIECZOREK & P. ŚWIĄTEK
- 713 Rapid discrimination of the common species of the stored product pest *Liposcelis* (Psocoptera: Liposcelididae) from China and the Czech Republic, based on PCR-RFLP analysis — M. QIN, Z.-H. LI, Z. KUČEROVÁ, Y. CAO & V. STEJSKAL
- 719 Taxonomical revision of the *Longitarsus capensis* species-group: An example of Mediterranean-southern African disjunct distributions (Coleoptera: Chrysomelidae) — M. BIONDI & P. D'ALESSANDRO
- 737 On the systematic position of the diving-beetle genus *Pachydrus* (Coleoptera: Dytiscidae: Hydrocorinae): Evidence from larval chaetotaxy and morphology — M.C. MICHAUT & P.L.M. TORRES
- 751 Revision of *Xyalophora* Kieffer and description of *Xyalophoroides* gen. n. (Hymenoptera: Figitidae: Figitinae) — M. JIMÉNEZ, J. PARETAS-MARTÍNEZ & J. PUJADE-VILLAR
- 771 Revision of *Phaeochiton* Kerzhner, 1964 (Heteroptera: Miridae: Phylini) — F.V. KONSTANTINOV
- 783 A new genus of isophlebioid damsel-dragonflies with "calopterygid"-like wing shape from the Middle Jurassic of China (Odonata: Isophlebiidae: Camptophlebiidae) — A. NEL, D.-Y. HUANG & Q.-B. LIN
- NOTES
- 789 *Polygraphus grandiclava* (Coleoptera: Curculionidae) collected from pine and cherry trees: A phylogenetic analysis — D. AVTZIS, M. KNÍŽEK, K. HELLERIGL & C. STAUFFER
- 793 A modified method for clearing, staining and mounting plant-inhabiting mites — F. FARAJI & F. BAKKER