

CONTENTS

ARTICLES

- 453 Sex ratio distortion in the *Nesolynx thymus* (Hymenoptera: Eulophidae), an ecto-pupal parasitoid of uzifly, *Exorista sorbillans* (Diptera: Tachinidae) — B.M. PRAKASH, A. PRATHIMA, H.C. HUCHESH, H. RAVIKUMAR, S. SAMPATHKUMAR & H.P. PUTTARAJU
- 457 Identifications of *cytochrome c* and *Apaf-1* and their mRNA expressions under heat stress in insecticide-susceptible and -resistant *Plutella xylostella* (Lepidoptera: Plutellidae) — L.J. ZHANG, J.F. HUANG, F. CHEN, Z.Y. YU, Z.L. WU & G. WU
- 469 Chromosomal characteristics and evolutionary relationships of the Palearctic black fly *Simulium carthusiense* (Diptera: Simuliidae) — P.H. ADLER & G. SEITZ
- 475 Cryptic diversity and signs of mitochondrial introgression in the *Agrilus viridis* species complex (Coleoptera: Buprestidae) — M. PENTINSAARI, M. MUTANEN & L. KAILA
- 487 Effects of cold acclimation, cooling rate and heat stress on cold tolerance of the potato tuber moth *Phthorimaea operculella* (Lepidoptera: Gelechiidae) — C. HEMMATI, S. MOHARRAMPOUR & A.A. TALEBI
- 495 Comparison of the developmental time of *Bracon hebetor* (Hymenoptera: Braconidae) reared on five different lepidopteran host species and its relationship with digestive enzymes — D. SAADAT, A.R. BANDANI & M. DASTRANJ
- 501 Growth and development in a lepidopteran with variable instar number, *Pseudocoremia suavis* (Geometridae), under standard rearing conditions and when parasitised by *Meteorus pulchricornis* (Hymenoptera: Braconidae) — E.I. BARRACLOUGH, E.P.J. BURGESS, A.M. KEAN & L.A. MALONE
- 513 Female fertility in *Hippodamia convergens* (Coleoptera: Coccinellidae) is maximized by polyandry, but reduced by continued male presence — M.H. BAYOUMY & J.P. MICHAUD
- 521 Development and thermal requirements of the Nearctic predator *Geocoris punctipes* (Hemiptera: Geocoridae) reared at constant and alternating temperatures and fed on *Anagasta kuehniella* (Lepidoptera: Pyralidae) eggs — A.M. CALIXTO, V.H.P. BUENO, F.C. MONTES & J.C. VAN LENTEREN
- 529 Rarity of blister beetles (Coleoptera: Meloidae) in Southern Africa correlates with their phylogeny and trophic habits, but not body size — M. PITZALIS, V. AMORE, F. MONTALTO, L. LUISELLI & M.A. BOLOGNA
- 537 Application of the mean individual biomass of ground beetles (Coleoptera: Carabidae) to assess the assemblage successions along areas of recent glacier retreats — M. GOBBI
- 543 Generalist-specialist continuum and life history traits of Central European butterflies (Lepidoptera) — are we missing a part of the picture? — A. BARTONOVA, J. BENES & M. KONVICKA
- 555 Species composition and dynamics in abundance of migrant and sedentary butterflies (Lepidoptera) at Gibraltar during the spring period — K.J. BENSUSAN, R. NESBIT, C.E. PEREZ, P. TRYJANOWSKI & P. ZDUNIAK
- 561 Fungi associated with the red-haired bark beetle, *Hylurgus ligniperda* (Coleoptera: Curculionidae) in the forest-steppe zone in eastern Ukraine — K. DAVYDENKO, R. VASAITIS, V. MESHKOVA & A. MENKIS
- 567 Determining the instar of a weevil larva (Coleoptera: Curculionidae) using a parsimonious method — A. MERVILLE, A. VALLIER, S. VENNER, A. SIBERCHICOT, D. FOUCHET, A. HEDDI & M.-C. BEL-VENNER
- 575 Variation in the shape of the wings and taxonomy of Eurasian populations of the *Calopteryx splendens* complex (Odonata: Calopterygidae) — S. SADEGHI & H.J. DUMONT

NOTES

- 584 More evidence for an admixture of the *Hyles euphorbiae* complex's main lineages in Mediterranean Europe (Lepidoptera: Sphingidae) — M.B. MENDE & A.K. HUNDSDOERFER
- 588 Pollen analysis of the crop contents of adult Oedemeridae (Coleoptera) in Bulgaria — J. ATANASSOVA & O. SRVILOV
- 594 Discovery of the first European parasitoid of the emerald ash borer *Agrilus planipennis* (Coleoptera: Buprestidae) — M.J. ORLOVA-BIENKOWSKAJA & S.A. BELOKOBYSKIJ