

CONTENTS

ARTICLES

- 465 Meiotic karyotypes and testis structure of 14 species of the planthopper tribe Issini (Hemiptera: Fulgoroidea, Issidae) — V.G. KUZNETSOVA, A. MARYAŃSKA-NADACHOWSKA & V.M. GNEZDILOV
- 481 (Ovo-)viviparity in the darkling beetle, *Alegoria castelnaui* (Tenebrioninae: Ulomini), from Guadeloupe — A.M. DUTRILLAUX, D. PLUOT-SIGWALT & B. DUTRILLAUX
- 487 A new look at discontinuous respiration in pupae of *Hyalophora cecropia* (Lepidoptera: Saturniidae): Haemocoelic pressure, extracardiac pulsations and O₂ consumption — K. SLÁMA
- 509 Analysis of the lipids mobilized by adipokinetic hormones in the firebug *Pyrrhocoris apterus* (Heteroptera: Pyrrhocoridae) — I. BÁRTŮ, A. TOMČALA, R. SOCHA, P. ŠIMEK & D. KODRÍK
- 521 Pre-diapause mating and overwintering of fertilized adult females: New aspects of the life cycle of the wing-polymorphic bug *Pyrrhocoris apterus* (Heteroptera: Pyrrhocoridae) — R. SOCHA
- 527 Activity of *Lygus lineolaris* (Heteroptera: Miridae) adults monitored around the periphery and inside a commercial vineyard — D. FLEURY, Y. MAUFFETTE, S. METHOT & C. VINCENT
- 535 Distribution, abundance and host association of two parasitoid species attacking frugivorous drosophilid larvae in central Japan — H. MITSUI & M.T. KIMURA
- 541 Identification of limonene as a potential kairomone of the harlequin ladybird *Harmonia axyridis* (Coleoptera: Coccinellidae) — A. ALHMEDI, E. HAUBRUGE & F. FRANCIS
- 549 Effects of starvation and mating status on the activity of the flea beetle, *Phyllotreta nemorum* (Coleoptera: Chrysomelidae) — K. OKU, K.M.C.A. VERMEER, P. VERBAARSCHOT & P.W. DE JONG
- 553 Carabid beetle assemblages associated with urban golf courses in the greater Helsinki area — J. SAARIKIVI, L. IDSTRÖM, S. VENN, J. NIEMELÄ & D.J. KOTZE
- 563 Environmental correlates of species richness of Sesiidae (Lepidoptera) in Europe — M. BĄKOWSKI, W. ULRICH & Z. LAŠTŮVKA
- 571 Ecological factors determining the density-distribution of Central European dragonflies (Odonata) — F. HARABIŠ & A. DOLNÝ
- 579 Effect of population density on the development of *Mesovelia furcata* (Mesoveliidae), *Microvelia reticulata* and *Velia caprai* (Veliidae) (Heteroptera: Gerromorpha) — T. DITRICH & M. PAPAČEK
- 589 Germ cell cluster formation and ovariole structure in *Puto albicans* and *Crypticerya morrilli* (Hemiptera: Coccinea). Phylogenetic implications — T. SZKLARZEWCZ, A. MICHALIK, A. CZAJA & S. SZYDŁOWSKA
- 597 Homologies of the head of Membracoidea based on nymphal morphology with notes on other groups of Auchenorrhyncha (Hemiptera) — D.A. DMITRIEV
- 615 New morphological data from the Jurassic of Inner Mongolia confirms the damselfly aspect of Protomyrmeleontiidae (Insecta: Odonatoptera) — D.-Y. HUANG, J.F. PETRULEVIČIUS & A. NEL
- 621 The *Tachyusa coarctata* species group revisited: Phylogenetic relationships and a new species (Coleoptera: Staphylinidae: Aleocharinae) — G. PAŠNIK
- 631 Re-evaluation of the genus *Phonochorion* (Orthoptera: Tettigoniidae: Phaneropterinae) — H. SEVGILI, S.S. CAĞLAR & İ.K. SAĞLAM
- 647 Revision of the *Poecilimon ornatus* group (Orthoptera: Phaneropteridae) with particular reference to the taxa in Bulgaria and Macedonia — D.P. CHOBANOV & K.-G. HELLER

NOTES

- 673 Characterization of polymorphic microsatellite loci in the neotropical plant-ant *Allomerus decemarticulatus* (Formicidae: Myrmicinae) and multiplexing with other microsatellites from the ant subfamily Myrmicinae — P.-J.G. MALÉ, A. LOISEAU, A. ESTOUP, A. QUILICHINI & J. ORIVEL
- 677 Infestation of *Narcissus poeticus* by a scathophagid fly, *Norellia melaleuca* (Diptera: Scathophagidae) — G. CHICHIRICÒ
- 681 Temperature-dependent development of dubas bug, *Ommatissus lybicus* (Hemiptera: Tropiduchidae), an endemic pest of date palm, *Phoenix dactylifera* — A.M. MOKHTAR & S.S. AL NABHANI
- 687 An improved method for the culture of wing discs of the wingless bagworm moth, *Eumeta variegata* (Lepidoptera: Psychidae) — S. NIITSU & S. LOBBIA