

CONTENTS

| | |
|---|----|
| POLAČIK M., JANÁČ M., VASSILEV M., TRICHKOVA T. Morphometric comparison of native and non-native populations of round goby <i>Neogobius melanostomus</i> from the River Danube | 1 |
| HALAČKA K., VÍTEK T., VETEŠNÍK L., SPURNÝ P. Epidermis structure and blood parameter differences between sculpin <i>Cottus gobio</i> and Siberian sculpin <i>Cottus poecilopus</i> from the Morava watershed | 9 |
| JŮZLOVÁ Z., RIEGERT J. Vocal mimicry in the song of the icterine warbler, <i>Hippolais icterina</i> (Sylviidae, Passeriformes)..... | 17 |
| KANDEMİR İ., SÖZEN M., MATUR F., KANKILIÇ T., MARTÍNKOVÁ N., ÇOLAK F., ÖZKURT S.Ö., ÇOLAK E. Phylogeny of species and cytotypes of mole rats (Spalacidae) in Turkey inferred from mitochondrial cytochrome <i>b</i> gene sequences | 25 |
| BANASZEK A., ZIOMEK J. Genetic variation and effective population size in an isolated population of the common hamster, <i>Cricetus cricetus</i> | 34 |
| MARKOV G., LANSZKI J. Diet composition of the golden jackal, <i>Canis aureus</i> in an agricultural environment | 44 |
| DUMA I.I., GIURGIU S. Circadian activity and nest use of <i>Dryomys nitedula</i> as revealed by infrared motion sensor cameras | 49 |
| XU W., XIA C., LIN J., YANG W., BLANK D.A., QIAO J., LIU W. Diet of <i>Gazella subgutturosa</i> (Güldenstaedt, 1780) and food overlap with domestic sheep in Xinjiang, China | 54 |
| BRAHMI K., KHECHEKHOUCHE E.A., MOSTEFAOUI O., DOUMANDJI S., BAZIZ B., AULAGNIER S. First quantitative data on the diet of the fennec fox, <i>Vulpes zerda</i> (Canidae, Carnivora), in Algeria | 61 |
| BAPPERT M.-T., BURDA H., BEGALL S. To mate or not to mate? Mate preference and fidelity in monogamous Ansell's mole-rats, <i>Fukomys anelli</i> , Bathyergidae | 71 |
| FROLÍKOVÁ M., STOPKOVÁ R., ANTALÍKOVÁ J., JOHNSON P.M., STOPKA P., DVOŘÁKOVÁ-HORTOVÁ K. Role of complement regulatory proteins CD46, CD55 and CD59 in reproduction | 84 |
| Book Review..... | 95 |