

CONTENTS

MOUSAVI-SABET H., ANVARIFAR H. Landmark-based morphometric variation between <i>Cobitis keyvani</i> and <i>Cobitis faridpaki</i> (Pisces: Cobitidae), with new habitat for <i>C. faridpaki</i> in the southern Caspian Sea basin	167
SUVOROV P., ŠÁLEK M. Character of surrounding habitat determines nest predation in suburban idle fields	176
ZHANG M., WANG X., DING Y. Flight responses of blue sheep in Ningxia Helan Mountain National Nature Reserve	185
KLENOVŠEK T., NOVAK T., ČAS M., TRILAR T., JANŽEKOVIČ F. Feeding ecology of three sympatric <i>Sorex</i> shrew species in montane forests of Slovenia	193
MARKOV G., NINOV N., ANDREEV R. Craniological variation of the Balkan chamois, <i>Rupicapra rupicapra balcanica</i> from Bulgaria	200
RUSIN M.Y., BANASZEK A., MISHTA A.V. The common hamster (<i>Cricetus cricetus</i>) in Ukraine: evidence for population decline	207
ALMEIDA D., RODOLFO N., SAYER C.D., COPP G.H. Seasonal use of ponds as foraging habitat by Eurasian otter with description of an alternative handling technique for common toad predation.....	214
ŠTOFIK J., MERGANIČ J., MERGANIČOVÁ K., SANIGA M. Seasonal changes in food composition of the brown bear (<i>Ursus arctos</i>) from the edge of its occurrence – Eastern Carpathians (Slovakia).....	222
MORI E., SFORZI A. Structure of phytobezoars found in the stomach of a crested porcupine, <i>Hystrix cristata</i> L., 1758	232
SUTOR A., SCHWARZ S. Seasonal habitat selection of raccoon dogs (<i>Nyctereutes procyonoides</i>) in Southern Brandenburg, Germany	235
Book Review	244