

# CONTENTS

LAMBRECHTS M.M., SCHATZ B. Ants and Paridae share nesting boxes in continental Mediterranean habitat.....	63
ŠÁLEK M., ZÁMEČNÍK V. Delayed nest predation: a possible tactic toward nests of open-nesting birds ....	67
MOHAPATRA R.K., PANDA S. Husbandry, behaviour and conservation breeding of Indian pangolin .....	73
ZHANG Y., LUO Z., LIU B. The effect of group size on the vigilance of Mongolian gazelle ( <i>Procapra gutturosa</i> ) .....	81
KONTSIOTIS V.J., BAKALLOUDIS D.E., TSIOMPANOUDIS A.C., XOFIS P. Body condition variation of wild rabbit population in the north-east Mediterranean island of Lemnos – Greece .....	87
MERTA D., MOCAŁA P., POMYKACZ M., FRĄCKOWIAK W. Autumn-winter diet and fat reserves of wild boars ( <i>Sus scrofa</i> ) inhabiting forest and forest-farmland environment in south-western Poland.....	95
DVOŘÁK S., BARTÁK V., MACHÁČEK Z., MATĚJŮ J. Home range size and spatio-temporal dynamics of male sika deer ( <i>Cervus nippon</i> ; Cervidae, Artiodactyla) in an introduced population .....	103
DI FEBBRARO M., LOY A. A new method based on indirect evidences to infer activity pattern in moles. A test on the blind mole in Central Apennines.....	116
SÁNDOR G., LÁSZLÓ R., NÁHLIK, A. Determination of time of conception of fallow deer in a Hungarian free range habitat.....	122
BLANK D.A., RUCKSTUHL K., YANG W. Secretion marking with preorbital glands in goitered gazelle, <i>Gazella subgutturosa</i> (Artiodactyla: Bovidae).....	127