

Contents in full

| | |
|---------------------------------------------------------------------------------------------------------------------|-----|
| The Author | v |
| A Note from the Publisher on Primers in Biology | vii |
| Preface | ix |
| Acknowledgements | xii |
| CHAPTER 1 The Cell Cycle | |
| 1-0 Overview: Cell Reproduction | 2 |
| Cell reproduction is a fundamental feature of all living things | |
| Cells reproduce in discrete steps | |
| The ordering of cell-cycle events is governed by an independent control system | |
| 1-1 Events of the Eukaryotic Cell Cycle | 4 |
| Chromosome duplication and segregation occur in distinct cell-cycle phases that are usually separated by gap phases | |
| Cytoplasmic components are duplicated throughout the cell cycle | |
| Cell growth is usually coordinated with cell division | |
| 1-2 Variations in Cell-Cycle Organization | 6 |
| Cell-cycle structure varies in different cells and organisms | |
| Multiple rounds of chromosome duplication or segregation can occur in the same cell cycle | |
| The symmetry of cell division varies in different cell types | |
| 1-3 The Cell-Cycle Control System | 8 |
| Cell-cycle events are governed by an independent control system | |
| The cell-cycle control system is based on oscillations in the activities of cyclin-dependent protein kinases | |
| Cell-cycle events are initiated at three regulatory checkpoints | |
| Cell-cycle progression in most cells can be blocked at checkpoints | |
| CHAPTER 2 Model Organisms in Cell-Cycle Analysis | |
| 2-0 Overview: Cell-Cycle Analysis in Diverse Eukaryotes | 12 |
| Mechanisms of cell-cycle control are similar in all eukaryotes | |
| Budding and fission yeasts provide powerful systems for the genetic analysis of eukaryotic cell-cycle control | |
| Early animal embryos are useful for the biochemical characterization of simple cell cycles | |
| Control of cell division in multicellular organisms can be dissected genetically in <i>Drosophila</i> | |
| Cultured cell lines provide a means of analyzing cell-cycle control in mammals | |
| 2-1 Life Cycles of Budding and Fission Yeasts | 14 |
| Budding yeast and fission yeast divide by different mechanisms | |
| Yeast cells alternate between haploid and diploid states and undergo sporulation in response to starvation | |
| 2-2 Genetic Analysis of Cell-Cycle Control in Yeast | 16 |
| Cell biological processes are readily dissected with yeast genetic methods | |
| Conditional mutants are used to analyze essential cell-cycle processes | |
| Homologous genes have different names in fission yeast and budding yeast | |
| 2-3 The Early Embryo of <i>Xenopus laevis</i> | 18 |
| The early embryonic divisions of <i>Xenopus</i> provide a simplified system for cell-cycle analysis | |
| Unfertilized eggs develop from diploid oocytes by meiosis | |
| The early embryonic cell cycle can be reconstituted in a test tube | |
| 2-4 The Fruit Fly <i>Drosophila melanogaster</i> | 20 |
| <i>Drosophila</i> allows genetic analysis of cell-cycle control in metazoans | |
| Cells of the early <i>Drosophila</i> embryo divide by a simplified cell cycle | |
| Gap phases are introduced in late embryogenesis | |

| | | |
|------------|---------------------------------------------------------------------------------------------------------------------------|----|
| | Adult fly structures develop from imaginal cells | |
| 2-5 | Mammalian Cell-Cycle Analysis | 22 |
| | Mammalian cell-cycle control can be analyzed in cells growing in culture | |
| | Mutations lead to immortalization and transformation of mammalian cells | |
| | Specific gene disruption is the ideal approach for assessing protein function in mammalian cells | |
| 2-6 | Methods in Cell-Cycle Analysis | 24 |
| | Cell-cycle position can be assessed by many approaches | |
| | Cell populations can be synchronized at specific cell-cycle stages | |
| | Complete understanding of cell-cycle control mechanisms requires the analysis of protein structure and enzymatic behavior | |

CHAPTER 3 The Cell-Cycle Control System

| | | |
|-------------|-----------------------------------------------------------------------------------------------------------------|----|
| 3-0 | Overview: The Cell-Cycle Control System | 28 |
| | The cell-cycle control system is a complex assembly of oscillating protein kinase activities | |
| | Multiple regulatory mechanisms govern Cdk activity during the cell cycle | |
| | The cell-cycle control system generates robust, switch-like and adaptable changes in Cdk activity | |
| 3-1 | Cyclin-Dependent Kinases | 30 |
| | The cyclin-dependent kinases are a small family of enzymes that require cyclin subunits for activity | |
| | The active site of cyclin-dependent kinases is blocked in the absence of cyclin | |
| 3-2 | Cyclins | 32 |
| | Cyclins are the key determinants of Cdk activity and can be classified in four groups | |
| | Cyclins contain a conserved helical core | |
| 3-3 | Control of Cdk Activity by Phosphorylation | 34 |
| | Full Cdk activity requires phosphorylation by the Cdk-activating kinase | |
| | Cdk function is regulated by inhibitory phosphorylation by Wee1 and dephosphorylation by Cdc25 | |
| 3-4 | The Structural Basis of Cdk Activation | 36 |
| | The conformation of the Cdk active site is dramatically rearranged by cyclin binding and phosphorylation by CAK | |
| 3-5 | Substrate Targeting by Cyclin-Cdk Complexes | 38 |
| | Cyclins are specialized for particular functions | |
| | Cyclins can interact directly with the substrates of the associated Cdk | |
| | Cyclins can direct the associated Cdk to specific subcellular locations | |
| | Cks1 may serve as an adaptor protein that targets Cdks to phosphoproteins | |
| 3-6 | Cdk Regulation by Inhibitory Subunits | 40 |
| | Cdk inhibitors help suppress Cdk activity in G1 | |
| | Cip/Kip proteins bind both subunits of the cyclin-Cdk complex | |
| | G1-Cdks are activated by Cip/Kip proteins and inhibited by INK4 proteins | |
| 3-7 | Biochemical Switches in Signaling Systems | 42 |
| | Components of the cell-cycle control system are assembled into biochemical switches | |
| | Switch-like behavior can be generated by various mechanisms | |
| | Bistability is required for an effective binary switch | |
| 3-8 | Switch-Like Activation of Cdk1 | 44 |
| | Cdk1 activation at mitosis is based on positive feedback | |
| | Cdk switches are robust as a result of multiple partly redundant mechanisms | |
| 3-9 | Protein Degradation in Cell-Cycle Control | 46 |
| | Many cell-cycle regulators are destroyed by ubiquitin-dependent proteolysis | |
| | SCF catalyzes ubiquitination of phosphorylated substrates using interchangeable substrate-targeting subunits | |
| 3-10 | The Anaphase-Promoting Complex | 48 |
| | The APC initiates anaphase and mitotic exit | |
| | Cdc20 activates the APC in anaphase | |
| | APC activity is maintained in G1 by Cdh1 | |
| | APC targets contain specific recognition sequences | |

| | |
|------------------------------------------------------------------------------------------------------|----|
| 3-11 Assembling and Regulating a Cell-Cycle Oscillator | 50 |
| Negative feedback can generate a repeating oscillator | |
| Regulated braking mechanisms allow the Cdk oscillator to be paused in G1 | |
| 3-12 Transcriptional Control of Cell-Cycle Regulators | 52 |
| A sequential program of gene expression contributes to cell-cycle control | |
| Expression of a large fraction of the genes in the yeast genome is regulated during the cell cycle | |
| Key gene regulatory proteins in yeast are activated at the major cell-cycle transitions | |
| The E2F family controls cell-cycle-dependent changes in gene expression in metazoans | |
| 3-13 Programming the Cell-Cycle Control System | 54 |
| The order of cell-cycle events is determined by regulatory interactions between multiple oscillators | |
| The cell-cycle control system is responsive to many external inputs | |

CHAPTER 4 Chromosome Duplication

| | |
|------------------------------------------------------------------------------------------------------|----|
| 4-0 Overview: Chromosome Duplication and its Control | 58 |
| DNA synthesis begins at replication origins | |
| The cell-cycle control system activates replication origins only once in each S phase | |
| Chromosome duplication requires duplication of chromatin structure | |
| 4-1 Basic Mechanisms of DNA Synthesis | 60 |
| The two strands of DNA are replicated by different mechanisms | |
| DNA replication begins with origin unwinding and primer synthesis | |
| Discontinuous DNA fragments are joined together by DNA ligase | |
| Telomerase synthesizes DNA at chromosome ends | |
| 4-2 The Replication Origin | 62 |
| Replication origins in budding yeast contain well defined DNA sequences | |
| Replication origins in animal chromosomes are defined by several factors in addition to DNA sequence | |
| 4-3 Assembly of the Prereplicative Complex at the Replication Origin | 64 |
| The replication origin interacts with a multisubunit protein complex | |
| The ORC and accessory proteins load the Mcm helicase onto origins | |
| Mcm loading involves ATP-dependent protein remodeling | |
| 4-4 Regulation of the Prereplicative Complex | 66 |
| Assembly of prereplicative complexes is restricted to G1 by multiple mechanisms | |
| Prereplicative complex components are destroyed or inhibited in yeast as a result of Cdk activity | |
| Pre-RC assembly is controlled in animals by both Cdks and the APC | |
| 4-5 Cyclins Required for Activation of Replication Origins in Yeast | 68 |
| Cdks and Cdc7 trigger the initiation of DNA replication | |
| In budding yeast, the cyclins Clb5 and Clb6 are key activators of replication origins | |
| Yeast cells lacking S cyclins can replicate their DNA | |
| 4-6 Cyclins Required for Activation of Replication Origins in Metazoans | 70 |
| Different cyclins control initiation of DNA replication in different stages of animal development | |
| Cyclin A is a major regulator of replication initiation in cultured mammalian cells | |
| DNA replication in frog embryos is triggered by cyclin E-Cdk2 | |
| Cyclin E-Cdk2 is a major regulator of DNA replication in <i>Drosophila</i> | |
| 4-7 Control of Replication by the Protein Kinase Cdc7-Dbf4 | 72 |
| Cdc7 triggers the activation of replication origins | |
| Cdc7 is activated during S phase by the regulatory subunit Dbf4 | |
| Dbf4 levels are regulated by multiple mechanisms | |
| 4-8 Activation of the Replication Origin | 74 |
| Replication begins with DNA unwinding at the origin | |
| Late-firing origins are regulated independently | |

| | | |
|-------------|----------------------------------------------------------------------------------------------------------------------|----|
| | Replication must be completed before chromosome segregation occurs | |
| 4-9 | Basic Chromatin Structure | 76 |
| | Chromatin is complex and dynamic | |
| | The basic unit of chromatin structure is the nucleosome | |
| | Higher-order chromatin structure is also controlled by non-histone proteins, histone H1 and histone modifications | |
| 4-10 | Histone Synthesis in S phase | 78 |
| | Histone synthesis rises sharply during S phase | |
| | Transcription of histone genes increases in S phase | |
| | Histone mRNA processing and stability increase in S phase | |
| | The level of free histones in the cell acts as a signal to link histone synthesis to DNA synthesis | |
| 4-11 | Nucleosome Assembly on Nascent DNA | 80 |
| | Nucleosomes are distributed to both new DNA strands behind the replication fork | |
| | Nucleosome assembly factors load histones on nascent DNA | |
| 4-12 | Heterochromatin at Telomeres and Centromeres | 82 |
| | Heterochromatin is inherited by epigenetic mechanisms | |
| | Telomeres are packaged in a heritable heterochromatin structure | |
| | The centromere nucleates a heritable and poorly understood form of heterochromatin | |
| 4-13 | Molecular Mechanisms of Heterochromatin Duplication | 84 |
| | Duplication of heterochromatin structure involves proteins that recognize and promote localized histone modification | |
| | The Sir proteins form a heritable polymer at telomeres in budding yeast | |
| | HP1 may nucleate heritable chromatin structure at the centromere and other regions | |
| | Sister-chromatid cohesion in S phase prepares the cell for mitosis | |

CHAPTER 5 Early Mitosis: Preparing the Chromosomes for Segregation

| | | |
|------------|--------------------------------------------------------------------------------------------------------|-----|
| 5-0 | Overview: The Events of Mitosis | 88 |
| | The central events of mitosis are sister-chromatid separation and segregation | |
| | The events of early mitosis set the stage for sister-chromatid segregation | |
| | The completion of mitosis begins with sister-chromatid segregation | |
| 5-1 | Overview: Principles of Mitotic Regulation | 90 |
| | Phosphorylation and proteolysis control progression through mitosis | |
| | Mitotic events must go to completion | |
| | Mitotic entry and exit are major regulatory transitions with differing importance in different species | |
| 5-2 | Cyclins that Promote Mitotic Entry in Yeast | 92 |
| | Cyclin–Cdk complexes trigger mitotic entry in all eukaryotes | |
| | Fission yeast cells trigger mitosis with a single mitotic cyclin | |
| | Two pairs of mitotic cyclins control budding yeast mitosis | |
| 5-3 | Cyclins that Promote Mitotic Entry in Metazoans | 94 |
| | Mitosis in metazoans is governed by cyclins A and B | |
| | Vertebrate mitosis is driven by multiple forms of cyclins A and B | |
| | The active cyclin B1–Cdk1 complex moves from cytoplasm to nucleus in late prophase | |
| | Vertebrate cyclins A and B drive different mitotic events | |
| 5-4 | Regulation of Mitotic Cdks by Wee1 and Cdc25 | 96 |
| | Cyclin B–Cdk1 complexes are activated rapidly in early M phase by dephosphorylation | |
| | Multiple Wee1-related kinases and Cdc25-related phosphatases govern Cdk1 activity in animal cells | |
| 5-5 | Switch-like Activation of Cyclin B–Cdk1 at Mitosis | 98 |
| | Mitotic Cdk1 activation involves multiple positive feedback loops | |
| | Cdc25B and cyclin A–Cdk help trigger cyclin B–Cdk1 activation | |
| 5-6 | Subcellular Localization of Mitotic Regulators | 100 |
| | Cyclin B1–Cdk1 is regulated by changes in its subcellular localization | |

| | | |
|--------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|-----|
| | Cyclin B1–Cdk1 location is controlled by phosphorylation of cyclin B1 | |
| | Cdc25C localization is regulated by phosphorylation | |
| | Cyclin B1–Cdk1 activation and nuclear accumulation are partly interdependent | |
| 5-7 | Protein Kinases of the Polo and Aurora Families | 102 |
| | Polo-like kinases (Plks) help control spindle assembly and mitotic exit | |
| | Spindle function and sister-chromatid segregation are controlled in part by aurora kinases | |
| 5-8 | Preparations for Mitosis: Sister-Chromatid Cohesion | 104 |
| | Sister chromatids are held together by two mechanisms | |
| | Cohesin is a key mediator of sister-chromatid cohesion | |
| | Cohesion is established during DNA replication | |
| | DNA decatenation prepares sister chromatids for separation | |
| 5-9 | Entry into Mitosis: Sister-Chromatid Condensation and Resolution | 106 |
| | Chromosomes are dramatically reorganized in mitosis | |
| | Condensin complexes drive chromosome condensation and resolution | |
| 5-10 | Regulation of Chromosome Condensation and Resolution | 108 |
| | Mitotic Cdks act on condensin to govern the timing of chromosome condensation | |
| | Sister-chromatid resolution is governed by Plk and aurora B in animal cells | |
| | | |
| CHAPTER 6 Assembly of the Mitotic Spindle | | |
| 6-0 | Overview: The Mitotic Spindle | 112 |
| | Chromosome segregation depends on the mitotic spindle | |
| | The mitotic spindle must be bipolar | |
| | Multiple mechanisms drive spindle assembly | |
| 6-1 | Microtubule Structure and Behavior | 114 |
| | Microtubules are polymers of tubulin subunits | |
| | Microtubules exhibit dynamic instability | |
| 6-2 | Microtubule Nucleation, Stability and Motility | 116 |
| | Cellular microtubules originate on preformed protein complexes that are usually concentrated in a microtubule-organizing center | |
| | Microtubule dynamics are governed by a variety of stabilizing and destabilizing proteins | |
| | Motor proteins move along microtubules | |
| 6-3 | The Centrosome and the Spindle Pole Body | 118 |
| | The centrosome cycle resembles the chromosome cycle | |
| | Centrosome behavior is determined by the centrioles | |
| | The yeast spindle pole body is embedded in the nuclear envelope | |
| 6-4 | Control of Centrosome Duplication | 120 |
| | Duplication of the centrosome and spindle pole body is initiated in late G1 by G1/S–Cdks | |
| | Centrosome duplication normally occurs once per cell cycle | |
| 6-5 | The Kinetochore | 122 |
| | The kinetochore is the major site of microtubule–chromosome attachment | |
| | The kinetochore provides a stable attachment to a dynamic microtubule plus end | |
| 6-6 | Early Steps in Spindle Assembly | 124 |
| | Spindle assembly begins in prophase | |
| | Mitotic microtubules are highly dynamic | |
| | Centrosome separation initiates spindle assembly | |
| | Centrosome maturation increases microtubule nucleation in mitosis | |
| 6-7 | Nuclear Envelope Breakdown | 126 |
| | The nuclear envelope is composed of two membranes on an underlying protein support | |
| | Nuclear envelope breakdown begins at nuclear pores | |
| | The endoplasmic reticulum and Golgi apparatus are reorganized in mitosis | |
| 6-8 | Mitotic Chromosome Function in Spindle Assembly | 128 |
| | Spindles self-organize around chromosomes | |
| | Microtubules can be stabilized by a gradient of Ran–GTP around chromosomes | |

| | |
|-----------------------------------------------------------------------------------------------|-----|
| 6-9 Attachment of Sister Chromatids to the Spindle | 130 |
| Centrosomes search for and capture kinetochores in prometaphase | |
| Some kinetochore microtubules originate at the kinetochore | |
| Chromosome attachment results in tension between sister kinetochores | |
| 6-10 Bi-Orientation of Sister Chromatids | 132 |
| Kinetochore–microtubule attachment is stabilized by tension | |
| Aurora B is required for the correction of syntelic attachments | |
| Merotelic attachments are processed by multiple mechanisms | |
| 6-11 Forces Driving Chromosome Movement | 134 |
| Multiple forces act on chromosomes in the spindle | |
| The kinetochore is a major source of poleward force | |
| Microtubule flux generates poleward force | |
| A polar ejection force is generated by chromosome arms | |
| 6-12 Chromosome Congression | 136 |
| Chromosome oscillations in prometaphase are generated by changes in the state of kinetochores | |
| Microtubule flux may promote chromosome congression | |

CHAPTER 7 The Completion of Mitosis

| | |
|------------------------------------------------------------------------------------------------------------|-----|
| 7-0 Overview: The Completion of Mitosis | 140 |
| The final events of mitosis occur in anaphase and telophase | |
| The metaphase-to-anaphase transition is initiated by ubiquitination and destruction of regulatory proteins | |
| Dephosphorylation of Cdk targets drives the events of late M phase | |
| APC ^{Cdc20} initiates Cdk inactivation | |
| 7-1 Initiation of Anaphase: Activation of the APC | 142 |
| APC ^{Cdc20} activation in early mitosis is essential for anaphase to occur | |
| Phosphorylation promotes APC ^{Cdc20} activation in early mitosis | |
| 7-2 Initiation of Anaphase: The Spindle Checkpoint | 144 |
| Unattached kinetochores generate a signal that prevents anaphase | |
| The spindle checkpoint monitors defects in microtubule attachment and kinetochore tension | |
| 7-3 Inhibition of APC^{Cdc20} by the Spindle Checkpoint | 146 |
| Unattached kinetochores catalyze the formation of inhibitory signaling complexes | |
| The spindle checkpoint signal is rapidly turned off once kinetochores are attached | |
| 7-4 Control of Sister-Chromatid Separation | 148 |
| Separase is inhibited before anaphase by securin | |
| In vertebrate cells Cdk1 inhibits separase by phosphorylation | |
| 7-5 Control of Late Mitosis in Budding Yeast | 150 |
| Cdk inactivation in mitosis in budding yeast is not due to APC ^{Cdc20} alone | |
| The protein phosphatase Cdc14 is required to complete mitosis in budding yeast | |
| 7-6 Control of Anaphase Events | 152 |
| The anaphase spindle segregates the chromosomes | |
| Dephosphorylation of Cdk targets governs anaphase spindle behavior | |
| 7-7 Control of Telophase | 154 |
| Dephosphorylation of Cdk substrates drives the final steps of mitosis | |
| Spindle disassembly is the central event of telophase | |
| Nuclear envelope assembly begins around individual chromosomes | |

CHAPTER 8 Cytokinesis

| | |
|-------------------------------------------------------------------------|-----|
| 8-0 Overview: Cytokinesis | 158 |
| Cytokinesis distributes daughter nuclei into separate cells | |
| Cytokinesis depends on a contractile ring and membrane deposition | |
| The cleavage plane is positioned between the daughter nuclei | |
| The timing of cytokinesis is coordinated with the completion of mitosis | |

| | | |
|------------------------------|-----------------------------------------------------------------------------------------------------|-----|
| 8-1 | The Actin–Myosin Ring | 160 |
| | Bundles of actin assemble at the site of division | |
| | Force is generated in the contractile ring by non-muscle myosin II | |
| | Actin filament formation depends on formins | |
| 8-2 | Assembly and Contraction of the Actin–Myosin Ring | 162 |
| | Contractile ring function depends on accessory factors whose importance varies in different species | |
| | Contraction of the actin–myosin ring is regulated by activation of myosin II | |
| | The GTPase Rho controls actin and myosin behavior at the cleavage site | |
| 8-3 | Membrane and Cell Wall Deposition at the Division Site | 164 |
| | Membrane deposition is required during cytokinesis | |
| | Membrane addition occurs in parallel with actin–myosin contraction | |
| 8-4 | The Positioning and Timing of Cytokinesis in Yeast | 166 |
| | Preparations for cytokinesis in budding yeast begin in late G1 | |
| | Fission yeast uses the nucleus to mark the division site in early mitosis | |
| 8-5 | The Positioning and Timing of Cytokinesis in Animal Cells | 168 |
| | Signals from the mitotic spindle determine the site of cleavage in animal cells | |
| | Multiple regulatory components at the central spindle help control cytokinesis | |
| | Cytokinesis is coordinated with mitosis by the spindle and Cdk1 inactivation | |
| 8-6 | Specialization of Cytokinesis in Animal Development | 170 |
| | Cytokinesis can be blocked or incomplete in some stages of development | |
| | Cellularization is a specialized form of cytokinesis | |
| 8-7 | Asymmetric Cell Division | 172 |
| | Asymmetric spindle positioning leads to daughter cells of unequal sizes | |
| | Unequal forces on the poles underlie asymmetric spindle positioning | |
| | The orientation of cell division is controlled by the mitotic spindle | |
| CHAPTER 9 Meiosis | | |
| 9-0 | Overview: Meiosis | 176 |
| | Sexual reproduction is based on the fusion of haploid cells | |
| | The meiotic program involves two rounds of chromosome segregation | |
| | Homologous recombination is an important feature of meiosis | |
| | Defects in meiosis lead to aneuploidy | |
| 9-1 | Regulation of Early Meiotic Events in Yeast | 178 |
| | The meiotic program is controlled at multiple checkpoints | |
| | The transcription factor Ime1 initiates the budding yeast meiotic program | |
| | Entry into the meiotic program is driven by the protein kinase Ime2 | |
| 9-2 | Homologous Recombination in Meiosis | 180 |
| | Homologous recombination is a central feature of meiotic prophase | |
| 9-3 | Homolog Pairing in Meiotic Prophase | 182 |
| | Stages of meiotic prophase are defined by cytological landmarks | |
| | Homolog pairing occurs in two successive stages | |
| 9-4 | Chiasma Formation in Late Meiotic Prophase | 184 |
| | A small number of recombination sites are selected for crossover formation in zygotene | |
| | Crossover sites nucleate the synaptonemal complex in some species | |
| | Chiasmata appear in diplotene | |
| 9-5 | Controlling Entry into the First Meiotic Division | 186 |
| | Meiosis I is initiated by M–Cdk activity | |
| | Entry into the first meiotic division of animal cells is controlled in diplotene | |
| | Ndt80 and Cdk1 promote entry into the meiotic divisions of budding yeast | |
| | Recombination defects block entry into meiosis I | |
| 9-6 | Chromosome Attachment in Meiosis I | 188 |
| | Homolog pairs are bi-oriented on the first meiotic spindle | |
| | Homolog bi-orientation depends on cohesion of sister-chromatid arms | |
| | Homolog linkage does not involve chiasmata in some species | |

| | |
|---------------------------------------------------------------------------------------------------------------------------|-----|
| 9-7 Chromosome Segregation in Meiosis I | 190 |
| Loss of sister-chromatid arm cohesion initiates anaphase I | |
| The spindle checkpoint system helps control anaphase I | |
| Centromeric cohesin is protected from cleavage in meiosis I | |
| 9-8 Finishing Meiosis | 192 |
| Meiosis I is followed by meiosis II | |
| Partial Cdk1 inactivation occurs after meiosis I | |
| The meiotic program is coordinated with gametogenesis | |
| | |
| CHAPTER 10 Control of Cell Proliferation and Growth | |
| 10-0 Overview: Control of Cell Proliferation and Growth | 196 |
| Cell proliferation is controlled at a checkpoint in late G1 | |
| Progression through Start depends on an irreversible wave of Cdk activity | |
| Progression through Start requires changes in gene expression | |
| Cell division is often coordinated with cell growth | |
| 10-1 Activation of Gene Expression at Start in Budding Yeast | 198 |
| The gene regulatory proteins SBF and MBF drive expression of Start-specific genes in yeast | |
| SBF and MBF are activated by Cln3–Cdk1 at Start | |
| Small changes in the amount of Cln3 help trigger cell-cycle entry | |
| SBF and MBF are inactivated in S phase by Clb–Cdk1 complexes | |
| 10-2 Activation of S–Cdks in Budding Yeast | 200 |
| G1/S–Cdks promote activation of S–Cdks | |
| Multisite phosphorylation of Sic1 generates switch-like S–Cdk activation | |
| G1/S– and S–Cdks collaborate to inactivate APC ^{Cdh1} after Start | |
| 10-3 Extracellular Control of Start in Yeast: Mating Factor Signaling | 202 |
| Yeast mating factors induce cell-cycle arrest in G1 | |
| Far1 has multiple functions in proliferating and arrested cells | |
| Far1 phosphorylation is triggered by a G-protein signaling pathway | |
| 10-4 Activation of Gene Expression at Start in Animals | 204 |
| E2F transcription factors help control G1/S gene expression in animals | |
| Stimulation of G1/S gene expression results from a combination of increased gene activation and decreased gene repression | |
| E2F function is regulated by pRB proteins | |
| 10-5 Regulation of E2F–pRB Complexes | 206 |
| G1/S gene expression at Start involves the replacement of repressor E2Fs with activator E2Fs | |
| Phosphorylation of pRB proteins releases E2F | |
| Multiple mechanisms of E2F activation provide robust regulation of Start | |
| 10-6 Mitogenic Signaling in Animal Cells | 208 |
| Extracellular mitogens control the rate of cell division in animals | |
| Activated mitogen receptors recruit signaling complexes to the cell membrane | |
| Ras and Myc are components of many mitogenic signaling pathways | |
| Activation of the PI3 kinase helps promote mitogenesis | |
| 10-7 Activation of G1–Cdks by Mitogens | 210 |
| Mitogenic signaling pathways lead to activation of cyclin D–Cdk complexes | |
| Mitogens control cyclin D–Cdk localization and destruction | |
| Mitogens and anti-mitogens control the concentrations of Cdk inhibitor proteins | |
| 10-8 Activation of G1/S– and S–Cdk Complexes in Animal Cells | 212 |
| G1/S–Cdk activation at Start depends on removal of the inhibitor p27 | |
| Cyclin A–Cdk2 activation is promoted in part by APC inhibition | |
| 10-9 Developmental Control of Cell Proliferation | 214 |
| Developmental signals limit cell division to specific embryonic regions | |
| Embryonic divisions are limited by depletion of key cell-cycle regulators | |
| 10-10 Overview: Coordination of Cell Division and Cell Growth | 216 |
| Cell division and cell growth are separate processes | |

| | |
|-------------------------------------------------------------------------------------------------------|-----|
| Cell growth is regulated by extracellular nutrients and growth factors | |
| Cell growth and division are coordinated by multiple mechanisms | |
| The size of a cell depends on its genomic content | |
| 10-11 Control of Cell Growth | 218 |
| Cell growth rate is determined primarily by the rate of protein synthesis | |
| Extracellular nutrients and growth factors stimulate cell growth by activating the protein kinase TOR | |
| TOR affects cell growth mainly by stimulating protein synthesis | |
| Growth factors stimulate protein synthesis through the activation of PI3 kinase | |
| 10-12 Coordination of Cell Growth and Division in Yeast | 220 |
| Yeast cell growth and division are tightly coupled | |
| Yeast cells monitor translation rates as an indirect indicator of cell size | |
| Growth thresholds are rapidly adjustable | |
| 10-13 Coordination of Growth and Division in Animal Cells | 222 |
| Growth and division are coordinated by multiple mechanisms in animal cells | |
| Division depends on growth in many animal cell types | |
| Animal cell growth and division are sometimes controlled independently | |
| 10-14 Control of Cell Death | 224 |
| Animal cell numbers are determined by a balance of cell birth and death | |
| Survival factors suppress the mitochondrial pathway of apoptosis | |
| DNA damage and other stresses can trigger apoptosis | |
| | |
| CHAPTER 11 The DNA Damage Response | |
| 11-0 Overview: The DNA Damage Response | 228 |
| The DNA damage response helps maintain the genome | |
| ATR and ATM are conserved protein kinases at the heart of the DNA damage response | |
| Replication defects trigger a DNA damage response | |
| 11-1 Detection and Repair of DNA Damage | 230 |
| DNA can be damaged in many ways | |
| Base and nucleotide excision repair systems repair nucleotide damage | |
| Double-strand breaks are repaired by two main mechanisms | |
| 11-2 The DNA Damage Response: Recruitment of ATR and ATM | 232 |
| ATR is required for the response to multiple forms of damage | |
| ATM is specialized for the response to unprocessed double-strand breaks | |
| 11-3 The DNA Damage Response: Adaptors and Chk1 and Chk2 | 234 |
| Protein complexes assemble at DNA damage sites to coordinate DNA repair and the damage response | |
| A PCNA-like complex is required for the ATR-mediated damage response | |
| Adaptor proteins link DNA damage to activation of Chk1 and Chk2 | |
| 11-4 Activation of p53 by DNA Damage | 236 |
| p53 is responsible for long-term inhibition of cell proliferation in animal cells | |
| The major regulators of p53 include Mdm2, p300 and ARF | |
| Damage response kinases phosphorylate p53 and Mdm2 | |
| 11-5 Effects of DNA Damage on Progression through Start | 238 |
| DNA damage blocks cell-cycle progression at multiple points | |
| DNA damage has minor effects on progression through Start in budding yeast | |
| DNA damage in vertebrate cells triggers a robust G1 arrest | |
| p53 has different effects in different cell types | |
| 11-6 Effects of DNA Damage at Replication Forks | 240 |
| A DNA damage response is initiated at replication forks during S phase | |
| ATR is the key initiator of the response to stalled replication forks | |
| The DNA damage response stabilizes the replication fork | |
| 11-7 Effects of DNA Damage on DNA Synthesis and Mitosis | 242 |
| DNA damage in S phase blocks replication origin firing | |

| | |
|-------------------------------------------------------------------------------------------------------------------|-----|
| DNA damage blocks mitotic entry in most eukaryotes | |
| DNA damage blocks anaphase in budding yeast | |
| 11-8 Responses to Mitogenic and Telomere Stress | 244 |
| Hyperproliferative signals trigger the activation of p53 | |
| Imbalances in mitogenic stimuli promote replicative senescence in mouse cells | |
| Telomere degeneration promotes cell-cycle arrest in human cells | |
| CHAPTER 12 The Cell Cycle in Cancer | |
| 12-0 Overview: Cell-Cycle Defects in Cancer | 248 |
| Cancer cells break the communal rules of tissues | |
| Cancer progression is an evolutionary process driven by gene mutation | |
| Genetic instability accelerates cancer progression | |
| 12-1 Gene Mutations that Drive Cancer | 250 |
| Mutations in oncogenes and tumor suppressors stimulate tumor progression | |
| Oncogenes can be activated by many different mechanisms | |
| Multiple mutations are required to cripple tumor suppressor genes | |
| Cancer can be initiated by mechanisms other than gene mutation | |
| 12-2 Tissue Specificity in Cancer | 252 |
| Cancers are a complex group of diseases | |
| The molecular basis of tumorigenesis can vary in different tissues | |
| 12-3 Stimulation of Cell-Cycle Entry in Cancer Cells | 254 |
| Tumor cells are independent of mitogens and resistant to anti-mitogens | |
| G1/S gene regulation is defective in most cancers | |
| Multiple mitogenic defects are required for tumor formation | |
| 12-4 Cell Growth and Survival in Tumors | 256 |
| Cell growth is stimulated in tumors | |
| Tumor cells are less dependent than normal cells on survival factors | |
| Differentiation is often inhibited in tumor cells | |
| Tumor cells are resistant to the hyperproliferation stress response | |
| 12-5 Genetic Instability in Cancer | 258 |
| Most cancer cells have unstable genomes | |
| Defects in the DNA damage response promote genetic instability in cancer | |
| Genetic instability sometimes results from an increased rate of point mutation | |
| Chromosomal instability is the major form of genetic instability | |
| 12-6 Telomeres and the Structural Instability of Chromosomes | 260 |
| Defective DNA damage responses can lead to chromosomal instability | |
| Degenerating telomeres can lead to chromosomal instability | |
| 12-7 Instability in Chromosome Number | 262 |
| Cancer cells often become aneuploid through a tetraploid intermediate | |
| Cancer cells often contain excessive numbers of centrosomes | |
| Mutations in mitotic spindle components contribute to chromosomal instability | |
| 12-8 Cancer Progression | 264 |
| There are many genetic routes to a malignant cancer | |
| Colon cancer progression usually begins with mutations in the gene <i>APC</i> | |
| Two forms of genetic instability drive colorectal cancer progression | |
| 12-9 Stopping Cancer | 266 |
| Reducing cancer mortality begins with prevention and early diagnosis | |
| Therapies must kill cancer cells but leave healthy cells intact | |
| A detailed understanding of the molecular basis of cancer may lead to rational and more specific cancer therapies | |
| Glossary | 269 |
| References | 275 |
| Index | 287 |