

Contents

1.1.1	Magnetism of Free Lanthanide Ions	2
1.1.2	Spin–Orbital Coupling and Anisotropy of Free Lanthanide Ions	6
1.1.3	Lanthanide Magnetism in Ligand Fields	8
1.2.1	Effective Barrier (U_{eff})	11
1.2.2	Blocking Temperature (T_B) and Hysteresis	18
1.2.3	Relaxation Dynamics	19
1.3.1	Angle-Resolved Magnetometry	27
1.3.2	Ab Initio Calculations	32
References		34
2	Lanthanide Single-Ion Molecular Magnets	41
2.1	Single-Ion Anisotropy and QTM	42
2.2	[LnPc ₂] and LnPOM SIMMs	46
2.2.1	Phthalocyanine-Type Ligands and the [LnPc ₂] Structure	46
2.2.2	The Magnetism of Seminal [LnPc ₂] ⁻ ·TBA ⁺	48
2.2.3	The Enhanced SMMs from the Redox Reactions	53
2.2.4	The Substituted [LnPc ₂] SIMMs	57
2.2.5	The [LnPc ₂]-like SIMMs	60
2.2.6	The LnPOM SIMMs	62
2.3	Lanthanide β-diketone Systems	64
2.3.1	The Magnetism of [Dy(acac) ₃ (H ₂ O) ₂]	66
2.3.2	The Modulations of Auxiliary Ligands	67
2.3.3	Chiral β-diketone Dy SIMMs	72
2.3.4	The SIMMs with Large Capping Ligands	73
2.3.5	The SIMMs with Radical Ligands	77

2.4	Organometallic SIMMs	80
2.5	Conclusion	84
	References	84
3	Dinuclear Lanthanide Single-Molecule Magnets	91
3.1	Magnetic Interactions	92
3.2	The Coupling of Two SIMMs	97
3.2.1	Triple-Decker Lanthanide Phthalocyanine System	98
3.2.2	Dinuclear β -Diketone SMMs	102
3.2.3	Dinuclear Lanthanide Radical SMMs	108
3.2.4	Organometallic Multidecker Ln-COT SMMs	110
3.3	The Choice of Bridging Ligands	112
3.4	Dinuclear SMMs with O-Bridges	116
3.4.1	Carboxylato-Bridged Dy_2 SMMs	118
3.4.2	Phenoxido-Bridged Dy_2 SMMs	120
3.4.3	Alkoxido-Bridged Dy_2 SMMs	122
3.5	Conclusion	123
	References	124
4	Single-Molecule Toroics and Multinuclear Lanthanide Single-Molecule Magnets	127
4.1	Single-Molecule Toroic	128
4.1.1	The Seminal Dy_3 SMT	130
4.1.2	New Dy_3 -SMTs Systems	132
4.1.3	Cyclic Dy_4 and Dy_6 SMT Systems	135
4.1.4	Coupled Dy_3 -SMTs Systems	137
4.1.5	Other Dy_3 Examples	142
4.2	High-Performance Multinuclear Lanthanide SMMs	144
4.2.1	Building Block Approach	145
4.2.2	Organometallic Approach	147
4.2.3	Multidentate Ligand Approach	150
	References	163
5	Hydrazone-Based Lanthanide Single-Molecule Magnets	167
5.1	Hydrazone Ligands	168
5.2	Two Model Complexes for Two-Step Relaxation	170
5.3	Hydrazone Ligands Based on Different Hydrazide Parts	172
5.3.1	Benzhydrazone Ligands	172
5.3.2	Isonicotinohydrazone Ligands	175
5.3.3	Picolinohydrazone Ligands	177
5.3.4	Pyrazine Acylhydrazone Ligands	179
5.3.5	Hydroxymethyl Picolinohydrazone Ligands	181
5.3.6	<i>o</i> -Vanillyl Hydrazone Ligands	183

5.3.7	Hydrazone–Oxime Ligands	186
5.3.8	Hydrazone Ligands from Bis-Acylhydrazide.	188
References		191
6	Conclusion and Perspective	195
6.1	New Type of Lanthanide SMM— $Dy_nSc_{3-n}N@C_{80}$ ($n = 1, 2, 3$)	196
6.2	The Organization of SMMs on Surfaces	198
6.2.1	Transition Metal SMMs on Au Surface	199
6.2.2	[LnPc ₂] SMMs on Surfaces	201
6.3	Molecular Spintronics	204
References		208



Jinkui Tang received his Ph.D. degree in chemistry from Nankai University under the guidance of Prof. De-Zheng Liao and then worked as a postdoctoral fellow at Universität Karlsruhe (TH) in the research group of Prof. Amic K. Powell and Universiteit Leiden with Prof. Jan Reedijk, before initiating his independent career at Changchun Institute of Applied Chemistry, Chinese Academy of Sciences in 2007. His current research focuses on the synthesis and characterization of new molecular magnetic materials (including single-molecule-magnets and single-molecule forces) based on highly anisotropic lanthanide ions. His work has resulted in about 150 peer-reviewed publications.



Peng Zhang obtained his B.Sc. in chemistry from Inner Mongolia University, China, in 2010. Now he is a Ph.D. student supervised by Prof. Jinkui Tang at Changchun Institute of Applied Chemistry, Chinese Academy of Sciences (CAS). His current work focuses on lanthanide-based molecular magnetism.