

Contents

| | |
|---------------------------------|----|
| Foreword | 13 |
| Acknowledgements | 15 |
| Introduction (A. Lebeda) | 17 |

Scientific contributions

I. General aspects, culture and management

| | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| J. Moravec, A. Lebeda and E. Kříštková (Czech Republic) History of growing and breeding of cucurbitaceous vegetables in Czech Lands | 21 |
| J. Lužný, A. Lebeda and E. Kříštková (Czech Republic) A dedication to Franz Frimmel, a Czech leader of cucurbit breeding | 39 |
| L. Nowaczyk and P. Nowaczyk (Poland) Cucumber fruit size and seed yield affected by growth regulators | 45 |
| N. Biriukova and E. Maslovskaya (Russia) The influence of cultivation conditions on parthenocarpy of cucumber | 51 |
| G. Baysal, R. Tipirdamaz and Y. Ekmekci (Turkey) Effects of salinity on some physiological parameters in three cultivars of cucumber (<i>Cucumis sativus</i>) | 57 |
| M. Edelstein, M. Ben-Hur, R. Cohen, Y. Burger and I. Ravina (Israel) Comparison of grafted and non-grafted melon plants under excess of boron and salinity stress | 63 |
| A.M. Granero, J.M.G. Sanz, J.L.M. Vidal, A.G. Frenich, A.R. Serrano and F.J.E. González (Spain) Sugars and volatiles composition of nectar of zucchini flowers | 69 |
| J.M.G. Sanz, A.R. Serrano and A.M. Granero (Spain) Pollination of zucchini culture by bumblebees: Advance of results of quality production | 75 |
| | |
| <i>II. Germplasm</i> | |
| V.S. Seshadri and T.A. More (India) History and antiquity of cucurbits in India | 81 |

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| W. van Dooijeweert (the Netherlands) The status of the cucumber (<i>Cucumis sativus</i>) collection of CGN | 91 |
| V. Vinter, A. Kříštková, A. Lebeda and E. Kříštková (Czech Republic) Descriptor lists for genetic resources of the genus <i>Cucumis</i> and cultivated species of the genus <i>Cucurbita</i> | 95 |
| H. Pavlikaki, C. Ponce Navarro and N. Fanourakis (Greece) Genetic relationships of different Greek landraces of cucumber (<i>Cucumis sativus</i>) as assessed by RAPDs | 101 |
| T.C. Andres (USA) Diversity in tropical pumpkin (<i>Cucurbita moschata</i>): a review of infraspecific classifications | 107 |
| T.C. Andres (USA) Diversity in tropical pumpkin (<i>Cucurbita moschata</i>): cultivar origin and history | 113 |
| E. Kříštková, A. Kříštková and V. Vinter (Czech Republic) Morphological variation of cultivated <i>Cucurbita</i> species | 119 |
| M.L. Gomez-Guillamón, E. Moriones, M.S. Luis-Arteaga, V. Carnide, A. Börner, N. Sari, K. Abak and J.M. Alvarez (Spain, Portugal, Germany, Turkey) Management, conservation and valorization on genetic resources of <i>Cucumis melo</i> and wild relatives | 129 |
| V. Carnide, S. Martins, F.J. Vences, L.E. Sáenz de Miera and M.R. Barroso (Portugal, Spain) Evaluation of Portuguese melon landraces conserved on farm by morphological traits and RAPDs | 135 |
| M.R. Barroso, S. Martins, F.J. Vences, L.E. Sáenz de Miera and V. Carnide (Portugal, Spain) Comparative analysis of melon landraces from South Portugal using RAPD markers | 143 |
| Y. Burger, Y. Yeselson, U. Saar, H.S. Paris, N. Katzir, Y. Tadmor and A.A. Schaffer (Israel) Screening of melon (<i>Cucumis melo</i>) germplasm for consistently high sucrose content and for high ascorbic acid content | 151 |
| M.A. Queiroz, M.L. Silva, L.M. Silveira, R.C.S. Dias, M.A.J.F. Ferreira, S.R.R. Ramos, R.L. Romão, J.G.A. Assis, F.F. Souza and M.C.C.L. Moura (Brazil) Pre-breeding in the watermelon germplasm bank of the Northeast of Brazil | 157 |

| | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| M. Koutsika-Sotiriou, E. Traka-Mavrona, A.L. Tsivelikas, G. Mpardas, A. Mpeis and E. Klonari (Greece) Use of genetic resources in a dual approach toward selecting improved scion/rootstock grafting combinations of melon (<i>Cucumis melo</i>) on <i>Cucurbita</i> spp. | 163 |
| A. López-Sesé and J.E. Staub (USA) Effects of seed maturation and temperature in germination of squash accessions: implications for gene flow | 169 |
| M. Sugiyama, K. Sugiyama, T. Ohara, M. Morishita and Y. Sakata (Japan) Characteristics and inheritance of a high hermaphroditic flower-bearing accession of watermelon (<i>Citrullus lanatus</i>) | 175 |
| Y. Tadmor, N. Katzir, S. King, A. Levi, A. Davis and J. Hirschberg (Israel, USA) Fruit coloration in watermelon: lessons from the tomato | 181 |
| <i>III. Diseases and pests, disease resistance</i> | |
| J.F. Chen, G. Moriarty and M. Jahn (China, USA) Some disease resistance tests in <i>Cucumis hystrix</i> and its progenies from interspecific hybridization with cucumber | 189 |
| T. Montoro, S. Sánchez-Campos, R. Camero, C.F. Marco, P. Corella and M.L. Gómez-Guillamón (Spain) Searching for resistance to cucumber vein yellowing virus in <i>Cucumis melo</i> | 197 |
| C. Mallor, J.M. Álvarez and M. Luis-Arteaga (Spain) Behaviour of <i>Cucumis melo</i> 'Cantaloup Haogen' against melon necrotic spot virus (MNSV) | 203 |
| J. Garcia-Mas, M. Morales, H. van Leeuwen, A. Monfort, P. Puigdomenech, P. Arús, C. Nieto, M.A. Aranda, C. Dogimont, G. Orjeda, M. Caboche and A. Bendahmane (Spain, France) A physical map covering the <i>Nsv</i> locus in melon | 209 |
| B.S. Daryono, S. Somowiyarjo and K.T. Natsuaki (Japan, Indonesia) Detection of resistant melons to the Indonesian isolate of KGMMV | 213 |
| J.D. McCreight (USA) Progress in breeding melon for resistance to lettuce infectious yellows virus | 219 |
| N. Guner and T.C. Wehner (USA) Resistance to a severe strain of zucchini yellow mosaic virus in watermelon | 223 |

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| J. Svoboda and J. Polák (Czech Republic) Preliminary evaluation of squash cultivars for resistance to a Czech isolate of zucchini yellow mosaic virus | 231 |
| M. Pachner and T. Lelley (Austria) Different genes for resistance to zucchini yellow mosaic virus (ZYMV) in <i>Cucurbita moschata</i> | 237 |
| H. Olczak-Woltman, M. Bakowska, M. Schollenberger and K. Niemirowicz-Szczytt (Poland) Cucumber screening for resistance to angular leaf spot | 245 |
| D.L. Hopkins (USA) Characteristics of resistance to <i>Acidovorax avenae</i> subsp. <i>citrullii</i> in the <i>Citrullus lanatus</i> accessions PI 482279 and PI 494817 | 251 |
| D. Kenigsbuch, D. Taler, M. Galperin, I. Benjamin and Y. Cohen (Israel) Plant <i>eR</i> genes encoding for glyoxalate aminotransferase enzymes confer resistance against downy mildew in melon | 257 |
| A. Lebeda and J. Urban (Czech Republic) Disease impact and pathogenicity variation in Czech populations of <i>Pseudoperonospora cubensis</i> | 267 |
| J. Urban and A. Lebeda (Czech Republic) Differential sensitivity to fungicides in Czech populations of <i>Pseudoperonospora cubensis</i> | 275 |
| A. Lebeda and B. Sedláková (Czech Republic) Disease impact and pathogenicity variation in Czech populations of cucurbit powdery mildews | 281 |
| B. Sedláková and A. Lebeda (Czech Republic) Variation in sensitivity to fungicides in Czech populations of cucurbit powdery mildews | 289 |
| T. Montoro, M. Salinas, J. Capel, M. Gómez-Guillamón and R. Lozano (Spain) Genetic variability in <i>Sphaerotheca fusca</i> as determined by AFLPs: the case of race 2 as a causal agent of powdery mildew in melon | 295 |
| R. Song, G. Gusmini and T.C. Wehner (China, USA) A summary of eleven preliminary studies of greenhouse and field testing methods for resistance to gummy stem blight in watermelon | 301 |
| L. Perche pied, C. Dogimont and M. Pitrat (France) Genetic analysis of resistance to <i>Fusarium oxysporum</i> f.sp. <i>melonis</i> race 1.2 in melon | 307 |

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| R. Cohen, Y. Burger, C. Horev, A. Porat, U. Saar and M. Edelstein (Israel) Reduction of <i>Monosporascus</i> wilt incidence using different Galia-type melons grafted onto <i>Cucurbita</i> rootstock | 313 |
| M. Grinberg, V. Soroker, E. Palevsky, I. Shomer and R. Perl-Treves (Israel) Response of cucumber to the broad mite (<i>Polyphagotarsonemus latus</i>) | 319 |
| J. Pauquet, E. Burget, L. Hagen, V. Chovelon, A. Le Menn, N. Valot, S. Desloire, M. Caboche, P. Rousselle, M. Pitrat, A. Bendahmane and C. Dogimont (France) | |
| Map-based cloning of the <i>Vat</i> gene from melon conferring resistance to both aphid colonization and aphid transmission of several viruses | 325 |
| IV. Breeding and genetics | |
| K. Bartoszak (Poland) | |
| Performance of pickling cucumber cultivars presently on the Polish National List | 333 |
| U. Kłosinska and E.U. Kozik (Poland) | |
| Suitability of new cucumber F ₁ hybrids for open-field cultivation | 337 |
| J. Berenji and D. Papp (Serbia, Hungary) | |
| The effect of ethephon on the sex expression of naked seeded oil pumpkin | 341 |
| N. Ercan, M. Temirkaynak, F. Şensoy and A.S. Şensoy (Turkey) | |
| Evaluation of some inbred lines of summer squash for plant, flower, fruit and seed properties | 345 |
| P. Gómez, A. Peñaranda, D. Garrido and M. Jamilena (Spain) | |
| Evaluation of flower abscission and sex expression in different cultivars of zucchini squash (<i>Cucurbita pepo</i>) | 347 |
| L. Perche pied, C. Périn, N. Giovinazzo, D. Besombes, C. Dogimont and M. Pitrat (France) | |
| Susceptibility to sulfur dusting and inheritance in melon | 353 |
| B. Kowalczyk (Poland) | |
| Distinctness, uniformity and stability testing of cucumber culti- vars in Poland | 359 |
| Z. Sun, R.L. Lower and J.E. Staub (USA) | |
| Generation means analysis of parthenocarpic characters in a pro- cessing cucumber (<i>Cucumis sativus</i>) population | 365 |
| J.E. Zalapa, J.E. Staub and J.D. McCreight (USA) | |
| Genetic analysis of branching in melon (<i>Cucumis melo</i>) | 373 |

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| J.E. Staub, J.E. Zalapa, M.K. Paris and J.D. McCreight (USA) Selection for lateral branch number in melon (<i>Cucumis melo</i>) | 381 |
| H.S. Paris, A. Hanan and F. Baumkoler (Israel) Assortment of five gene loci in <i>Cucurbita pepo</i> | 389 |
| L. Wessel-Beaver, H.E. Cuevas, T.C. Andres and D.R. Piperno (USA) Genetic compatibility between <i>Cucurbita moschata</i> and <i>C. argyrosperma</i> | 393 |
| M.D. Robbins and J.E. Staub (USA) Strategies for selection of multiple quantitatively inherited yield components in cucumber | 401 |
| V. Tissue culture, biotechnology, molecular genetics and mapping | |
| J. Sztangret, J. Wronka, T. Galecka, A. Korzeniewska and K. Niemirowicz-Szczytt (Poland) Cucumber (<i>Cucumis sativus</i>) haploids developed from parthenocarpic hybrids | 411 |
| D. Sklov, A. Lebeda and B. Navrtilov (Czech Republic) Embryo and ovule cultures in <i>Cucumis</i> species and their utilization in interspecific hybridization | 415 |
| H. Ezura and Y. Akasaka-Kennedy (Japan) Somatic embryogenesis in a model cultivar, PI 161375 (<i>Cucumis melo</i> subsp. <i>agrestis</i>), of melon | 431 |
| E. Kiss-Baba, S. Penczel, V. Zarka, Gy.D. Bisztray and I. Velich (Hungary) Regeneration ability of some Hungarian melon varieties | 437 |
| J. Gajdov, A. Lebeda and B. Navrtilov (Czech Republic) Protoplast cultures of <i>Cucumis</i> and <i>Cucurbita</i> spp. | 441 |
| G. Bartoszewski, S. Malepszy and M.J. Havey (Poland, USA) Mosaic (MSC) cucumbers regenerated from independent cell cultures possess different mitochondrial rearrangements | 455 |
| S. Curuk, S. Cetiner, C. Elman, X. Xia, Y. Wang, A. Yeheskel, L. Zilberstein, R. Perl-Treves, A.A. Watad and V. Gaba (Turkey, Israel) Wounding by vortexing with carborundum facilitates <i>Agrobacterium</i> -mediated transformation of melon (<i>Cucumis melo</i>) | 459 |
| A. AtarEs, B. Garcia-Sogo, B. Pineda, P. Ellul and V. Moreno (Spain) Transformation of melon via PEG-induced direct DNA uptake into protoplasts | 465 |

| | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Z. Gao, M. Petreikov, Y. Burger, S. Shen and A.A. Schaffer (Israel) Stachyose to sucrose metabolism in sweet melon (<i>Cucumis melo</i>) fruit mesocarp during the sucrose accumulation stage | 471 |
| S-M. Chung and J.E. Staub (USA) Consensus chloroplast primer analysis: A molecular tool for evolutionary studies in Cucurbitaceae | 477 |
| Y. Brotman, I. Kovalski, C. Dogimont, M. Pitrat, N. Katzir and R. Perl-Treves (Israel, France) Molecular mapping of the melon <i>Fom-1/Prv</i> locus | 485 |
| Y. Yariv, V. Portnoy, Y. Burger, Y. Benyamin, E. Lewinsohn, Y. Tadmor, U. Ravid, R. White, J. Giovannoni, A.A. Schaffer and N. Katzir (Israel, USA) Isolation and characterization of fruit-related genes in melon (<i>Cucumis melo</i>) using SSH and macroarray techniques | 491 |
| I. Eduardo, P. Ar's and A.J. Monforte (Spain) Genetics of fruit quality in melon. Verification of QTLs involved in fruit shape with near-isogenic lines (NILs) | 499 |
| N. Fukino, M. Kuzuya, M. Kunihisa and S. Matsumoto (Japan) Characterization of simple sequence repeats (SSRs) and development of SSR markers in melon (<i>Cucumis melo</i>) | 503 |
| A. Zraidi and T. Lelley (Austria) Genetic map for pumpkin <i>Cucurbita pepo</i> using random amplified polymorphic DNA markers | 507 |
| A. Levi, C.E. Thomas, J. Thies, A. Simmons, Y. Xu, X. Zhang, O.U.K. Reddy, Y. Tadmor, N. Katzir, T. Trebitsh, S. King, A. Davis, J. Fauve and T. Wehner (USA, China, Israel, France) Developing a genetic linkage map for watermelon: polymorphism, segregation and distribution of markers | 515 |
| P.A. Rajagopalan, T. Saraf-Levy, A. Lizhe and R. Perl-Treves (Israel, China) Increased femaleness in transgenic cucumbers that overexpress an ethylene receptor | 525 |
| Authors index | 533 |
| List of participants | 537 |