## Pflügers Archiv European Journal of Physiology

GENERAL INFORMATION > springeronline.com

ELECTRONIC CONTENT



Volume 450 • Number 6 • September 2005		Hede SE, Amstrup J, Klaerke DA, Novak I: P2Y <sub>2</sub> and P2Y <sub>4</sub> receptors regulate pancreatic Ca <sup>2+</sup> -activated K <sup>+</sup> channels differently	429
Invited Review	the state	channels directionly	
Cummins EP, Taylor CT:		Skeletal Muscle	
Hypoxia-responsive transcription factors	363	Whitman SA, Wacker MJ, Richmond SR, Godard MP: Contributions of the ubiquitin–proteasome pathway and	
Cardiovascular System		apoptosis to human skeletal muscle wasting with age	437
Balanos GM, Talbot NP, Robbins PA, Dorrington KL: Separating the direct effect of hypoxia from the indirect effect of changes in cardiac output on the maximum pressure		Acknowledgement to Referees	447
difference across the tricuspid valve in healthy humans	372	Abstracts	
Cell and Molecular Physiology		Belgian Society of Fundamental and Clinical Physiology and Pharmacology:	
Knott TK, Velázquez-Marrero C, Lemos JR:		Abstracts of the Spring Meeting	
ATP elicits inward currents in isolated vasopressinergic neurohypophysial terminals via P2X <sub>2</sub> and P2X <sub>3</sub> receptors	381	23 April 2005, Brussels, (Université Libre de Bruxelles), Belgium	R1
Endocrinology		Forthcoming articles in print	
Carnio EC, Stabile AM, Batalhão ME, Silva JS, Antunes-Rodrigues J, Branco LGS, Magder S: Vasopressin release during endotoxaemic shock in mice lacking		published online and citable with the DOI (Digital Object Identified be found at springerlink.com	er) can
inducible nitric oxide synthase	390		
Misler S, Dickey A, Barnett DW:			
Maintenance of stimulus-secretion coupling and single beta-cell			
function in cryopreserved-thawed human islets of Langerhans	395	the set of the	
Epithelial Transport			
Musa-Aziz R, Mello-Aires M:		and the second	
Action of ANG II and ANP on colon epithelial cells	405		
and the second second second second		Indexed in/abstracted by Current Contents, Index Medicus, and EMBASE/Excerpta Medica	
Ion Channels, Transporters	ni. Ni		St. Sala
Holm LM, Jahn TP, Møller ALB, Schjoerring JK, Ferri D, Klaerke DA, Zeuthen T:		Cover illustration: Diagram representing hypoxia responsive transcriptio tors. Included below each transcription factor, in brackets, are some of the	
NH <sub>3</sub> and NH <sub>4</sub> <sup>+</sup> permeability in aquaporin-expressing <i>Xenopus</i> oocytes	415	relevant references. For details, see article by Cummins and Taylor (2005) Pflügers Arch–Eur of Physiol 450: 363–371	
		A CONTRACT OF A CONTRACT.	



