

Contents

Introduction	7
I. Material and Methods	9
II. Evolution of parasitic trematodes from free-living protrematodes	12
III. Primitive life-cycle of trematodes (primary diheteroxenia)	25
1. Life-cycle without adolescaria	25
2. Life-cycle with adolescaria (models - <i>Notocotylus attenuatus</i> , <i>Philophthalmus</i> sp.)	25
IV. Three-host life-cycle of trematodes bound to water environment	36
1. Origin of tri- and tetraheteroxenia	36
2. Significance of the second intermediate host	39
3. Adaptation of cercaria to penetration into the second intermediate host, modification of cyst wall and differentiation of metacercaria (models - <i>Moliniella anceps</i> , <i>Echinoparyphium</i> <i>aconiatum</i> , <i>Plagiorchis laricola</i>)	40
V. Life-cycles of trematodes bound to terrestrial environment	53
1. Phylogensis of terrestrial snails and their adaptation to the existence on land	53
2. Adaptation of trematode larvae to parasitism in terrestrial snails	56
VI. Three-host life-cycle of trematodes bound to terrestrial environment (models - <i>Dicrocoelium</i> <i>lanceatum</i> , <i>Eurytrema pancreaticum</i> , <i>Corrigia</i> <i>corrigia</i> , <i>Brachylaimus aequans</i>)	60
VII. Secondary two-host life-cycles of trematodes bound to terrestrial environment (models - <i>Hasstilesia ovis</i> , <i>Leucochloridium paradoxum</i> , <i>L. perturbatum</i>)	79
VIII. Conclusion	88
IX. References	91