

# BRIEF CONTENTS

## Part I THE MOLECULAR DESIGN OF LIFE

- 1 Biochemistry: An Evolving Science 1
- 2 Protein Composition and Structure 27
- 3 Exploring Proteins and Proteomes 65
- 4 DNA, RNA, and the Flow of Genetic Information 105
- 5 Exploring Genes and Genomes 135
- 6 Exploring Evolution and Bioinformatics 169
- 7 Hemoglobin: Portrait of a Protein in Action 191
- 8 Enzymes: Basic Concepts and Kinetics 215
- 9 Catalytic Strategies 251
- 10 Regulatory Strategies 285
- 11 Carbohydrates 315
- 12 Lipids and Cell Membranes 341
- 13 Membrane Channels and Pumps 367
- 14 Signal-Transduction Pathways 397

## Part II TRANSDUCING AND STORING ENERGY

- 15 Metabolism: Basic Concepts and Design 423
- 16 Glycolysis and Gluconeogenesis 449
- 17 The Citric Acid Cycle 495
- 18 Oxidative Phosphorylation 523
- 19 The Light Reactions of Photosynthesis 565
- 20 The Calvin Cycle and the Pentose Phosphate Pathway 589
- 21 Glycogen Metabolism 617
- 22 Fatty Acid Metabolism 643
- 23 Protein Turnover and Amino Acid Catabolism 681

## Part III SYNTHESIZING THE MOLECULES OF LIFE

- 24 The Biosynthesis of Amino Acids 713
- 25 Nucleotide Biosynthesis 743
- 26 The Biosynthesis of Membrane Lipids and Steroids 767
- 27 The Integration of Metabolism 801
- 28 DNA Replication, Repair, and Recombination 827
- 29 RNA Synthesis and Processing 859
- 30 Protein Synthesis 893
- 31 The Control of Gene Expression in Prokaryotes 925
- 32 The Control of Gene Expression in Eukaryotes 941

## Part IV RESPONDING TO ENVIRONMENTAL CHANGES

- 33 Sensory Systems 961
- 34 The Immune System 981
- 35 Molecular Motors 1011
- 36 Drug Development 1033

# CONTENTS

Preface v

## Part I THE MOLECULAR DESIGN OF LIFE 1

### CHAPTER 1 Biochemistry: An Evolving Science 1

- 1.1 Biochemical Unity Underlies Biological Diversity 1
- 1.2 DNA Illustrates the Interplay Between Form and Function 4
  - DNA is constructed from four building blocks 4
  - Two single strands of DNA combine to form a double helix 5
  - DNA structure explains heredity and the storage of information 5
- 1.3 Concepts from Chemistry Explain the Properties of Biological Molecules 6
  - The formation of the DNA double helix as a key example 6
  - The double helix can form from its component strands 6
  - Covalent and noncovalent bonds are important for the structure and stability of biological molecules 6
  - The double helix is an expression of the rules of chemistry 9
  - The laws of thermodynamics govern the behavior of biochemical systems 10
  - Heat is released in the formation of the double helix 12
  - Acid–base reactions are central in many biochemical processes 13
  - Acid–base reactions can disrupt the double helix 14
  - Buffers regulate pH in organisms and in the laboratory 15
- 1.4 The Genomic Revolution Is Transforming Biochemistry, Medicine, and Other Fields 17
  - Genome sequencing has transformed biochemistry and other fields 17
  - Environmental factors influence human biochemistry 20
  - Genome sequences encode proteins and patterns of expression 21
- APPENDIX: Visualizing Molecular Structures I: Small Molecules 22

### CHAPTER 2 Protein Composition and Structure 27

- 2.1 Proteins Are Built from a Repertoire of 20 Amino Acids 29
- 2.2 Primary Structure: Amino Acids Are Linked by Peptide Bonds to Form Polypeptide Chains 35
  - Proteins have unique amino acid sequences specified by genes 37
  - Polypeptide chains are flexible yet conformationally restricted 38
- 2.3 Secondary Structure: Polypeptide Chains Can Fold into Regular Structures Such As the Alpha Helix, the Beta Sheet, and Turns and Loops 40
  - The alpha helix is a coiled structure stabilized by intrachain hydrogen bonds 40
  - Beta sheets are stabilized by hydrogen bonding between polypeptide strands 42



Polypeptide chains can change direction by making reverse turns and loops	44	<b>3.3 Mass Spectrometry Is a Powerful Technique for the Identification of Peptides and Proteins</b>	<b>85</b>
Fibrous proteins provide structural support for cells and tissues	44	Peptides can be sequenced by mass spectrometry	87
<b>2.4 Tertiary Structure: Water-Soluble Proteins Fold into Compact Structures with Nonpolar Cores</b>	<b>46</b>	Proteins can be specifically cleaved into small peptides to facilitate analysis	88
<b>2.5 Quaternary Structure: Polypeptide Chains Can Assemble into Multisubunit Structures</b>	<b>48</b>	Genomic and proteomic methods are complementary	89
<b>2.6 The Amino Acid Sequence of a Protein Determines Its Three-Dimensional Structure</b>	<b>49</b>	The amino acid sequence of a protein provides valuable information	90
Amino acids have different propensities for forming $\alpha$ helices, $\beta$ sheets, and turns	51	Individual proteins can be identified by mass spectrometry	91
Protein folding is a highly cooperative process	52	<b>3.4 Peptides Can Be Synthesized by Automated Solid-Phase Methods</b>	<b>92</b>
Proteins fold by progressive stabilization of intermediates rather than by random search	53	<b>3.5 Three-Dimensional Protein Structure Can Be Determined by X-ray Crystallography and NMR Spectroscopy</b>	<b>95</b>
Prediction of three-dimensional structure from sequence remains a great challenge	54	X-ray crystallography reveals three-dimensional structure in atomic detail	95
Some proteins are inherently unstructured and can exist in multiple conformations	55	Nuclear magnetic resonance spectroscopy can reveal the structures of proteins in solution	97
Protein misfolding and aggregation are associated with some neurological diseases	56	<b>CHAPTER 4 DNA, RNA, and the Flow of Genetic Information</b>	<b>105</b>
Protein modification and cleavage confer new capabilities	57	<b>4.1 A Nucleic Acid Consists of Four Kinds of Bases Linked to a Sugar-Phosphate Backbone</b>	<b>106</b>
APPENDIX: Visualizing Molecular Structures II: Proteins	61	RNA and DNA differ in the sugar component and one of the bases	106
<b>CHAPTER 3 Exploring Proteins and Proteomes</b>	<b>65</b>	Nucleotides are the monomeric units of nucleic acids	107
The proteome is the functional representation of the genome	66	DNA molecules are very long and have directionality	108
<b>3.1 The Purification of Proteins Is an Essential First Step in Understanding Their Function</b>	<b>66</b>	<b>4.2 A Pair of Nucleic Acid Strands with Complementary Sequences Can Form a Double-Helical Structure</b>	<b>109</b>
The assay: How do we recognize the protein that we are looking for?	67	The double helix is stabilized by hydrogen bonds and van der Waals interactions	109
Proteins must be released from the cell to be purified	67	DNA can assume a variety of structural forms	111
Proteins can be purified according to solubility, size, charge, and binding affinity	68	Z-DNA is a left-handed double helix in which backbone phosphates zigzag	112
Proteins can be separated by gel electrophoresis and displayed	71	Some DNA molecules are circular and supercoiled	113
A protein purification scheme can be quantitatively evaluated	75	Single-stranded nucleic acids can adopt elaborate structures	113
Ultracentrifugation is valuable for separating biomolecules and determining their masses	76	<b>4.3 The Double Helix Facilitates the Accurate Transmission of Hereditary Information</b>	<b>114</b>
Protein purification can be made easier with the use of recombinant DNA technology	78	Differences in DNA density established the validity of the semiconservative replication hypothesis	115
<b>3.2 Immunology Provides Important Techniques with Which to Investigate Proteins</b>	<b>79</b>	The double helix can be reversibly melted	116
Antibodies to specific proteins can be generated	79	<b>4.4 DNA Is Replicated by Polymerases That Take Instructions from Templates</b>	<b>117</b>
Monoclonal antibodies with virtually any desired specificity can be readily prepared	80	DNA polymerase catalyzes phosphodiester-bridge formation	117
Proteins can be detected and quantified by using an enzyme-linked immunosorbent assay	82	The genes of some viruses are made of RNA	118
Western blotting permits the detection of proteins separated by gel electrophoresis	83	<b>4.5 Gene Expression Is the Transformation of DNA Information into Functional Molecules</b>	<b>119</b>
Fluorescent markers make the visualization of proteins in the cell possible	84	Several kinds of RNA play key roles in gene expression	119



All cellular RNA is synthesized by RNA polymerases	120
RNA polymerases take instructions from DNA templates	121
Transcription begins near promoter sites and ends at terminator sites	122
Transfer RNAs are the adaptor molecules in protein synthesis	123

#### 4.6 Amino Acids Are Encoded by Groups of Three Bases Starting from a Fixed Point 124

Major features of the genetic code	125
Messenger RNA contains start and stop signals for protein synthesis	126
The genetic code is nearly universal	126

#### 4.7 Most Eukaryotic Genes Are Mosaics of Introns and Exons 127

RNA processing generates mature RNA	127
Many exons encode protein domains	128

### CHAPTER 5 Exploring Genes and Genomes 135

#### 5.1 The Exploration of Genes Relies on Key Tools 136

Restriction enzymes split DNA into specific fragments	137
Restriction fragments can be separated by gel electrophoresis and visualized	137
DNA can be sequenced by controlled termination of replication	138
DNA probes and genes can be synthesized by automated solid-phase methods	139
Selected DNA sequences can be greatly amplified by the polymerase chain reaction	141
PCR is a powerful technique in medical diagnostics, forensics, and studies of molecular evolution	142
The tools for recombinant DNA technology have been used to identify disease-causing mutations	143

#### 5.2 Recombinant DNA Technology Has Revolutionized All Aspects of Biology 143

Restriction enzymes and DNA ligase are key tools in forming recombinant DNA molecules	143
Plasmids and $\lambda$ phage are choice vectors for DNA cloning in bacteria	144
Bacterial and yeast artificial chromosomes	147
Specific genes can be cloned from digests of genomic DNA	147
Complementary DNA prepared from mRNA can be expressed in host cells	149
Proteins with new functions can be created through directed changes in DNA	150
Recombinant methods enable the exploration of the functional effects of disease-causing mutations	152

#### 5.3 Complete Genomes Have Been Sequenced and Analyzed 152

The genomes of organisms ranging from bacteria to multicellular eukaryotes have been sequenced	153
The sequence of the human genome has been completed	154

Next-generation sequencing methods enable the rapid determination of a complete genome sequence	155
Comparative genomics has become a powerful research tool	156

#### 5.4 Eukaryotic Genes Can Be Quantitated and Manipulated with Considerable Precision 157

Gene-expression levels can be comprehensively examined	157
New genes inserted into eukaryotic cells can be efficiently expressed	159
Transgenic animals harbor and express genes introduced into their germ lines	160
Gene disruption and genome editing provide clues to gene function and opportunities for new therapies	160
RNA interference provides an additional tool for disrupting gene expression	162
Tumor-inducing plasmids can be used to introduce new genes into plant cells	163
Human gene therapy holds great promise for medicine	164

### CHAPTER 6 Exploring Evolution and Bioinformatics 169

#### 6.1 Homologs Are Descended from a Common Ancestor 170

#### 6.2 Statistical Analysis of Sequence Alignments Can Detect Homology 171

The statistical significance of alignments can be estimated by shuffling	173
Distant evolutionary relationships can be detected through the use of substitution matrices	174
Databases can be searched to identify homologous sequences	177

#### 6.3 Examination of Three-Dimensional Structure Enhances Our Understanding of Evolutionary Relationships 177

Tertiary structure is more conserved than primary structure	178
Knowledge of three-dimensional structures can aid in the evaluation of sequence alignments	179
Repeated motifs can be detected by aligning sequences with themselves	180
Convergent evolution illustrates common solutions to biochemical challenges	181
Comparison of RNA sequences can be a source of insight into RNA secondary structures	182

#### 6.4 Evolutionary Trees Can Be Constructed on the Basis of Sequence Information 183

Horizontal gene transfer events may explain unexpected branches of the evolutionary tree	184
--	-----

#### 6.5 Modern Techniques Make the Experimental Exploration of Evolution Possible 185

Ancient DNA can sometimes be amplified and sequenced	185
Molecular evolution can be examined experimentally	185



**CHAPTER 7 Hemoglobin: Portrait of a Protein in Action** 191**7.1 Myoglobin and Hemoglobin Bind Oxygen at Iron Atoms in Heme** 192

Changes in heme electronic structure upon oxygen binding are the basis for functional imaging studies 193

The structure of myoglobin prevents the release of reactive oxygen species 194

Human hemoglobin is an assembly of four myoglobin-like subunits 195

**7.2 Hemoglobin Binds Oxygen Cooperatively** 195

Oxygen binding markedly changes the quaternary structure of hemoglobin 197

Hemoglobin cooperativity can be potentially explained by several models 198

Structural changes at the heme groups are transmitted to the  $\alpha_1\beta_1$ - $\alpha_2\beta_2$  interface 200

2,3-Bisphosphoglycerate in red cells is crucial in determining the oxygen affinity of hemoglobin 200

Carbon monoxide can disrupt oxygen transport by hemoglobin 201

**7.3 Hydrogen Ions and Carbon Dioxide Promote the Release of Oxygen: The Bohr Effect** 202**7.4 Mutations in Genes Encoding Hemoglobin Subunits Can Result in Disease** 204

Sickle-cell anemia results from the aggregation of mutated deoxyhemoglobin molecules 205

Thalassemia is caused by an imbalanced production of hemoglobin chains 207

The accumulation of free alpha-hemoglobin chains is prevented 207

Additional globins are encoded in the human genome 208

**APPENDIX: Binding Models Can Be Formulated in Quantitative Terms: The Hill Plot and the Concerted Model** 210**CHAPTER 8 Enzymes: Basic Concepts and Kinetics** 215**8.1 Enzymes are Powerful and Highly Specific Catalysts** 216

Many enzymes require cofactors for activity 217

Enzymes can transform energy from one form into another 217

**8.2 Gibbs Free Energy Is a Useful Thermodynamic Function for Understanding Enzymes** 218

The free-energy change provides information about the spontaneity but not the rate of a reaction 218

The standard free-energy change of a reaction is related to the equilibrium constant 219

Enzymes alter only the reaction rate and not the reaction equilibrium 220

**8.3 Enzymes Accelerate Reactions by Facilitating the Formation of the Transition State** 221

The formation of an enzyme-substrate complex is the first step in enzymatic catalysis 222

The active sites of enzymes have some common features 223

The binding energy between enzyme and substrate is important for catalysis 225

**8.4 The Michaelis-Menten Model Accounts for the Kinetic Properties of Many Enzymes** 225

Kinetics is the study of reaction rates 225

The steady-state assumption facilitates a description of enzyme kinetics 226

Variations in  $K_M$  can have physiological consequences 228 $K_M$  and  $V_{max}$  values can be determined by several means 228 $K_M$  and  $V_{max}$  values are important enzyme characteristics 229 $k_{cat}/K_M$  is a measure of catalytic efficiency 230

Most biochemical reactions include multiple substrates 231

Allosteric enzymes do not obey Michaelis-Menten kinetics 233

**8.5 Enzymes Can Be Inhibited by Specific Molecules** 234

The different types of reversible inhibitors are kinetically distinguishable 235

Irreversible inhibitors can be used to map the active site 237

Penicillin irreversibly inactivates a key enzyme in bacterial cell-wall synthesis 239

Transition-state analogs are potent inhibitors of enzymes 240

Catalytic antibodies demonstrate the importance of selective binding of the transition state to enzymatic activity 241

**8.6 Enzymes Can Be Studied One Molecule at a Time** 242**APPENDIX: Enzymes are Classified on the Basis of the Types of Reactions That They Catalyze** 245**CHAPTER 9 Catalytic Strategies** 251

A few basic catalytic principles are used by many enzymes 252

**9.1 Proteases Facilitate a Fundamentally Difficult Reaction** 253

Chymotrypsin possesses a highly reactive serine residue 253

Chymotrypsin action proceeds in two steps linked by a covalently bound intermediate 254

Serine is part of a catalytic triad that also includes histidine and aspartate 255

Catalytic triads are found in other hydrolytic enzymes 258

The catalytic triad has been dissected by site-directed mutagenesis 260

Cysteine, aspartyl, and metalloproteases are other major classes of peptide-cleaving enzymes 260

Protease inhibitors are important drugs 263



**9.2 Carbonic Anhydrases Make a Fast Reaction Faster** 264

- Carbonic anhydrase contains a bound zinc ion essential for catalytic activity 265
- Catalysis entails zinc activation of a water molecule 265
- A proton shuttle facilitates rapid regeneration of the active form of the enzyme 267

**9.3 Restriction Enzymes Catalyze Highly Specific DNA-Cleavage Reactions** 269

- Cleavage is by in-line displacement of 3'-oxygen from phosphorus by magnesium-activated water 269
- Restriction enzymes require magnesium for catalytic activity 271
- The complete catalytic apparatus is assembled only within complexes of cognate DNA molecules, ensuring specificity 272
- Host-cell DNA is protected by the addition of methyl groups to specific bases 274
- Type II restriction enzymes have a catalytic core in common and are probably related by horizontal gene transfer 275

**9.4 Myosins Harness Changes in Enzyme Conformation to Couple ATP Hydrolysis to Mechanical Work** 275

- ATP hydrolysis proceeds by the attack of water on the gamma-phosphoryl group 276
- Formation of the transition state for ATP hydrolysis is associated with a substantial conformational change 277
- The altered conformation of myosin persists for a substantial period of time 278
- Scientists can watch single molecules of myosin move 279
- Myosins are a family of enzymes containing P-loop structures 280

**CHAPTER 10 Regulatory Strategies** 285**10.1 Aspartate Transcarbamoylase Is Allosterically Inhibited by the End Product of Its Pathway** 286

- Allosterically regulated enzymes do not follow Michaelis–Menten kinetics 287
- ATCase consists of separable catalytic and regulatory subunits 287
- Allosteric interactions in ATCase are mediated by large changes in quaternary structure 288
- Allosteric regulators modulate the T-to-R equilibrium 291

**10.2 Isozymes Provide a Means of Regulation Specific to Distinct Tissues and Developmental Stages** 292**10.3 Covalent Modification Is a Means of Regulating Enzyme Activity** 293

- Kinases and phosphatases control the extent of protein phosphorylation 294
- Phosphorylation is a highly effective means of regulating the activities of target proteins 296

Cyclic AMP activates protein kinase A by altering the quaternary structure 297

ATP and the target protein bind to a deep cleft in the catalytic subunit of protein kinase A 298

**10.4 Many Enzymes Are Activated by Specific Proteolytic Cleavage** 299

- Chymotrypsinogen is activated by specific cleavage of a single peptide bond 299
- Proteolytic activation of chymotrypsinogen leads to the formation of a substrate-binding site 300
- The generation of trypsin from trypsinogen leads to the activation of other zymogens 301
- Some proteolytic enzymes have specific inhibitors 302
- Blood clotting is accomplished by a cascade of zymogen activations 303
- Prothrombin requires a vitamin K-dependent modification for activation 304
- Fibrinogen is converted by thrombin into a fibrin clot 304
- Vitamin K is required for the formation of  $\gamma$ -carboxyglutamate 306
- The clotting process must be precisely regulated 307
- Hemophilia revealed an early step in clotting 308

**CHAPTER 11 Carbohydrates** 315**11.1 Monosaccharides Are the Simplest Carbohydrates** 316

- Many common sugars exist in cyclic forms 318
- Pyranose and furanose rings can assume different conformations 320
- Glucose is a reducing sugar 321
- Monosaccharides are joined to alcohols and amines through glycosidic bonds 322
- Phosphorylated sugars are key intermediates in energy generation and biosyntheses 322

**11.2 Monosaccharides Are Linked to Form Complex Carbohydrates** 323

- Sucrose, lactose, and maltose are the common disaccharides 323
- Glycogen and starch are storage forms of glucose 324
- Cellulose, a structural component of plants, is made of chains of glucose 324

**11.3 Carbohydrates Can Be Linked to Proteins to Form Glycoproteins** 325

- Carbohydrates can be linked to proteins through asparagine (*N*-linked) or through serine or threonine (*O*-linked) residues 326
- The glycoprotein erythropoietin is a vital hormone 327
- Glycosylation functions in nutrient sensing 327
- Proteoglycans, composed of polysaccharides and protein, have important structural roles 327
- Proteoglycans are important components of cartilage 328
- Mucins are glycoprotein components of mucus 329
- Protein glycosylation takes place in the lumen of the endoplasmic reticulum and in the Golgi complex 330



Specific enzymes are responsible for oligosaccharide assembly	331
Blood groups are based on protein glycosylation patterns	331
Errors in glycosylation can result in pathological conditions	332
Oligosaccharides can be “sequenced”	332

#### 11.4 Lectins Are Specific Carbohydrate-Binding

<b>Proteins</b>	<b>333</b>
Lectins promote interactions between cells	334
Lectins are organized into different classes	334
Influenza virus binds to sialic acid residues	335

### CHAPTER 12 Lipids and Cell Membranes 341

Many common features underlie the diversity of biological membranes	342
---	-----

#### 12.1 Fatty Acids Are Key Constituents of Lipids 342

Fatty acid names are based on their parent hydrocarbons	342
Fatty acids vary in chain length and degree of unsaturation	343

#### 12.2 There Are Three Common Types of Membrane Lipids 344

Phospholipids are the major class of membrane lipids	344
Membrane lipids can include carbohydrate moieties	345
Cholesterol is a lipid based on a steroid nucleus	346
Archaeal membranes are built from ether lipids with branched chains	346
A membrane lipid is an amphipathic molecule containing a hydrophilic and a hydrophobic moiety	347

#### 12.3 Phospholipids and Glycolipids Readily Form Bimolecular Sheets in Aqueous Media 348

Lipid vesicles can be formed from phospholipids	348
Lipid bilayers are highly impermeable to ions and most polar molecules	349

#### 12.4 Proteins Carry Out Most Membrane Processes 350

Proteins associate with the lipid bilayer in a variety of ways	351
Proteins interact with membranes in a variety of ways	351
Some proteins associate with membranes through covalently attached hydrophobic groups	354
Transmembrane helices can be accurately predicted from amino acid sequences	354

#### 12.5 Lipids and Many Membrane Proteins Diffuse Rapidly in the Plane of the Membrane 356

The fluid mosaic model allows lateral movement but not rotation through the membrane	357
Membrane fluidity is controlled by fatty acid composition and cholesterol content	357
Lipid rafts are highly dynamic complexes formed between cholesterol and specific lipids	358
All biological membranes are asymmetric	358

### CHAPTER 13 Membrane Channels and Pumps 367

The expression of transporters largely defines the metabolic activities of a given cell type	368
--	-----

#### 13.1 The Transport of Molecules Across a Membrane May Be Active or Passive 368

Many molecules require protein transporters to cross membranes	368
Free energy stored in concentration gradients can be quantified	369

#### 13.2 Two Families of Membrane Proteins Use ATP Hydrolysis to Pump Ions and Molecules Across Membranes 370

P-type ATPases couple phosphorylation and conformational changes to pump calcium ions across membranes	370
--	-----

Digitalis specifically inhibits the $\text{Na}^+ - \text{K}^+$ pump by blocking its dephosphorylation	373
---	-----

P-type ATPases are evolutionarily conserved and play a wide range of roles	374
--	-----

Multidrug resistance highlights a family of membrane pumps with ATP-binding cassette domains	374
--	-----

#### 13.3 Lactose Permease Is an Archetype of Secondary Transporters That Use One Concentration Gradient to Power the Formation of Another 376

#### 13.4 Specific Channels Can Rapidly Transport Ions Across Membranes 378

Action potentials are mediated by transient changes in $\text{Na}^+$ and $\text{K}^+$ permeability	378
--	-----

Patch-clamp conductance measurements reveal the activities of single channels	379
---	-----

The structure of a potassium ion channel is an archetype for many ion-channel structures	379
--	-----

The structure of the potassium ion channel reveals the basis of ion specificity	380
---	-----

The structure of the potassium ion channel explains its rapid rate of transport	383
---	-----

Voltage gating requires substantial conformational changes in specific ion-channel domains	383
--	-----

A channel can be inactivated by occlusion of the pore: the ball-and-chain model	384
---	-----

The acetylcholine receptor is an archetype for ligand-gated ion channels	385
--	-----

Action potentials integrate the activities of several ion channels working in concert	387
---	-----

Disruption of ion channels by mutations or chemicals can be potentially life-threatening	388
--	-----

#### 13.5 Gap Junctions Allow Ions and Small Molecules to Flow Between Communicating Cells 389

#### 13.6 Specific Channels Increase the Permeability of Some Membranes to Water 390



<b>CHAPTER 14 Signal-Transduction Pathways</b>	<b>397</b>		
Signal transduction depends on molecular circuits	398		
<b>14.1 Heterotrimeric G Proteins Transmit Signals and Reset Themselves</b>	<b>399</b>		
Ligand binding to 7TM receptors leads to the activation of heterotrimeric G proteins	400		
Activated G proteins transmit signals by binding to other proteins	402		
Cyclic AMP stimulates the phosphorylation of many target proteins by activating protein kinase A	403		
G proteins spontaneously reset themselves through GTP hydrolysis	403		
Some 7TM receptors activate the phosphoinositide cascade	404		
Calcium ion is a widely used second messenger	405		
Calcium ion often activates the regulatory protein calmodulin	407		
<b>14.2 Insulin Signaling: Phosphorylation Cascades Are Central to Many Signal-Transduction Processes</b>	<b>407</b>		
The insulin receptor is a dimer that closes around a bound insulin molecule	408		
Insulin binding results in the cross-phosphorylation and activation of the insulin receptor	408		
The activated insulin-receptor kinase initiates a kinase cascade	409		
Insulin signaling is terminated by the action of phosphatases	411		
<b>14.3 EGF Signaling: Signal-Transduction Pathways Are Poised to Respond</b>	<b>411</b>		
EGF binding results in the dimerization of the EGF receptor	411		
The EGF receptor undergoes phosphorylation of its carboxyl-terminal tail	413		
EGF signaling leads to the activation of Ras, a small G protein	413		
Activated Ras initiates a protein kinase cascade	414		
EGF signaling is terminated by protein phosphatases and the intrinsic GTPase activity of Ras	414		
<b>14.4 Many Elements Recur with Variation in Different Signal-Transduction Pathways</b>	<b>415</b>		
<b>14.5 Defects in Signal-Transduction Pathways Can Lead to Cancer and Other Diseases</b>	<b>416</b>		
Monoclonal antibodies can be used to inhibit signal-transduction pathways activated in tumors	416		
Protein kinase inhibitors can be effective anticancer drugs	417		
Cholera and whooping cough are the result of altered G-protein activity	417		
<b>Part II TRANSDUCING AND STORING ENERGY</b>			
<b>CHAPTER 15 Metabolism: Basic Concepts and Design</b>	<b>423</b>		
<b>15.1 Metabolism Is Composed of Many Coupled, Interconnecting Reactions</b>	<b>424</b>		
Metabolism consists of energy-yielding and energy-requiring reactions	424	A thermodynamically unfavorable reaction can be driven by a favorable reaction	425
		<b>15.2 ATP Is the Universal Currency of Free Energy in Biological Systems</b>	<b>426</b>
		ATP hydrolysis is exergonic	426
		ATP hydrolysis drives metabolism by shifting the equilibrium of coupled reactions	427
		The high phosphoryl potential of ATP results from structural differences between ATP and its hydrolysis products	429
		Phosphoryl-transfer potential is an important form of cellular energy transformation	430
		<b>15.3 The Oxidation of Carbon Fuels Is an Important Source of Cellular Energy</b>	<b>432</b>
		Compounds with high phosphoryl-transfer potential can couple carbon oxidation to ATP synthesis	432
		Ion gradients across membranes provide an important form of cellular energy that can be coupled to ATP synthesis	433
		Phosphates play a prominent role in biochemical processes	434
		Energy from foodstuffs is extracted in three stages	434
		<b>15.4 Metabolic Pathways Contain Many Recurring Motifs</b>	<b>435</b>
		Activated carriers exemplify the modular design and economy of metabolism	435
		Many activated carriers are derived from vitamins	438
		Key reactions are reiterated throughout metabolism	440
		Metabolic processes are regulated in three principal ways	442
		Aspects of metabolism may have evolved from an RNA world	444
		<b>CHAPTER 16 Glycolysis and Gluconeogenesis</b>	<b>449</b>
		Glucose is generated from dietary carbohydrates	450
		Glucose is an important fuel for most organisms	451
		<b>16.1 Glycolysis Is an Energy-Conversion Pathway in Many Organisms</b>	<b>451</b>
		Hexokinase traps glucose in the cell and begins glycolysis	451
		Fructose 1,6-bisphosphate is generated from glucose 6-phosphate	453
		The six-carbon sugar is cleaved into two three-carbon fragments	454
		Mechanism: Triose phosphate isomerase salvages a three-carbon fragment	455
		The oxidation of an aldehyde to an acid powers the formation of a compound with high phosphoryl-transfer potential	457
		Mechanism: Phosphorylation is coupled to the oxidation of glyceraldehyde 3-phosphate by a thioester intermediate	458
		ATP is formed by phosphoryl transfer from 1,3-bisphosphoglycerate	459
		Additional ATP is generated with the formation of pyruvate	460
		Two ATP molecules are formed in the conversion of glucose into pyruvate	461



NAD <sup>+</sup> is regenerated from the metabolism of pyruvate	462	Flexible linkages allow lipoamide to move between different active sites	500
Fermentations provide usable energy in the absence of oxygen	464	<b>17.2 The Citric Acid Cycle Oxidizes Two-Carbon Units</b>	<b>501</b>
The binding site for NAD <sup>+</sup> is similar in many dehydrogenases	465	Citrate synthase forms citrate from oxaloacetate and acetyl coenzyme A	502
Fructose is converted into glycolytic intermediates by fructokinase	465	Mechanism: The mechanism of citrate synthase prevents undesirable reactions	502
Excessive fructose consumption can lead to pathological conditions	466	Citrate is isomerized into isocitrate	504
Galactose is converted into glucose 6-phosphate	466	Isocitrate is oxidized and decarboxylated to alpha-ketoglutarate	504
Many adults are intolerant of milk because they are deficient in lactase	467	Succinyl coenzyme A is formed by the oxidative decarboxylation of alpha-ketoglutarate	505
Galactose is highly toxic if the transferase is missing	468	A compound with high phosphoryl-transfer potential is generated from succinyl coenzyme A	505
<b>16.2 The Glycolytic Pathway Is Tightly Controlled</b>	<b>469</b>	Mechanism: Succinyl coenzyme A synthetase transforms types of biochemical energy	506
Glycolysis in muscle is regulated to meet the need for ATP	469	Oxaloacetate is regenerated by the oxidation of succinate	507
The regulation of glycolysis in the liver illustrates the biochemical versatility of the liver	472	The citric acid cycle produces high-transfer-potential electrons, ATP, and CO <sub>2</sub>	508
A family of transporters enables glucose to enter and leave animal cells	473	<b>17.3 Entry to the Citric Acid Cycle and Metabolism Through It Are Controlled</b>	<b>510</b>
Aerobic glycolysis is a property of rapidly growing cells	474	The pyruvate dehydrogenase complex is regulated allosterically and by reversible phosphorylation	511
Cancer and endurance training affect glycolysis in a similar fashion	476	The citric acid cycle is controlled at several points	512
<b>16.3 Glucose Can Be Synthesized from Noncarbohydrate Precursors</b>	<b>476</b>	Defects in the citric acid cycle contribute to the development of cancer	513
Gluconeogenesis is not a reversal of glycolysis	478	<b>17.4 The Citric Acid Cycle Is a Source of Biosynthetic Precursors</b>	<b>514</b>
The conversion of pyruvate into phosphoenolpyruvate begins with the formation of oxaloacetate	478	The citric acid cycle must be capable of being rapidly replenished	514
Oxaloacetate is shuttled into the cytoplasm and converted into phosphoenolpyruvate	480	The disruption of pyruvate metabolism is the cause of beriberi and poisoning by mercury and arsenic	515
The conversion of fructose 1,6-bisphosphate into fructose 6-phosphate and orthophosphate is an irreversible step	480	The citric acid cycle may have evolved from preexisting pathways	516
The generation of free glucose is an important control point	481	<b>17.5 The Glyoxylate Cycle Enables Plants and Bacteria to Grow on Acetate</b>	<b>516</b>
Six high-transfer-potential phosphoryl groups are spent in synthesizing glucose from pyruvate	481	<b>CHAPTER 18 Oxidative Phosphorylation</b>	<b>523</b>
<b>16.4 Gluconeogenesis and Glycolysis Are Reciprocally Regulated</b>	<b>482</b>	<b>18.1 Eukaryotic Oxidative Phosphorylation Takes Place in Mitochondria</b>	<b>524</b>
Energy charge determines whether glycolysis or gluconeogenesis will be most active	482	Mitochondria are bounded by a double membrane	524
The balance between glycolysis and gluconeogenesis in the liver is sensitive to blood-glucose concentration	483	Mitochondria are the result of an endosymbiotic event	525
Substrate cycles amplify metabolic signals and produce heat	485	<b>18.2 Oxidative Phosphorylation Depends on Electron Transfer</b>	<b>526</b>
Lactate and alanine formed by contracting muscle are used by other organs	485	The electron-transfer potential of an electron is measured as redox potential	526
Glycolysis and gluconeogenesis are evolutionarily intertwined	487	A 1.14-volt potential difference between NADH and molecular oxygen drives electron transport through the chain and favors the formation of a proton gradient	528
<b>CHAPTER 17 The Citric Acid Cycle</b>	<b>495</b>	<b>18.3 The Respiratory Chain Consists of Four Complexes: Three Proton Pumps and a Physical Link to the Citric Acid Cycle</b>	<b>529</b>
The citric acid cycle harvests high-energy electrons	496	Iron-sulfur clusters are common components of the electron transport chain	531
<b>17.1 The Pyruvate Dehydrogenase Complex Links Glycolysis to the Citric Acid Cycle</b>	<b>497</b>		
Mechanism: The synthesis of acetyl coenzyme A from pyruvate requires three enzymes and five coenzymes	498		



The high-potential electrons of NADH enter the respiratory chain at NADH-Q oxidoreductase	532
Ubiquinol is the entry point for electrons from FADH <sub>2</sub> of flavoproteins	533
Electrons flow from ubiquinol to cytochrome <i>c</i> through Q-cytochrome <i>c</i> oxidoreductase	533
The Q cycle funnels electrons from a two-electron carrier to a one-electron carrier and pumps protons	535
Cytochrome <i>c</i> oxidase catalyzes the reduction of molecular oxygen to water	535
Toxic derivatives of molecular oxygen such as superoxide radicals are scavenged by protective enzymes	538
Electrons can be transferred between groups that are not in contact	540
The conformation of cytochrome <i>c</i> has remained essentially constant for more than a billion years	541
<b>18.4 A Proton Gradient Powers the Synthesis of ATP</b>	<b>541</b>
ATP synthase is composed of a proton-conducting unit and a catalytic unit	543
Proton flow through ATP synthase leads to the release of tightly bound ATP: The binding-change mechanism	544
Rotational catalysis is the world's smallest molecular motor	546
Proton flow around the <i>c</i> ring powers ATP synthesis	546
ATP synthase and G proteins have several common features	548
<b>18.5 Many Shuttles Allow Movement Across Mitochondrial Membranes</b>	<b>549</b>
Electrons from cytoplasmic NADH enter mitochondria by shuttles	549
The entry of ADP into mitochondria is coupled to the exit of ATP by ATP-ADP translocase	550
Mitochondrial transporters for metabolites have a common tripartite structure	551
<b>18.6 The Regulation of Cellular Respiration Is Governed Primarily by the Need for ATP</b>	<b>552</b>
The complete oxidation of glucose yields about 30 molecules of ATP	552
The rate of oxidative phosphorylation is determined by the need for ATP	553
ATP synthase can be regulated	554
Regulated uncoupling leads to the generation of heat	554
Oxidative phosphorylation can be inhibited at many stages	556
Mitochondrial diseases are being discovered	557
Mitochondria play a key role in apoptosis	557
Power transmission by proton gradients is a central motif of bioenergetics	558
<b>CHAPTER 19 The Light Reactions of Photosynthesis</b>	<b>565</b>
Photosynthesis converts light energy into chemical energy	566
<b>19.1 Photosynthesis Takes Place in Chloroplasts</b>	<b>567</b>
The primary events of photosynthesis take place in thylakoid membranes	567
Chloroplasts arose from an endosymbiotic event	568
<b>19.2 Light Absorption by Chlorophyll Induces Electron Transfer</b>	<b>568</b>
A special pair of chlorophylls initiate charge separation	569
Cyclic electron flow reduces the cytochrome of the reaction center	572
<b>19.3 Two Photosystems Generate a Proton Gradient and NADPH in Oxygenic Photosynthesis</b>	<b>572</b>
Photosystem II transfers electrons from water to plastoquinone and generates a proton gradient	572
Cytochrome <i>bf</i> links photosystem II to photosystem I	575
Photosystem I uses light energy to generate reduced ferredoxin, a powerful reductant	575
Ferredoxin-NADP <sup>+</sup> reductase converts NADP <sup>+</sup> into NADPH	576
<b>19.4 A Proton Gradient across the Thylakoid Membrane Drives ATP Synthesis</b>	<b>578</b>
The ATP synthase of chloroplasts closely resembles those of mitochondria and prokaryotes	578
The activity of chloroplast ATP synthase is regulated	579
Cyclic electron flow through photosystem I leads to the production of ATP instead of NADPH	580
The absorption of eight photons yields one O <sub>2</sub> , two NADPH, and three ATP molecules	581
<b>19.5 Accessory Pigments Funnel Energy into Reaction Centers</b>	<b>581</b>
Resonance energy transfer allows energy to move from the site of initial absorbance to the reaction center	582
The components of photosynthesis are highly organized	583
Many herbicides inhibit the light reactions of photosynthesis	584
<b>19.6 The Ability to Convert Light into Chemical Energy Is Ancient</b>	<b>584</b>
Artificial photosynthetic systems may provide clean, renewable energy	585
<b>CHAPTER 20 The Calvin Cycle and the Pentose Phosphate Pathway</b>	<b>589</b>
<b>20.1 The Calvin Cycle Synthesizes Hexoses from Carbon Dioxide and Water</b>	<b>590</b>
Carbon dioxide reacts with ribulose 1,5-bisphosphate to form two molecules of 3-phosphoglycerate	591
Rubisco activity depends on magnesium and carbamate	592
Rubisco activase is essential for rubisco activity	593
Rubisco also catalyzes a wasteful oxygenase reaction: Catalytic imperfection	593
Hexose phosphates are made from phosphoglycerate, and ribulose 1,5-bisphosphate is regenerated	594
Three ATP and two NADPH molecules are used to bring carbon dioxide to the level of a hexose	597
Starch and sucrose are the major carbohydrate stores in plants	597
<b>20.2 The Activity of the Calvin Cycle Depends on Environmental Conditions</b>	<b>598</b>



Rubisco is activated by light-driven changes in proton and magnesium ion concentrations	598
Thioredoxin plays a key role in regulating the Calvin cycle	599
The C <sub>4</sub> pathway of tropical plants accelerates photosynthesis by concentrating carbon dioxide	599
Crassulacean acid metabolism permits growth in arid ecosystems	601

### **20.3 The Pentose Phosphate Pathway Generates NADPH and Synthesizes Five-Carbon Sugars** 601

Two molecules of NADPH are generated in the conversion of glucose 6-phosphate into ribulose 5-phosphate	602
The pentose phosphate pathway and glycolysis are linked by transketolase and transaldolase	602
Mechanism: Transketolase and transaldolase stabilize carbanionic intermediates by different mechanisms	605

### **20.4 The Metabolism of Glucose 6-Phosphate by the Pentose Phosphate Pathway Is Coordinated with Glycolysis** 607

The rate of the pentose phosphate pathway is controlled by the level of NADP <sup>+</sup>	607
The flow of glucose 6-phosphate depends on the need for NADPH, ribose 5-phosphate, and ATP	608
The pentose phosphate pathway is required for rapid cell growth	610
Through the looking-glass: The Calvin cycle and the pentose phosphate pathway are mirror images	610

### **20.5 Glucose 6-Phosphate Dehydrogenase Plays a Key Role in Protection Against Reactive Oxygen Species** 610

Glucose 6-phosphate dehydrogenase deficiency causes a drug-induced hemolytic anemia	610
A deficiency of glucose 6-phosphate dehydrogenase confers an evolutionary advantage in some circumstances	612

## **CHAPTER 21 Glycogen Metabolism** 617

Glycogen metabolism is the regulated release and storage of glucose	618
---	-----

### **21.1 Glycogen Breakdown Requires the Interplay of Several Enzymes** 619

Phosphorylase catalyzes the phosphorolytic cleavage of glycogen to release glucose 1-phosphate	619
Mechanism: Pyridoxal phosphate participates in the phosphorolytic cleavage of glycogen	620
A debranching enzyme also is needed for the breakdown of glycogen	621
Phosphoglucomutase converts glucose 1-phosphate into glucose 6-phosphate	622
The liver contains glucose 6-phosphatase, a hydrolytic enzyme absent from muscle	622

### **21.2 Phosphorylase Is Regulated by Allosteric Interactions and Reversible Phosphorylation** 623

Liver phosphorylase produces glucose for use by other tissues	623
---	-----

Muscle phosphorylase is regulated by the intracellular energy charge	625
--	-----

Biochemical characteristics of muscle fiber types differ	625
--	-----

Phosphorylation promotes the conversion of phosphorylase <i>b</i> to phosphorylase <i>a</i>	626
---	-----

Phosphorylase kinase is activated by phosphorylation and calcium ions	626
---	-----

### **21.3 Epinephrine and Glucagon Signal the Need for Glycogen Breakdown** 627

G proteins transmit the signal for the initiation of glycogen breakdown	627
---	-----

Glycogen breakdown must be rapidly turned off when necessary	629
--	-----

The regulation of glycogen phosphorylase became more sophisticated as the enzyme evolved	629
--	-----

### **21.4 Glycogen Is Synthesized and Degraded by Different Pathways** 630

UDP-glucose is an activated form of glucose	630
---	-----

Glycogen synthase catalyzes the transfer of glucose from UDP-glucose to a growing chain	630
---	-----

A branching enzyme forms $\alpha$ -1,6 linkages	631
---	-----

Glycogen synthase is the key regulatory enzyme in glycogen synthesis	632
--	-----

Glycogen is an efficient storage form of glucose	632
--	-----

### **21.5 Glycogen Breakdown and Synthesis Are Reciprocally Regulated** 632

Protein phosphatase 1 reverses the regulatory effects of kinases on glycogen metabolism	633
---	-----

Insulin stimulates glycogen synthesis by inactivating glycogen synthase kinase	635
--	-----

Glycogen metabolism in the liver regulates the blood-glucose level	635
--	-----

A biochemical understanding of glycogen-storage diseases is possible	637
--	-----

## **CHAPTER 22 Fatty Acid Metabolism** 643

Fatty acid degradation and synthesis mirror each other in their chemical reactions	644
--	-----

### **22.1 Triacylglycerols Are Highly Concentrated Energy Stores** 645

Dietary lipids are digested by pancreatic lipases	645
---	-----

Dietary lipids are transported in chylomicrons	646
--	-----

### **22.2 The Use of Fatty Acids as Fuel Requires Three Stages of Processing** 647

Triacylglycerols are hydrolyzed by hormone-stimulated lipases	647
---	-----

Free fatty acids and glycerol are released into the blood	648
---	-----

Fatty acids are linked to coenzyme A before they are oxidized	648
---	-----

Carnitine carries long-chain activated fatty acids into the mitochondrial matrix	649
--	-----

Acetyl CoA, NADH, and FADH <sub>2</sub> are generated in each round of fatty acid oxidation	650
---	-----



The complete oxidation of palmitate yields 106 molecules of ATP	652	<b>23.2 Protein Turnover Is Tightly Regulated</b>	<b>683</b>
<b>22.3 Unsaturated and Odd-Chain Fatty Acids Require Additional Steps for Degradation</b>	<b>652</b>	Ubiquitin tags proteins for destruction	683
An isomerase and a reductase are required for the oxidation of unsaturated fatty acids	652	The proteasome digests the ubiquitin-tagged proteins	685
Odd-chain fatty acids yield propionyl CoA in the final thiolysis step	654	The ubiquitin pathway and the proteasome have prokaryotic counterparts	686
Vitamin B <sub>12</sub> contains a corrin ring and a cobalt atom	654	Protein degradation can be used to regulate biological function	687
Mechanism: Methylmalonyl CoA mutase catalyzes a rearrangement to form succinyl CoA	655	<b>23.3 The First Step in Amino Acid Degradation Is the Removal of Nitrogen</b>	<b>687</b>
Fatty acids are also oxidized in peroxisomes	656	Alpha-amino groups are converted into ammonium ions by the oxidative deamination of glutamate	687
Ketone bodies are formed from acetyl CoA when fat breakdown predominates	657	Mechanism: Pyridoxal phosphate forms Schiff-base intermediates in aminotransferases	689
Ketone bodies are a major fuel in some tissues	658	Aspartate aminotransferase is an archetypal pyridoxal-dependent transaminase	690
Animals cannot convert fatty acids into glucose	660	Blood levels of aminotransferases serve a diagnostic function	691
Some fatty acids may contribute to the development of pathological conditions	661	Pyridoxal phosphate enzymes catalyze a wide array of reactions	691
<b>22.4 Fatty Acids Are Synthesized by Fatty Acid Synthase</b>	<b>661</b>	Serine and threonine can be directly deaminated	692
Fatty acids are synthesized and degraded by different pathways	661	Peripheral tissues transport nitrogen to the liver	692
The formation of malonyl CoA is the committed step in fatty acid synthesis	662	<b>23.4 Ammonium Ion Is Converted into Urea in Most Terrestrial Vertebrates</b>	<b>693</b>
Intermediates in fatty acid synthesis are attached to an acyl carrier protein	662	The urea cycle begins with the formation of carbamoyl phosphate	693
Fatty acid synthesis consists of a series of condensation, reduction, dehydration, and reduction reactions	662	Carbamoyl phosphate synthetase is the key regulatory enzyme for urea synthesis	694
Fatty acids are synthesized by a multifunctional enzyme complex in animals	664	Carbamoyl phosphate reacts with ornithine to begin the urea cycle	694
The synthesis of palmitate requires 8 molecules of acetyl CoA, 14 molecules of NADPH, and 7 molecules of ATP	666	The urea cycle is linked to gluconeogenesis	696
Citrate carries acetyl groups from mitochondria to the cytoplasm for fatty acid synthesis	666	Urea-cycle enzymes are evolutionarily related to enzymes in other metabolic pathways	696
Several sources supply NADPH for fatty acid synthesis	667	Inherited defects of the urea cycle cause hyperammonemia and can lead to brain damage	697
Fatty acid metabolism is altered in tumor cells	667	Urea is not the only means of disposing of excess nitrogen	698
<b>22.5 The Elongation and Unsaturation of Fatty Acids are Accomplished by Accessory Enzyme Systems</b>	<b>668</b>	<b>23.5 Carbon Atoms of Degraded Amino Acids Emerge as Major Metabolic Intermediates</b>	<b>698</b>
Membrane-bound enzymes generate unsaturated fatty acids	668	Pyruvate is an entry point into metabolism for a number of amino acids	699
Eicosanoid hormones are derived from polyunsaturated fatty acids	669	Oxaloacetate is an entry point into metabolism for aspartate and asparagine	700
Variations on a theme: Polyketide and nonribosomal peptide synthetases resemble fatty acid synthase	670	Alpha-ketoglutarate is an entry point into metabolism for five-carbon amino acids	700
<b>22.6 Acetyl CoA Carboxylase Plays a Key Role in Controlling Fatty Acid Metabolism</b>	<b>670</b>	Succinyl coenzyme A is a point of entry for several nonpolar amino acids	701
Acetyl CoA carboxylase is regulated by conditions in the cell	671	Methionine degradation requires the formation of a key methyl donor, S-adenosylmethionine	701
Acetyl CoA carboxylase is regulated by a variety of hormones	671	The branched-chain amino acids yield acetyl CoA, acetoacetate, or propionyl CoA	701
<b>CHAPTER 23 Protein Turnover and Amino Acid Catabolism</b>	<b>681</b>	Oxygenases are required for the degradation of aromatic amino acids	703
<b>23.1 Proteins are Degraded to Amino Acids</b>	<b>682</b>	<b>23.6 Inborn Errors of Metabolism Can Disrupt Amino Acid Degradation</b>	<b>705</b>
The digestion of dietary proteins begins in the stomach and is completed in the intestine	682	Phenylketonuria is one of the most common metabolic disorders	706
Cellular proteins are degraded at different rates	682	Determining the basis of the neurological symptoms of phenylketonuria is an active area of research	706



## Part III SYNTHESIZING THE MOLECULES OF LIFE

### CHAPTER 24 The Biosynthesis of Amino Acids 713

Amino acid synthesis requires solutions to three key biochemical problems 714

#### 24.1 Nitrogen Fixation: Microorganisms Use ATP and a Powerful Reductant to Reduce Atmospheric Nitrogen to Ammonia 714

The iron–molybdenum cofactor of nitrogenase binds and reduces atmospheric nitrogen 715

Ammonium ion is assimilated into an amino acid through glutamate and glutamine 717

#### 24.2 Amino Acids Are Made from Intermediates of the Citric Acid Cycle and Other Major Pathways 719

Human beings can synthesize some amino acids but must obtain others from their diet 719

Aspartate, alanine, and glutamate are formed by the addition of an amino group to an alpha-ketoacid 720

A common step determines the chirality of all amino acids 721

The formation of asparagine from aspartate requires an adenylated intermediate 721

Glutamate is the precursor of glutamine, proline, and arginine 722

3-Phosphoglycerate is the precursor of serine, cysteine, and glycine 722

Tetrahydrofolate carries activated one-carbon units at several oxidation levels 723

S-Adenosylmethionine is the major donor of methyl groups 724

Cysteine is synthesized from serine and homocysteine 726

High homocysteine levels correlate with vascular disease 726

Shikimate and chorismate are intermediates in the biosynthesis of aromatic amino acids 727

Tryptophan synthase illustrates substrate channeling in enzymatic catalysis 729

#### 24.3 Feedback Inhibition Regulates Amino Acid Biosynthesis 730

Branched pathways require sophisticated regulation 731

The sensitivity of glutamine synthetase to allosteric regulation is altered by covalent modification 732

#### 24.4 Amino Acids Are Precursors of Many Biomolecules 734

Glutathione, a gamma-glutamyl peptide, serves as a sulfhydryl buffer and an antioxidant 734

Nitric oxide, a short-lived signal molecule, is formed from arginine 735

Porphyrins are synthesized from glycine and succinyl coenzyme A 736

Porphyrins accumulate in some inherited disorders of porphyrin metabolism 737

### CHAPTER 25 Nucleotide Biosynthesis 743

Nucleotides can be synthesized by de novo or salvage pathways 744

#### 25.1 The Pyrimidine Ring Is Assembled de Novo or Recovered by Salvage Pathways 744

Bicarbonate and other oxygenated carbon compounds are activated by phosphorylation 745

The side chain of glutamine can be hydrolyzed to generate ammonia 745

Intermediates can move between active sites by channeling 745

Orotate acquires a ribose ring from PRPP to form a pyrimidine nucleotide and is converted into uridylylate 746

Nucleotide mono-, di-, and triphosphates are interconvertible 747

CTP is formed by amination of UTP 747

Salvage pathways recycle pyrimidine bases 748

#### 25.2 Purine Bases Can Be Synthesized de Novo or Recycled by Salvage Pathways 748

The purine ring system is assembled on ribose phosphate 749

The purine ring is assembled by successive steps of activation by phosphorylation followed by displacement 749

AMP and GMP are formed from IMP 751

Enzymes of the purine synthesis pathway associate with one another in vivo 752

Salvage pathways economize intracellular energy expenditure 752

#### 25.3 Deoxyribonucleotides Are Synthesized by the Reduction of Ribonucleotides Through a Radical Mechanism 753

Mechanism: A tyrosyl radical is critical to the action of ribonucleotide reductase 753

Stable radicals other than tyrosyl radical are employed by other ribonucleotide reductases 755

Thymidylate is formed by the methylation of deoxyuridylylate 755

Dihydrofolate reductase catalyzes the regeneration of tetrahydrofolate, a one-carbon carrier 756

Several valuable anticancer drugs block the synthesis of thymidylate 757

#### 25.4 Key Steps in Nucleotide Biosynthesis Are Regulated by Feedback Inhibition 758

Pyrimidine biosynthesis is regulated by aspartate transcarbamoylase 758

The synthesis of purine nucleotides is controlled by feedback inhibition at several sites 758

The synthesis of deoxyribonucleotides is controlled by the regulation of ribonucleotide reductase 759

#### 25.5 Disruptions in Nucleotide Metabolism Can Cause Pathological Conditions 760

The loss of adenosine deaminase activity results in severe combined immunodeficiency 760

Gout is induced by high serum levels of urate 761

Lesch–Nyhan syndrome is a dramatic consequence of mutations in a salvage-pathway enzyme 761

Folic acid deficiency promotes birth defects such as spina bifida 762



## CHAPTER 26 The Biosynthesis of Membrane Lipids and Steroids 767

### 26.1 Phosphatidate Is a Common Intermediate in the Synthesis of Phospholipids and Triacylglycerols 768

- The synthesis of phospholipids requires an activated intermediate 769
- Some phospholipids are synthesized from an activated alcohol 770
- Phosphatidylcholine is an abundant phospholipid 770
- Excess choline is implicated in the development of heart disease 771
- Base-exchange reactions can generate phospholipids 771
- Sphingolipids are synthesized from ceramide 772
- Gangliosides are carbohydrate-rich sphingolipids that contain acidic sugars 772
- Sphingolipids confer diversity on lipid structure and function 773
- Respiratory distress syndrome and Tay–Sachs disease result from the disruption of lipid metabolism 774
- Ceramide metabolism stimulates tumor growth 774
- Phosphatidic acid phosphatase is a key regulatory enzyme in lipid metabolism 775

### 26.2 Cholesterol Is Synthesized from Acetyl Coenzyme A in Three Stages 776

- The synthesis of mevalonate, which is activated as isopentenyl pyrophosphate, initiates the synthesis of cholesterol 776
- Squalene (C<sub>30</sub>) is synthesized from six molecules of isopentenyl pyrophosphate (C<sub>5</sub>) 777
- Squalene cyclizes to form cholesterol 778

### 26.3 The Complex Regulation of Cholesterol Biosynthesis Takes Place at Several Levels 779

- Lipoproteins transport cholesterol and triacylglycerols throughout the organism 782
- Low-density lipoproteins play a central role in cholesterol metabolism 784
- The absence of the LDL receptor leads to hypercholesterolemia and atherosclerosis 784
- Mutations in the LDL receptor prevent LDL release and result in receptor destruction 785
- Cycling of the LDL receptor is regulated 787
- HDL appears to protect against atherosclerosis 787
- The clinical management of cholesterol levels can be understood at a biochemical level 788

### 26.4 Important Derivatives of Cholesterol Include Bile Salts and Steroid Hormones 788

- Letters identify the steroid rings and numbers identify the carbon atoms 790
- Steroids are hydroxylated by cytochrome P450 monooxygenases that use NADPH and O<sub>2</sub> 790
- The cytochrome P450 system is widespread and performs a protective function 791
- Pregnenolone, a precursor of many other steroids, is formed from cholesterol by cleavage of its side chain 792

- Pregesterone and corticosteroids are synthesized from pregnenolone 792
- Androgens and estrogens are synthesized from pregnenolone 792
- Vitamin D is derived from cholesterol by the ring-splitting activity of light 794

## CHAPTER 27 The Integration of Metabolism 801

### 27.1 Caloric Homeostasis Is a Means of Regulating Body Weight 802

### 27.2 The Brain Plays a Key Role in Caloric Homeostasis 804

- Signals from the gastrointestinal tract induce feelings of satiety 804
- Leptin and insulin regulate long-term control over caloric homeostasis 805
- Leptin is one of several hormones secreted by adipose tissue 806
- Leptin resistance may be a contributing factor to obesity 806
- Dieting is used to combat obesity 807

### 27.3 Diabetes Is a Common Metabolic Disease Often Resulting from Obesity 807

- Insulin initiates a complex signal-transduction pathway in muscle 808
- Metabolic syndrome often precedes type 2 diabetes 809
- Excess fatty acids in muscle modify metabolism 810
- Insulin resistance in muscle facilitates pancreatic failure 810
- Metabolic derangements in type 1 diabetes result from insulin insufficiency and glucagon excess 812

### 27.4 Exercise Beneficially Alters the Biochemistry of Cells 813

- Mitochondrial biogenesis is stimulated by muscular activity 813
- Fuel choice during exercise is determined by the intensity and duration of activity 813

### 27.5 Food Intake and Starvation Induce Metabolic Changes 816

- The starved–fed cycle is the physiological response to a fast 816
- Metabolic adaptations in prolonged starvation minimize protein degradation 818

### 27.6 Ethanol Alters Energy Metabolism in the Liver 819

- Ethanol metabolism leads to an excess of NADH 820
- Excess ethanol consumption disrupts vitamin metabolism 821

## CHAPTER 28 DNA Replication, Repair, and Recombination 827

### 28.1 DNA Replication Proceeds by the Polymerization of Deoxyribonucleoside Triphosphates Along a Template 828



DNA polymerases require a template and a primer	829
All DNA polymerases have structural features in common	829
Two bound metal ions participate in the polymerase reaction	829
The specificity of replication is dictated by complementarity of shape between bases	830
An RNA primer synthesized by primase enables DNA synthesis to begin	831
One strand of DNA is made continuously, whereas the other strand is synthesized in fragments	831
DNA ligase joins ends of DNA in duplex regions	832
The separation of DNA strands requires specific helicases and ATP hydrolysis	832
<b>28.2 DNA Unwinding and Supercoiling Are Controlled by Topoisomerases</b>	<b>833</b>
The linking number of DNA, a topological property, determines the degree of supercoiling	835
Topoisomerases prepare the double helix for unwinding	836
Type I topoisomerases relax supercoiled structures	836
Type II topoisomerases can introduce negative supercoils through coupling to ATP hydrolysis	837
<b>28.3 DNA Replication Is Highly Coordinated</b>	<b>839</b>
DNA replication requires highly processive polymerases	839
The leading and lagging strands are synthesized in a coordinated fashion	840
DNA replication in <i>Escherichia coli</i> begins at a unique site	842
DNA synthesis in eukaryotes is initiated at multiple sites	843
Telomeres are unique structures at the ends of linear chromosomes	844
Telomeres are replicated by telomerase, a specialized polymerase that carries its own RNA template	845
<b>28.4 Many Types of DNA Damage Can Be Repaired</b>	<b>845</b>
Errors can arise in DNA replication	846
Bases can be damaged by oxidizing agents, alkylating agents, and light	846
DNA damage can be detected and repaired by a variety of systems	847
The presence of thymine instead of uracil in DNA permits the repair of deaminated cytosine	849
Some genetic diseases are caused by the expansion of repeats of three nucleotides	850
Many cancers are caused by the defective repair of DNA	850
Many potential carcinogens can be detected by their mutagenic action on bacteria	852
<b>28.5 DNA Recombination Plays Important Roles in Replication, Repair, and Other Processes</b>	<b>852</b>
RecA can initiate recombination by promoting strand invasion	853
Some recombination reactions proceed through Holliday-junction intermediates	854

<b>CHAPTER 29 RNA Synthesis and Processing</b>	<b>859</b>
RNA synthesis comprises three stages: Initiation, elongation, and termination	860
<b>29.1 RNA Polymerases Catalyze Transcription</b>	<b>861</b>
RNA chains are formed de novo and grow in the 5'-to-3' direction	862
RNA polymerases backtrack and correct errors	863
RNA polymerase binds to promoter sites on the DNA template to initiate transcription	864
Sigma subunits of RNA polymerase recognize promoter sites	865
RNA polymerases must unwind the template double helix for transcription to take place	865
Elongation takes place at transcription bubbles that move along the DNA template	866
Sequences within the newly transcribed RNA signal termination	866
Some messenger RNAs directly sense metabolite concentrations	867
The <i>rho</i> protein helps to terminate the transcription of some genes	868
Some antibiotics inhibit transcription	869
Precursors of transfer and ribosomal RNA are cleaved and chemically modified after transcription in prokaryotes	870
<b>29.2 Transcription in Eukaryotes Is Highly Regulated</b>	<b>871</b>
Three types of RNA polymerase synthesize RNA in eukaryotic cells	872
Three common elements can be found in the RNA polymerase II promoter region	874
The TFIID protein complex initiates the assembly of the active transcription complex	874
Multiple transcription factors interact with eukaryotic promoters	875
Enhancer sequences can stimulate transcription at start sites thousands of bases away	876
<b>29.3 The Transcription Products of Eukaryotic Polymerases Are Processed</b>	<b>876</b>
RNA polymerase I produces three ribosomal RNAs	877
RNA polymerase III produces transfer RNA	877
The product of RNA polymerase II, the pre-mRNA transcript, acquires a 5' cap and a 3' poly(A) tail	878
Small regulatory RNAs are cleaved from larger precursors	879
RNA editing changes the proteins encoded by mRNA	879
Sequences at the ends of introns specify splice sites in mRNA precursors	880
Splicing consists of two sequential transesterification reactions	881
Small nuclear RNAs in spliceosomes catalyze the splicing of mRNA precursors	882
Transcription and processing of mRNA are coupled	883
Mutations that affect pre-mRNA splicing cause disease	884
Most human pre-mRNAs can be spliced in alternative ways to yield different proteins	885
<b>29.4 The Discovery of Catalytic RNA was Revealing in Regard to Both Mechanism and Evolution</b>	<b>886</b>



**CHAPTER 30 Protein Synthesis 893****30.1 Protein Synthesis Requires the Translation of Nucleotide Sequences into Amino Acid Sequences 894**

- The synthesis of long proteins requires a low error frequency 894
- Transfer RNA molecules have a common design 895
- Some transfer RNA molecules recognize more than one codon because of wobble in base-pairing 897

**30.2 Aminoacyl Transfer RNA Synthetases Read the Genetic Code 898**

- Amino acids are first activated by adenylation 898
- Aminoacyl-tRNA synthetases have highly discriminating amino acid activation sites 899
- Proofreading by aminoacyl-tRNA synthetases increases the fidelity of protein synthesis 900
- Synthetases recognize various features of transfer RNA molecules 901
- Aminoacyl-tRNA synthetases can be divided into two classes 901

**30.3 The Ribosome Is the Site of Protein Synthesis 902**

- Ribosomal RNAs (5S, 16S, and 23S rRNA) play a central role in protein synthesis 903
- Ribosomes have three tRNA-binding sites that bridge the 30s and 50s subunits 905
- The start signal is usually AUG preceded by several bases that pair with 16S rRNA 905
- Bacterial protein synthesis is initiated by formylmethionyl transfer RNA 906
- Formylmethionyl-tRNA<sub>f</sub> is placed in the P site of the ribosome in the formation of the 70S initiation complex 907
- Elongation factors deliver aminoacyl-tRNA to the ribosome 907
- Peptidyl transferase catalyzes peptide-bond synthesis 908
- The formation of a peptide bond is followed by the GTP-driven translocation of tRNAs and mRNA 909
- Protein synthesis is terminated by release factors that read stop codons 910

**30.4 Eukaryotic Protein Synthesis Differs from Bacterial Protein Synthesis Primarily in Translation Initiation 911**

- Mutations in initiation factor 2 cause a curious pathological condition 913

**30.5 A Variety of Antibiotics and Toxins Can Inhibit Protein Synthesis 913**

- Some antibiotics inhibit protein synthesis 914
- Diphtheria toxin blocks protein synthesis in eukaryotes by inhibiting translocation 914
- Ricin fatally modifies 28S ribosomal RNA 915

**30.6 Ribosomes Bound to the Endoplasmic Reticulum Manufacture Secretory and Membrane Proteins 915**

- Protein synthesis begins on ribosomes that are free in the cytoplasm 916
- Signal sequences mark proteins for translocation across the endoplasmic reticulum membrane 916

Transport vesicles carry cargo proteins to their final destination 918

**CHAPTER 31 The Control of Gene Expression in Prokaryotes 925****31.1 Many DNA-Binding Proteins Recognize Specific DNA Sequences 926**

The helix-turn-helix motif is common to many prokaryotic DNA-binding proteins 927

**31.2 Prokaryotic DNA-Binding Proteins Bind Specifically to Regulatory Sites in Operons 927**

- An operon consists of regulatory elements and protein-encoding genes 928
- The *lac* repressor protein in the absence of lactose binds to the operator and blocks transcription 929
- Ligand binding can induce structural changes in regulatory proteins 930
- The operon is a common regulatory unit in prokaryotes 930
- Transcription can be stimulated by proteins that contact RNA polymerase 931

**31.3 Regulatory Circuits Can Result in Switching Between Patterns of Gene Expression 932**

- The  $\lambda$  repressor regulates its own expression 932
- A circuit based on the  $\lambda$  repressor and Cro forms a genetic switch 933
- Many prokaryotic cells release chemical signals that regulate gene expression in other cells 933
- Biofilms are complex communities of prokaryotes 934

**31.4 Gene Expression Can Be Controlled at Posttranscriptional Levels 935**

- Attenuation is a prokaryotic mechanism for regulating transcription through the modulation of nascent RNA secondary structure 935

**CHAPTER 32 The Control of Gene Expression in Eukaryotes 941****32.1 Eukaryotic DNA Is Organized into Chromatin 943**

- Nucleosomes are complexes of DNA and histones 943
- DNA wraps around histone octamers to form nucleosomes 943

**32.2 Transcription Factors Bind DNA and Regulate Transcription Initiation 945**

- A range of DNA-binding structures are employed by eukaryotic DNA-binding proteins 945
- Activation domains interact with other proteins 946
- Multiple transcription factors interact with eukaryotic regulatory regions 946
- Enhancers can stimulate transcription in specific cell types 946
- Induced pluripotent stem cells can be generated by introducing four transcription factors into differentiated cells 947

**32.3 The Control of Gene Expression Can Require Chromatin Remodeling 948**



The methylation of DNA can alter patterns of gene expression	949	Mechanosensory channels have been identified in <i>Drosophila</i> and vertebrates	976
Steroids and related hydrophobic molecules pass through membranes and bind to DNA-binding receptors	949	<b>33.5 Touch Includes the Sensing of Pressure, Temperature, and Other Factors</b>	<b>977</b>
Nuclear hormone receptors regulate transcription by recruiting coactivators to the transcription complex	950	Studies of capsaicin reveal a receptor for sensing high temperatures and other painful stimuli	977
Steroid-hormone receptors are targets for drugs	951	<b>CHAPTER 34 The Immune System</b>	<b>981</b>
Chromatin structure is modulated through covalent modifications of histone tails	952	Innate immunity is an evolutionarily ancient defense system	982
Histone deacetylases contribute to transcriptional repression	953	The adaptive immune system responds by using the principles of evolution	984
<b>32.4 Eukaryotic Gene Expression Can Be Controlled at Posttranscriptional Levels</b>	<b>954</b>	<b>34.1 Antibodies Possess Distinct Antigen-Binding and Effector Units</b>	<b>985</b>
Genes associated with iron metabolism are translationally regulated in animals	954	<b>34.2 Antibodies Bind Specific Molecules Through Hypervariable Loops</b>	<b>988</b>
Small RNAs regulate the expression of many eukaryotic genes	956	The immunoglobulin fold consists of a beta-sandwich framework with hypervariable loops	988
<b>Part IV RESPONDING TO ENVIRONMENTAL CHANGES</b>		X-ray analyses have revealed how antibodies bind antigens	989
<b>CHAPTER 33 Sensory Systems</b>	<b>961</b>	Large antigens bind antibodies with numerous interactions	990
<b>33.1 A Wide Variety of Organic Compounds Are Detected by Olfaction</b>	<b>962</b>	<b>34.3 Diversity Is Generated by Gene Rearrangements</b>	<b>991</b>
Olfaction is mediated by an enormous family of seven-transmembrane-helix receptors	962	J (joining) genes and D (diversity) genes increase antibody diversity	991
Odorants are decoded by a combinatorial mechanism	964	More than $10^8$ antibodies can be formed by combinatorial association and somatic mutation	992
<b>33.2 Taste Is a Combination of Senses That Function by Different Mechanisms</b>	<b>966</b>	The oligomerization of antibodies expressed on the surfaces of immature B cells triggers antibody secretion	993
Sequencing of the human genome led to the discovery of a large family of 7TM bitter receptors	967	Different classes of antibodies are formed by the hopping of $V_H$ genes	994
A heterodimeric 7TM receptor responds to sweet compounds	968	<b>34.4 Major-Histocompatibility-Complex Proteins Present Peptide Antigens on Cell Surfaces for Recognition by T-Cell Receptors</b>	<b>995</b>
Umami, the taste of glutamate and aspartate, is mediated by a heterodimeric receptor related to the sweet receptor	969	Peptides presented by MHC proteins occupy a deep groove flanked by alpha helices	996
Salty tastes are detected primarily by the passage of sodium ions through channels	969	T-cell receptors are antibody-like proteins containing variable and constant regions	998
Sour tastes arise from the effects of hydrogen ions (acids) on channels	969	CD8 on cytotoxic T cells acts in concert with T-cell receptors	998
<b>33.3 Photoreceptor Molecules in the Eye Detect Visible Light</b>	<b>970</b>	Helper T cells stimulate cells that display foreign peptides bound to class II MHC proteins	1000
Rhodopsin, a specialized 7TM receptor, absorbs visible light	970	Helper T cells rely on the T-cell receptor and CD4 to recognize foreign peptides on antigen-presenting cells	1000
Light absorption induces a specific isomerization of bound 11- <i>cis</i> -retinal	971	MHC proteins are highly diverse	1002
Light-induced lowering of the calcium level coordinates recovery	972	Human immunodeficiency viruses subvert the immune system by destroying helper T cells	1003
Color vision is mediated by three cone receptors that are homologs of rhodopsin	973	<b>34.5 The Immune System Contributes to the Prevention and the Development of Human Diseases</b>	<b>1004</b>
Rearrangements in the genes for the green and red pigments lead to "color blindness"	974	T cells are subjected to positive and negative selection in the thymus	1004
<b>33.4 Hearing Depends on the Speedy Detection of Mechanical Stimuli</b>	<b>975</b>	Autoimmune diseases result from the generation of immune responses against self-antigens	1005
Hair cells use a connected bundle of stereocilia to detect tiny motions	975		



The immune system plays a role in cancer prevention	1005	Drugs must have suitable properties to reach their targets	1036
Vaccines are a powerful means to prevent and eradicate disease	1006	Toxicity can limit drug effectiveness	1040
<b>CHAPTER 35 Molecular Motors</b>	<b>1011</b>	<b>36.2 Drug Candidates Can Be Discovered by Serendipity, Screening, or Design</b>	<b>1041</b>
<b>35.1 Most Molecular-Motor Proteins Are Members of the P-Loop NTPase Superfamily</b>	<b>1012</b>	Serendipitous observations can drive drug development	1041
Molecular motors are generally oligomeric proteins with an ATPase core and an extended structure	1012	Natural products are a valuable source of drugs and drug leads	1043
ATP binding and hydrolysis induce changes in the conformation and binding affinity of motor proteins	1014	Screening libraries of synthetic compounds expands the opportunity for identification of drug leads	1044
<b>35.2 Myosins Move Along Actin Filaments</b>	<b>1016</b>	Drugs can be designed on the basis of three-dimensional structural information about their targets	1046
Actin is a polar, self-assembling, dynamic polymer	1016	<b>36.3 Analyses of Genomes Hold Great Promise for Drug Discovery</b>	<b>1048</b>
Myosin head domains bind to actin filaments	1018	Potential targets can be identified in the human proteome	1048
Motions of single motor proteins can be directly observed	1018	Animal models can be developed to test the validity of potential drug targets	1049
Phosphate release triggers the myosin power stroke	1019	Potential targets can be identified in the genomes of pathogens	1050
Muscle is a complex of myosin and actin	1019	Genetic differences influence individual responses to drugs	1050
The length of the lever arm determines motor velocity	1022	<b>36.4 The Clinical Development of Drugs Proceeds Through Several Phases</b>	<b>1051</b>
<b>35.3 Kinesin and Dynein Move Along Microtubules</b>	<b>1022</b>	Clinical trials are time consuming and expensive	1052
Microtubules are hollow cylindrical polymers	1022	The evolution of drug resistance can limit the utility of drugs for infectious agents and cancer	1053
Kinesin motion is highly processive	1024	<b>Answers to Problems</b>	<b>A1</b>
<b>35.4 A Rotary Motor Drives Bacterial Motion</b>	<b>1026</b>	<b>Selected Readings</b>	<b>B1</b>
Bacteria swim by rotating their flagella	1026	<b>Index</b>	<b>C1</b>
Proton flow drives bacterial flagellar rotation	1026		
Bacterial chemotaxis depends on reversal of the direction of flagellar rotation	1028		
<b>CHAPTER 36 Drug Development</b>	<b>1033</b>		
<b>36.1 The Development of Drugs Presents Huge Challenges</b>	<b>1034</b>		
Drug candidates must be potent and selective modulators of their targets	1035		