

Contents

| | | |
|----------|---|-------------|
| | Editorial | <i>xi</i> |
| | Acknowledgment | <i>xiii</i> |
| 1 | Introduction to Single-Molecule Magnets | 1 |
| | <i>Małgorzata Holyńska</i> | |
| 1.1 | What Is a Single-Molecule Magnet? | 2 |
| 1.1.1 | Single-Chain Magnets (SCMs) | 2 |
| 1.1.2 | Single-Ion Magnets (SIMs) | 4 |
| 1.1.3 | Single-Toroid Magnets (STMs) | 5 |
| 1.2 | Historical Aspects | 6 |
| 1.3 | Recent Progress | 12 |
| 1.3.1 | SMMs Based on Actinides | 12 |
| 1.3.2 | Organometallic SMMs | 15 |
| 1.3.3 | Rational Design of SMMs | 17 |
| 1.3.4 | Quantum Computing | 19 |
| 1.3.5 | SMMs in Molecular Machines | 20 |
| 1.3.6 | Magnetic Refrigerants | 23 |
| 1.3.7 | Applications in Other Disciplines | 25 |
| | Acknowledgment | 28 |
| | References | 28 |
| 2 | Unique Magnetic Properties | 41 |
| | <i>Michael Pissas, Vassilis Psycharis, Catherine Raptopoulou, and Yiannis Sanakis</i> | |
| 2.1 | Introduction | 42 |
| 2.2 | Basic Electromagnetic Definitions | 43 |
| 2.3 | Magnetostatic Energy (Magnetic Work) | 45 |
| 2.4 | Thermodynamic Relations | 48 |
| 2.5 | Definition of ac Magnetic Susceptibility | 49 |
| 2.6 | Representative Results | 63 |
| 2.6.1 | ac Susceptibility Measurements in Tris(Acetylacetonato)iron(III), (Fe(acac) ₃) | 63 |
| 2.6.2 | ac Susceptibility Measurements in a One-Dimensional Chain Based on Mn ₆ Units | 64 |
| 2.6.3 | Spin Relaxation in a Ferromagnetically Coupled Triangular Cu ₃ Complex | 69 |

| | | |
|----------|---|-----------|
| 2.7 | Technical Aspects of the ac Susceptibility Measurements | 69 |
| 2.8 | Intermolecular Interactions | 73 |
| 2.9 | Conclusions | 78 |
| | References | 78 |
| 3 | Magnetic Modeling of Single-molecule Magnets | 87 |
| | <i>Vassilis Tangoulis and Nikolia Lalioti</i> | |
| 3.1 | Introduction | 87 |
| 3.2 | Atoms in a Magnetic Field | 90 |
| 3.2.1 | Free Atoms in a Magnetic Field | 90 |
| 3.2.1.1 | Lande g -Factor | 91 |
| 3.2.2 | Brillouin Theory | 93 |
| 3.2.2.1 | $J = 1/2$ Quantum Moment | 93 |
| 3.2.2.2 | General Quantum Case | 93 |
| 3.2.3 | Energy Spectrum | 94 |
| 3.2.3.1 | One Electron Case | 94 |
| 3.2.3.2 | Many Electrons Case | 95 |
| 3.2.3.3 | Pauli Principle – The Two Electrons Case | 95 |
| 3.2.3.4 | (L, S) -Multiplets – The Two Electrons Case | 96 |
| 3.2.3.5 | (L, S) -Coupling | 96 |
| 3.2.4 | Crystal Fields | 99 |
| 3.2.5 | Single-ion Anisotropy | 100 |
| 3.2.5.1 | Heavy Rare-Earths Case | 100 |
| 3.2.5.2 | Expressions of H_{cf} | 101 |
| 3.2.5.3 | Kramer's Theorem | 102 |
| 3.3 | Magnetic Modeling Tools | 103 |
| 3.3.1 | PHI v.3.0: Software for the Analysis of Anisotropic Monomeric and Exchange-coupled Polynuclear d- and f-Block Complexes | 103 |
| 3.3.1.1 | General Theory | 103 |
| 3.3.1.2 | Hamiltonian Formalism for the Exchange Coupling | 105 |
| 3.3.1.3 | Calculation of Thermodynamic Properties | 106 |
| 3.3.1.4 | Irreducible Tensor Operators (ITOs) Method | 107 |
| 3.3.1.5 | The Isotropic Case: Stepladder Manganese(III) Inverse-[9-MC-3]-metallacrown | 108 |
| 3.3.1.6 | Anisotropic Exchange Coupling | 110 |
| 3.3.2 | Monte Carlo Simulations: The ALPS Project Release v.2.0: Open Source Software for Strongly Correlated Systems | 115 |
| 3.3.2.1 | Stochastic Series Expansion (SSE) Quantum Monte Carlo Algorithm | 115 |
| 3.3.2.2 | Local vs Nonlocal Updates | 117 |
| 3.3.2.3 | Thermalization Process | 118 |
| 3.3.2.4 | ALPS Project: Definition of Input Files | 118 |
| 3.3.2.5 | The Case of Heterometallic Molecular Wheels | 120 |
| 3.3.2.6 | The Case of High-nuclearity Copper Cages: Tricorne Cu_{21} and Saddle-like Cyclic Cu_{16} | 123 |
| 3.3.2.7 | Other Examples. The Case of a $\text{Mn}^{\text{III}}_6\text{Mn}^{\text{II}}_6$ Molecular Wheel | 127 |
| | References | 131 |

- 4 Insight into Magnetic and Electronic Properties Through HFEPR Studies 135**
J. Krzystek and Joshua Telsler
- 4.1 Introduction: Magnetic vs Electronic Properties of Transition Metal Ions Including SMMs and SIMs 136
- 4.2 Basics of HFEPR as Applied to SIMs and Other Transition Metal Complexes 138
- 4.2.1 Spin Hamiltonian 138
- 4.2.2 Methodology of Extracting ZFS and *g* Information from HFEPR Spectra 140
- 4.3 Applicability of HFEPR to Investigating SMMs and SIMs 143
- 4.3.1 Polynuclear Clusters 143
- 4.3.2 Dimers 143
- 4.3.3 Mononuclear Complexes 144
- 4.3.4 Limitations to HFEPR 146
- 4.3.5 Techniques Alternative to HFEPR 146
- 4.4 Interplay Between Spin Hamiltonian Parameters and Crystal/Ligand-Field Parameters. From Simple Ligand Field to Sophisticated Quantum Chemical Calculations 147
- 4.4.1 Recapitulation 167
- Appendix: National High Magnetic Field Laboratory 169
- Acknowledgment 169
- References 169
- 5 Other Techniques to Study Single-Molecule Magnets 173**
Yiannis Sanakis, Vassilis Psycharis, Michael Pissas, and Catherine Raptopoulou
- 5.1 Introduction 174
- 5.2 The Mössbauer Effect 174
- 5.3 The Basic Principles of Mössbauer Spectroscopy 175
- 5.4 Hyperfine Interactions 176
- 5.4.1 The Isomer Shift 176
- 5.4.2 Quadrupole Splitting 178
- 5.4.3 Magnetic Hyperfine Interactions 179
- 5.4.4 General Remarks 181
- 5.5 Relaxation Phenomena and Dynamics 181
- 5.5.1 Mixed-Valence Systems 183
- 5.6 Application of Mössbauer Spectroscopy to Single-Molecule Magnets 185
- 5.6.1 $[\text{Fe}^{\text{III}}_8\text{O}_2(\text{OH})_{12}(\text{tacn})_6]\text{Br}_8 \cdot 9\text{H}_2\text{O}$ 185
- 5.6.2 $(\text{pyrH})_5[\text{Fe}^{\text{III}}_{13}\text{O}_4\text{F}_{24}(\text{OMe})_{12}] \cdot 4\text{H}_2\text{O} \cdot \text{MeOH}$ 187
- 5.6.3 $[\text{Fe}^{\text{III}}_{11}\text{O}_7(\text{dea})_3(\text{piv})_{12}]\text{Cl} \cdot 5\text{MeCN}$ 189
- 5.6.4 $[\text{HFe}^{\text{III}}_{19}\text{O}_{14}(\text{OEt})_{30}]$ 190
- 5.6.5 $[\text{Fe}^{\text{III}}_4(\text{OMe})_6(\text{dpm})_6]$ 191
- 5.6.6 $\{\text{Fe}^{\text{III}}[\text{Fe}^{\text{III}}(\text{L}^1)_2]_3\}$ 194
- 5.6.7 $[\text{Fe}^{\text{II}}_2(\text{acpyptentO})(\text{NCO})_3]$ 195
- 5.6.8 $[\text{Fe}^{\text{II}}_9(\text{X})_2(\text{O}_2\text{CMe})_8\{(2\text{-py})_2\text{CO}_2\}_4]$ ($\text{X} = \text{N}_3^-, \text{NCO}^-, \text{OH}^-$) 196
- 5.6.9 $[\text{Fe}^{\text{II}}_7(\text{OMe})_6(\text{Hbmsae})_6]\text{Cl}_2 \cdot 6\text{H}_2\text{O}$ 197

- 5.6.10 $[\text{Fe}^{\text{II}}\text{Fe}^{\text{III}}(\text{L})(\text{O}_2\text{CMe})_2](\text{ClO}_4)$ 199
- 5.6.11 $[(\text{Me}_3\text{TPyA})_2\text{Fe}^{\text{II}}_2(\text{L})](\text{BAR}^{\text{F}}_4)_2$ and $[(\text{Me}_3\text{TPyA})_2\text{Fe}^{\text{II/III}}_2(\text{L})](\text{BAR}^{\text{F}}_4)_3 \cdot \text{CH}_2\text{Cl}_2$ 201
- 5.6.12 $[(18\text{-C-}6)\text{K}(\text{thf})_2][(\text{t}^{\text{bs}}\text{L})\text{Fe}_3]$ and $[(\text{crypt-}222)\text{K}][(\text{t}^{\text{bs}}\text{L})\text{Fe}_3]$ 202
- 5.7 Application of Mössbauer Spectroscopy to Single-Ion Magnets 204
- 5.7.1 $[\text{M}(\text{solvent})_n][(\text{tpa}^{\text{R}})\text{Fe}^{\text{II}}]$ 204
- 5.7.2 $[\text{K}(\text{crypr-}222)][\text{Fe}^{\text{I}}\{\text{C}(\text{SiMe}_3)_3\}_2]$ 207
- 5.7.3 $[\text{Fe}^{\text{II}}\{\text{C}(\text{SiMe}_3)_3\}_2]$ 208
- 5.7.4 $[\text{Fe}^{\text{II}}\{\text{N}(\text{SiMe}_3)(\text{Dipp})\}_2]$ 209
- 5.7.5 $[\text{Fe}^{\text{II}}\{\text{OC}_6\text{H}_3\text{-}2,6\text{-}(\text{C}_6\text{H}_3\text{-}i\text{Pr}_2)_2\}_2]$ 210
- 5.7.6 $[\text{Fe}^{\text{I}}(\text{cAAC})_2\text{Cl}]$ 211
- 5.7.7 $[\text{Fe}^{\text{I}}(\text{cAAC})_2][\text{B}(\text{C}_6\text{F}_5)_4]$ 212
- 5.7.8 $[\text{K}(\text{L})][\text{Fe}^{\text{I}}\{\text{N}(\text{SiMe}_3)_3\}_2]$ 213
- 5.7.9 $[\text{Fe}^{\text{II}}(\text{Eind})_2]$ 214
- 5.8 Application of Mössbauer Spectroscopy to Fe/4f Single-Molecule Magnets 215
- 5.8.1 $[\text{Fe}^{\text{III}}_4\text{Dy}^{\text{III}}_4(\text{teaH})_8(\text{N}_3)_8(\text{H}_2\text{O})] \cdot \text{H}_2\text{O} \cdot 4\text{MeCN}$ 215
- 5.8.2 $[\text{Fe}^{\text{III}}_2\text{Ln}^{\text{III}}_2(\text{OH})_2(\text{teaH})_2(\text{O}_2\text{CCPh})_6] \cdot 3\text{MeCN}$ ($\text{Ln}^{\text{III}} = \text{Ce}^{\text{III}}$ to Yb^{III}) 217
- 5.8.3 $[\text{Fe}^{\text{III}}_4\text{Ln}^{\text{III}}_2(\text{teaH})_4(\text{N}_3)_7(\text{piv})_3] \cdot (\text{Ln}^{\text{III}} = \text{Y}^{\text{III}}, \text{Gd}^{\text{III}}, \text{Tb}^{\text{III}}, \text{Dy}^{\text{III}}, \text{Ho}^{\text{III}}, \text{Er}^{\text{III}})$ 218
- 5.8.4 $[\text{Fe}^{\text{III}}_4\text{Dy}^{\text{III}}_2(\text{OH})_2(n\text{-bdea})_4(\text{C}_6\text{H}_5\text{CO}_2)_8] \cdot \text{MeCN}$ 219
- 5.8.5 $[\text{Fe}^{\text{III}}_4\text{Dy}^{\text{III}}_2(\text{OH})_2(n\text{-bdea})_4((\text{CH}_3)_3\text{CCO}_2)_6(\text{N}_3)_2] \cdot 3\text{MeCN}$ 221
- 5.8.6 $[\text{Fe}_7\text{Dy}_3(\mu_4\text{-O})_2(\mu_3\text{-OH})_2(\text{mdea})_7(\mu\text{-benzoate})_4(\text{N}_3)_6] \cdot 2\text{H}_2\text{O} \cdot 7\text{MeOH}$ 222
- 5.8.7 $[\text{Fe}_4\text{Dy}_2(\mu_4\text{-O})_2(\text{NO}_3)_2(\text{piv})_6(\text{Hedte})_2] \cdot 4\text{MeCN} \cdot \text{C}_6\text{H}_5\text{OH}$ 223
- 5.8.8 $[\text{Fe}^{\text{III}}_2\text{Dy}_2(\mu_3\text{-OH})_2(\text{teg})_2(\text{N}_3)_2(\text{C}_6\text{H}_5\text{CO}_2)_4]$ 224
- 5.8.9 $[\text{Fe}^{\text{III}}_2\text{Dy}_2(\mu_3\text{-OH})_2(\text{pmide})_2(p\text{-Me-C}_6\text{H}_5\text{CO}_2)_6]$ 225
- 5.8.10 $[\text{Fe}^{\text{III}}_2\text{Dy}^{\text{III}}_2(\text{OH})_2(\text{L}^1)_2(\text{HL}^2)_2(\text{NO}_3)_4(\text{H}_2\text{O})_{1.5}(\text{MeOH})_{0.5}] \cdot 6\text{MeCN}$ 225
- 5.8.11 $[\text{Fe}^{\text{III}}_2\text{Ln}_2(\text{H}_2\text{L})_4(\text{NO}_3)_2](\text{ClO}_4)_2 \cdot 2\text{MeOH} \cdot 2\text{H}_2\text{O}$ ($\text{Ln} = \text{Gd}^{\text{III}}, \text{Dy}^{\text{III}}, \text{Tb}^{\text{III}}$) 228
- 5.8.12 $[\text{Fe}^{\text{III}}_3\text{Ln}(\mu_3\text{-O})_2(\text{CCl}_3\text{CO}_2)_8(\text{H}_2\text{O})(\text{thf})_3] \cdot x(\text{thf}) \cdot y(\text{heptane})$ ($\text{Ln}^{\text{III}} = \text{Ce}^{\text{III}}\text{-Ho}^{\text{III}}, \text{Lu}^{\text{III}}, \text{Y}^{\text{III}}$) 229
- 5.8.13 $[\text{Fe}^{\text{II}}_2\text{Dy}(\text{L})_2(\text{H}_2\text{O})](\text{ClO}_4)_2 \cdot 2\text{H}_2\text{O}$ 230
- 5.9 Application of Mössbauer Spectroscopy to Cyanide-Bridged Complexes 231
- 5.10 Other Spectroscopic Techniques Used to Study Iron-Based SMMs 234
- 5.11 Conclusions 236
- References 237
- 6 Synthesis and Chemistry of Single-molecule Magnets 245**
Zoi G. Lada, Eugenia Katsoulakou, and Spyros P. Perlepes
- 6.1 General Introduction for the Synthesis of SMMs and SIMs-Organization of the Chapter 246
- 6.2 Synthetic Aspects for Polynuclear 3d Metal SMMs 247

- 6.2.1 Approaches Using Simple 3d Metal Starting Materials 250
- 6.2.2 Approaches Using Preformed Coordination Clusters or SMMs as Starting Materials – Retention of Nuclearity 254
- 6.2.3 Approaches Using Preformed Coordination Clusters or SMMs as Starting Materials – Change of Nuclearity 258
- 6.3 Synthetic Aspects for Dinuclear and Polynuclear 4f Metal Complexes with SMM Properties 262
- 6.3.1 O-Bridged Groups 264
- 6.3.2 Chlorido Bridges 265
- 6.3.3 Monoatomic and Polyatomic N-based Ligands 266
- 6.3.4 Sulfur-bridged SMMs 267
- 6.3.5 Organometallic Bridges 267
- 6.3.6 Radical-bridged Lanthanide(III) SMMs 268
- 6.3.7 Multidecker Lanthanide(III)-Phthalocyanine SMMs 270
- 6.4 Synthetic Aspects for Dinuclear and Polynuclear Actinide SMMs 271
- 6.5 Synthesis of 3d/4f-, 3d/5f-, 4f/5f-Metal and Other Heterometallic SMMs 273
- 6.5.1 3d/4f-Metal SMMs 273
- 6.5.2 3d/5f-Metal SMMs 279
- 6.5.3 4f/5f-Metal Clusters and SMMs 281
- 6.5.4 Other Heterometallic SMMs – the Synthetic Utility of the Cyano Ligand 282
- 6.6 Synthesis of 3d Metal SIMs 286
- 6.7 Synthetic Methodology for 4f Metal SIMs 289
- 6.7.1 Phthalocyanine-based 4f Metal SIMs 290
- 6.7.2 Non-phthalocyanine 4f Metal SIMs 291
- 6.8 Synthetic Routes for 5f Metal SIMs 296
- 6.9 Concluding Comments in Brief–Prognosis for the Future 301
- References 303
- 7 Breakthrough in Radical-bridged Single-molecule Magnets 315**
Constantinos Efthymiou, Meghan Winterlich, and Constantina Papatriantafyllopoulou
- 7.1 General Information About Organic Radicals and Their Magnetic Properties 316
- 7.2 3d Metal Radical SMMs 318
- 7.2.1 Nitroxide Radical SMMs 318
- 7.2.2 Carbene Radical SMMs 321
- 7.2.3 Benzosemiquinonoid and Nindigo Radical SMMs 323
- 7.3 4f Metal Radical SMMs 325
- 7.3.1 Phthalocyanine Radical SMMs 325
- 7.3.2 Nitroxide Radical SMMs 331
- 7.3.3 N_2^{3-} Radical SMMs 336
- 7.3.4 Other 4f Radical SMMs 338
- 7.4 3d–4f Metal Radical SMMs 340
- 7.5 5f Metal Radical SMMs 342

| | | |
|----------|---|------------|
| 7.6 | Conclusions | 343 |
| | References | 345 |
| 8 | Assembly of Polynuclear Single-molecule Magnets | 353 |
| | <i>Kieran Griffiths, Vladislav A. Blatov, and George E. Kostakis</i> | |
| 8.1 | Introduction | 353 |
| 8.2 | History | 354 |
| 8.3 | Topological Methods in Crystal Chemistry and Coordination Chemistry | 356 |
| 8.3.1 | General Overview of ToposPro | 357 |
| 8.3.2 | Example of the ToposPro Analysis of Polynuclear Coordination Clusters | 360 |
| 8.4 | Polynuclear Coordination Clusters Assembly and Topology | 362 |
| 8.5 | 3d–4f PCCs | 363 |
| 8.5.1 | Synthetic Approach for 3d–4f PCCs | 363 |
| 8.5.2 | 3d–4f SMMs PCCs | 366 |
| 8.6 | Assembly Examples and Graph Comparison | 368 |
| 8.7 | Targeting for New Topologies | 373 |
| 8.8 | Synthetic Aspects in Recent Examples | 377 |
| 8.9 | Perspective | 380 |
| | References | 380 |
| | Annexure | 389 |
| | About the Authors | 415 |
| | Index | 421 |