

# Summary of contents

<b>Preface</b>	v
<b>Learning from this book</b>	vii
<b>About the authors</b>	x
<b>List of boxes</b>	xxiii
<b>Reviewer acknowledgments</b>	xxv
<b>Chapter 1</b> History and basic concepts	1
<b>Chapter 2</b> Development of the <i>Drosophila</i> body plan	37
<b>Chapter 3</b> Vertebrate development I: life cycles and experimental techniques	94
<b>Chapter 4</b> Vertebrate development II: <i>Xenopus</i> and zebrafish	142
<b>Chapter 5</b> Vertebrate development III: chick and mouse—completing the body plan	183
<b>Chapter 6</b> Development of nematodes and sea urchins	235
<b>Chapter 7</b> Morphogenesis: change in form in the early embryo	271
<b>Chapter 8</b> Cell differentiation and stem cells	333
<b>Chapter 9</b> Germ cells, fertilization, and sex determination	397
<b>Chapter 10</b> Organogenesis	435
<b>Chapter 11</b> Development of the nervous system	505
<b>Chapter 12</b> Growth, post-embryonic development, and regeneration	553
<b>Chapter 13</b> Plant development	609
<b>Chapter 14</b> Evolution and development	651
<b>Glossary</b>	702
<b>Index</b>	725

# Contents

<i>Preface</i>	v
<i>Learning from this book</i>	vii
<i>About the authors</i>	x
<i>List of boxes</i>	xxiii
<i>Reviewer acknowledgments</i>	xxv
<b>Chapter 1 History and basic concepts</b>	<b>1</b>
<b>The origins of developmental biology</b>	<b>3</b>
<b>1.1</b> Aristotle first defined the problem of epigenesis versus preformation	3
■ <b>Box 1A</b> Basic stages of <i>Xenopus laevis</i> development	4
<b>1.2</b> Cell theory changed how people thought about embryonic development and heredity	4
<b>1.3</b> Two main types of development were originally proposed	6
■ <b>Cell Biology Box 1B</b> The mitotic cell cycle	7
<b>1.4</b> The discovery of induction showed that one group of cells could determine the development of neighboring cells	8
<b>1.5</b> Developmental biology emerged from the coming together of genetics and embryology	8
<b>1.6</b> Development is studied mainly through selected model organisms	9
<b>1.7</b> The first developmental genes were identified as spontaneous mutations	11
<b>Summary</b>	13
<b>A conceptual tool kit</b>	13
<b>1.8</b> Development involves the emergence of pattern, change in form, cell differentiation, and growth	14
■ <b>Cell Biology Box 1C</b> Germ layers	15
<b>1.9</b> Cell behavior provides the link between gene action and developmental processes	17
<b>1.10</b> Genes control cell behavior by specifying which proteins are made	17
<b>1.11</b> The expression of developmental genes is under tight control	19
■ <b>Experimental Box 1D</b> Visualizing gene expression in embryos	20
<b>1.12</b> Development is progressive and the fates of cells become determined at different times	22
<b>1.13</b> Inductive interactions make cells different from each other	24
■ <b>Cell Biology Box 1E</b> Signal transduction and intracellular signaling pathways	26
<b>1.14</b> The response to inductive signals depends on the state of the cell	26
<b>1.15</b> Patterning can involve the interpretation of positional information	27
■ <b>Medical Box 1F</b> When development goes awry	28
<b>1.16</b> Lateral inhibition can generate spacing patterns	30
<b>1.17</b> Localization of cytoplasmic determinants and asymmetric cell division can make daughter cells different from each other	30
<b>1.18</b> The embryo contains a generative rather than a descriptive program	32
<b>1.19</b> The reliability of development is achieved by various means	32
<b>1.20</b> The complexity of embryonic development is due to the complexity of cells themselves	33
<b>1.21</b> Development is a central element in evolution	33
<b>Summary</b>	34
<b>Summary to Chapter 1</b>	35
<b>Chapter 2 Development of the <i>Drosophila</i> body plan</b>	<b>37</b>
<b><i>Drosophila</i> life cycle and overall development</b>	<b>38</b>
<b>2.1</b> The early <i>Drosophila</i> embryo is a multinucleate syncytium	38
<b>2.2</b> Cellularization is followed by gastrulation and segmentation	40
<b>2.3</b> After hatching, the <i>Drosophila</i> larva develops through several larval stages, pupates, and then undergoes metamorphosis to become an adult	41
<b>2.4</b> Many developmental genes were identified in <i>Drosophila</i> through large-scale genetic screening for induced mutations	42
■ <b>Experimental Box 2A</b> Mutagenesis and genetic screening strategy for identifying developmental mutants in <i>Drosophila</i>	43
<b>Summary</b>	44
<b>Setting up the body axes</b>	44
<b>2.5</b> The body axes are set up while the <i>Drosophila</i> embryo is still a syncytium	45

<b>2.6</b> Maternal factors set up the body axes and direct the early stage of <i>Drosophila</i> development	46	<b>Summary</b>	75
<b>2.7</b> Three classes of maternal genes specify the antero-posterior axis	47	<b>Segmentation genes and segment patterning</b>	75
<b>2.8</b> Bicoid protein provides an antero-posterior gradient of a morphogen	48	<b>2.23</b> Expression of the <i>engrailed</i> gene defines the boundary of a parasegment, which is also a boundary of cell lineage restriction	76
<b>2.9</b> The posterior pattern is controlled by the gradients of Nanos and Caudal proteins	50	<b>2.24</b> Segmentation genes stabilize parasegment boundaries	77
<b>2.10</b> The anterior and posterior extremities of the embryo are specified by activation of a cell-surface receptor	51	<b>2.25</b> Signals generated at the parasegment boundary delimit and pattern the future segments	78
<b>2.11</b> The dorso-ventral polarity of the embryo is specified by localization of maternal proteins in the egg vitelline envelope	52	■ <b>Cell Biology Box 2E</b> The Hedgehog signaling pathway	80
<b>2.12</b> Positional information along the dorso-ventral axis is provided by the Dorsal protein	53	■ <b>Experimental Box 2F</b> Mutants in denticle pattern provided clues to the logic of segment patterning	81
■ <b>Cell Biology Box 2B</b> The Toll signaling pathway: a multifunctional pathway	54	<b>Summary</b>	83
<b>Summary</b>	54	<b>Specification of segment identity</b>	83
<b>Localization of maternal determinants during oogenesis</b>	55	<b>2.26</b> Segment identity in <i>Drosophila</i> is specified by Hox genes	84
<b>2.13</b> The antero-posterior axis of the <i>Drosophila</i> egg is specified by signals from the preceding egg chamber and by interactions of the oocyte with follicle cells	56	<b>2.27</b> Homeotic selector genes of the bithorax complex are responsible for diversification of the posterior segments	85
■ <b>Cell Biology Box 2C</b> The JAK-STAT signaling pathway	58	<b>2.28</b> The Antennapedia complex controls specification of anterior regions	86
<b>2.14</b> Localization of maternal mRNAs to either end of the egg depends on the reorganization of the oocyte cytoskeleton	58	<b>2.29</b> The order of Hox gene expression corresponds to the order of genes along the chromosome	87
<b>2.15</b> The dorso-ventral axis of the egg is specified by movement of the oocyte nucleus followed by signaling between oocyte and follicle cells	60	<b>2.30</b> The <i>Drosophila</i> head region is specified by genes other than the Hox genes	87
<b>Summary</b>	61	<b>Summary</b>	88
<b>Patterning the early embryo</b>	62	<b>Summary to Chapter 2</b>	89
<b>2.16</b> The expression of zygotic genes along the dorso-ventral axis is controlled by Dorsal protein	62	<b>Chapter 3 Vertebrate development I: life cycles and experimental techniques</b>	<b>94</b>
<b>2.17</b> The Decapentaplegic protein acts as a morphogen to pattern the dorsal region	64	<b>Vertebrate life cycles and outlines of development</b>	95
<b>2.18</b> The antero-posterior axis is divided up into broad regions by gap gene expression	66	<b>3.1</b> The frog <i>Xenopus laevis</i> is the model amphibian for studying development of the body plan	98
<b>2.19</b> Bicoid protein provides a positional signal for the anterior expression of zygotic <i>hunchback</i>	<b>67</b>	<b>3.2</b> The zebrafish embryo develops around a large mass of yolk	102
<b>2.20</b> The gradient in Hunchback protein activates and represses other gap genes	68	<b>3.3</b> Birds and mammals resemble each other and differ from <i>Xenopus</i> in some important features of early development	105
■ <b>Experimental Box 2D</b> Targeted gene expression and misexpression screening	69	<b>3.4</b> The early chicken embryo develops as a flat disc of cells overlying a massive yolk	106
<b>Summary</b>	70	<b>3.5</b> The mouse egg has no yolk and early development involves the allocation of cells to form the placenta and extra-embryonic membranes	110
<b>Activation of the pair-rule genes and the establishment of parasegments</b>	71	<b>Experimental approaches to studying vertebrate development</b>	115
<b>2.21</b> Parasegments are delimited by expression of pair-rule genes in a periodic pattern	71	<b>3.6</b> Gene expression in embryos can be mapped by <i>in situ</i> nucleic acid hybridization	116
<b>2.22</b> Gap gene activity positions stripes of pair-rule gene expression	73	■ <b>Experimental Box 3A</b> Gene-expression profiling by DNA microarrays and RNA seq	117
		<b>3.7</b> Fate mapping and lineage tracing reveal which cells in which parts of the early embryo give rise to particular adult structures	118

<b>3.8</b> Not all techniques are equally applicable to all vertebrates	120	<b>4.7</b> Mesoderm induction occurs during a limited period in the blastula stage	155
<b>3.9</b> Developmental genes can be identified by spontaneous mutation and by large-scale mutagenesis screens	121	<b>4.8</b> Zygotic gene expression is turned on at the mid-blastula transition	156
■ <b>Experimental Box 3B</b> Large-scale mutagenesis screens for recessive mutations in zebrafish	123	<b>4.9</b> Mesoderm-inducing and patterning signals are produced by the vegetal region, the organizer, and the ventral mesoderm	157
<b>3.10</b> Transgenic techniques enable animals to be produced with mutations in specific genes	124	<b>4.10</b> Members of the TGF- $\beta$ family have been identified as mesoderm inducers	158
■ <b>Experimental Box 3C</b> The Cre/ <i>loxP</i> system: a strategy for making gene knock-outs in mice	127	■ <b>Experimental Box 4D</b> Investigating receptor function using dominant-negative proteins	159
■ <b>Experimental Box 3D</b> The CRISPR-Cas9 genome-editing system	128	<b>4.11</b> The zygotic expression of mesoderm-inducing and patterning signals is activated by the combined actions of maternal VegT and Wnt signaling	159
<b>3.11</b> Gene function can also be tested by transient transgenesis and gene silencing	130	<b>4.12</b> Threshold responses to gradients of signaling proteins are likely to pattern the mesoderm	161
<b>Human embryonic development</b>	131	<b>Summary</b>	162
<b>3.12</b> The early development of a human embryo is similar to that of the mouse	131	<b>The Spemann organizer and neural induction</b>	163
■ <b>Medical Box 3E</b> Preimplantation genetic diagnosis	134	■ <b>Cell Biology Box 4E</b> The fibroblast growth factor signaling pathway	163
<b>3.13</b> The timing of formation and the anatomy of the human placenta differs from that in the mouse	135	<b>4.13</b> Signals from the organizer pattern the mesoderm dorso-ventrally by antagonizing the effects of ventral signals	164
<b>3.14</b> Some studies of human development are possible but are subject to strict laws	136	<b>4.14</b> The antero-posterior axis of the embryo emerges during gastrulation	165
■ <b>Box 3F</b> Identical twins	137	<b>4.15</b> The neural plate is induced in the ectoderm	168
<b>Summary to Chapter 3</b>	138	<b>4.16</b> The nervous system is patterned along the antero-posterior axis by signals from the mesoderm	170
<b>Chapter 4 Vertebrate development II: <i>Xenopus</i> and zebrafish</b>	<b>142</b>	<b>4.17</b> The final body plan emerges by the end of gastrulation and neurulation	171
<b>Setting up the body axes</b>	143	<b>Summary</b>	172
<b>4.1</b> The animal-vegetal axis is maternally determined in <i>Xenopus</i>	143	<b>Development of the body plan in zebrafish</b>	172
■ <b>Cell Biology Box 4A</b> Intercellular protein signals in vertebrate development	145	<b>4.18</b> The body axes in zebrafish are established by maternal determinants	173
■ <b>Cell Biology Box 4B</b> The Wnt/ $\beta$ -catenin signaling pathway	146	<b>4.19</b> The germ layers are specified in the zebrafish blastoderm by similar signals to those in <i>Xenopus</i>	173
<b>4.2</b> Local activation of Wnt/ $\beta$ -catenin signaling specifies the future dorsal side of the embryo	147	<b>4.20</b> The shield in zebrafish is the embryonic organizer	176
<b>4.3</b> Signaling centers develop on the dorsal side of the blastula	149	■ <b>Box 4F</b> A zebrafish gene regulatory network	176
<b>Summary</b>	150	<b>Summary to Chapter 4</b>	178
<b>The origin and specification of the germ layers</b>	150	<b>Chapter 5 Vertebrate development III: chick and mouse—completing the body plan</b>	<b>183</b>
<b>4.4</b> The fate map of the <i>Xenopus</i> blastula makes clear the function of gastrulation	151	<b>Development of the body plan in chick and mouse and generation of the spinal cord</b>	184
<b>4.5</b> Cells of the early <i>Xenopus</i> embryo do not yet have their fates determined and regulation is possible	152	<b>5.1</b> The antero-posterior polarity of the chick blastoderm is related to the primitive streak	184
<b>4.6</b> Endoderm and ectoderm are specified by maternal factors, whereas mesoderm is induced from ectoderm by signals from the vegetal region	152	<b>5.2</b> Early stages in mouse development establish separate cell lineages for the embryo and the extra-embryonic structures	186
■ <b>Cell Biology Box 4C</b> Signaling by members of the TGF- $\beta$ family of growth factors	155		

<b>5.3</b> Movement of the anterior visceral endoderm indicates the definitive antero-posterior axis in the mouse embryo	190
<b>5.4</b> The fate maps of vertebrate embryos are variations on a basic plan	192
■ <b>Cell Biology Box 5A</b> Fine-tuning Nodal signaling	193
<b>5.5</b> Mesoderm induction and patterning in the chick and mouse occurs during primitive streak formation	195
<b>5.6</b> The node that develops at the anterior end of the streak in chick and mouse embryos is equivalent to the Spemann organizer in <i>Xenopus</i>	196
<b>5.7</b> Neural induction in chick and mouse is initiated by FGF signaling with inhibition of BMP signaling being required in a later step	198
■ <b>Cell Biology Box 5B</b> Chromatin-remodeling complexes	201
<b>5.8</b> Axial structures in chick and mouse are generated from self-renewing cell populations	202
<b>Summary</b>	204
<b>Somite formation and antero-posterior patterning</b>	205
■ <b>Cell Biology Box 5C</b> Retinoic acid: a small-molecule intercellular signal	206
<b>5.9</b> Somites are formed in a well-defined order along the antero-posterior axis	206
■ <b>Cell Biology Box 5D</b> The Notch signaling pathway	211
<b>5.10</b> Identity of somites along the antero-posterior axis is specified by Hox gene expression	213
■ <b>Box 5E</b> The Hox genes	214
<b>5.11</b> Deletion or overexpression of Hox genes causes changes in axial patterning	217
<b>5.12</b> Hox gene expression is activated in an anterior to posterior pattern	219
<b>5.13</b> The fate of somite cells is determined by signals from the adjacent tissues	221
<b>Summary</b>	223
<b>The origin and patterning of neural crest</b>	223
<b>5.14</b> Neural crest cells arise from the borders of the neural plate and migrate to give rise to a wide range of different cell types	223
<b>5.15</b> Neural crest cells migrate from the hindbrain to populate the branchial arches	225
<b>Summary</b>	226
<b>Determination of left-right asymmetry</b>	227
<b>5.16</b> The bilateral symmetry of the early embryo is broken to produce left-right asymmetry of internal organs	227
<b>5.17</b> Left-right symmetry breaking may be initiated within cells of the early embryo	229
<b>Summary</b>	230
<b>Summary to Chapter 5</b>	230
<b>Chapter 6 Development of nematodes and sea urchins</b>	<b>235</b>
<b>Nematodes</b>	236
■ <b>Cell Biology Box 6A</b> Apoptotic pathways in nematodes, <i>Drosophila</i> , and mammals	238
<b>6.1</b> The cell lineage of <i>Caenorhabditis elegans</i> is largely invariant	239
<b>6.2</b> The antero-posterior axis in <i>Caenorhabditis elegans</i> is determined by asymmetric cell division	239
■ <b>Experimental Box 6B</b> Gene silencing by antisense RNA and RNA interference	241
<b>6.3</b> The dorso-ventral axis in <i>Caenorhabditis elegans</i> is determined by cell-cell interactions	242
<b>6.4</b> Both asymmetric divisions and cell-cell interactions specify cell fate in the early nematode embryo	245
<b>6.5</b> Cell differentiation in the nematode is closely linked to the pattern of cell division	246
<b>6.6</b> Hox genes specify positional identity along the antero-posterior axis in <i>Caenorhabditis elegans</i>	247
<b>6.7</b> The timing of events in nematode development is under genetic control that involves microRNAs	248
■ <b>Box 6C</b> Gene silencing by microRNAs	250
<b>6.8</b> Vulval development is initiated through the induction of a small number of cells by short-range signals from a single inducing cell	251
<b>Summary</b>	253
<b>Echinoderms</b>	254
<b>6.9</b> The sea urchin embryo develops into a free-swimming larva	255
<b>6.10</b> The sea urchin egg is polarized along the animal-vegetal axis	256
<b>6.11</b> The sea urchin fate map is finely specified, yet considerable regulation is possible	257
<b>6.12</b> The vegetal region of the sea urchin embryo acts as an organizer	258
<b>6.13</b> The sea urchin vegetal region is demarcated by the nuclear accumulation of $\beta$ -catenin	260
<b>6.14</b> The animal-vegetal axis and the oral-aboral axis can be considered to correspond to the antero-posterior and dorso-ventral axes of other deuterostomes	261
<b>6.15</b> The pluteus skeleton develops from the primary mesenchyme	262
<b>6.16</b> The oral-aboral axis in sea urchins is related to the plane of the first cleavage	263
<b>6.17</b> The oral ectoderm acts as an organizing region for the oral-aboral axis	264

■ <b>Experimental Box 6D</b> The gene regulatory network for sea urchin endomesoderm specification	265
<b>Summary</b>	266
<b>Summary to Chapter 6</b>	267
<b>Chapter 7 Morphogenesis: change in form in the early embryo</b>	<b>271</b>
<b>Cell adhesion</b>	273
■ <b>Cell Biology Box 7A</b> Cell-adhesion molecules and cell junctions	274
<b>7.1</b> Sorting out of dissociated cells demonstrates differences in cell adhesiveness in different tissues	275
<b>7.2</b> Cadherins can provide adhesive specificity	276
<b>7.3</b> The activity of the cytoskeleton regulates the mechanical properties of cells and their interactions with each other	277
■ <b>Cell Biology Box 7B</b> The cytoskeleton, cell-shape change, and cell movement	278
<b>7.4</b> Transitions of tissues from an epithelial to a mesenchymal state, and vice versa, involve changes in adhesive junctions	279
<b>Summary</b>	280
<b>Cleavage and formation of the blastula</b>	280
<b>7.5</b> The orientation of the mitotic spindle determines the plane of cleavage at cell division	281
<b>7.6</b> The positioning of the spindle within the cell also determines whether daughter cells will be the same or different sizes	283
<b>7.7</b> Cells become polarized in the sea urchin blastula and the mouse morula	285
<b>7.8</b> Fluid accumulation as a result of tight-junction formation and ion transport forms the blastocoel of the mammalian blastocyst	287
<b>Summary</b>	288
<b>Gastrulation movements</b>	289
<b>7.9</b> Gastrulation in the sea urchin involves an epithelial-to-mesenchymal transition, cell migration, and invagination of the blastula wall	289
<b>7.10</b> Mesoderm invagination in <i>Drosophila</i> is due to changes in cell shape controlled by genes that pattern the dorso-ventral axis	293
<b>7.11</b> Germ-band extension in <i>Drosophila</i> involves myosin-dependent remodeling of cell junctions and cell intercalation	295
<b>7.12</b> Planar cell polarity confers directionality on a tissue	296
<b>7.13</b> Gastrulation in amphibians and fish involves involution, epiboly, and convergent extension	299
■ <b>Box 7C</b> Convergent extension	302
<b>7.14</b> <i>Xenopus</i> notochord development illustrates the dependence of medio-lateral cell elongation and cell intercalation on a pre-existing antero-posterior polarity	305
<b>7.15</b> Gastrulation in chick and mouse embryos involves the separation of individual cells from the epiblast and their ingression through the primitive streak	306
<b>Summary</b>	309
<b>Neural tube formation</b>	311
<b>7.16</b> Neural tube formation is driven by changes in cell shape and convergent extension	311
■ <b>Cell Biology Box 7D</b> Eph receptors and their ephrin ligands	313
■ <b>Medical Box 7E</b> Neural tube defects	314
<b>Summary</b>	315
<b>Formation of tubes and branching morphogenesis</b>	316
<b>7.17</b> The <i>Drosophila</i> tracheal system is a prime example of branching morphogenesis	316
<b>7.18</b> The vertebrate vascular system develops by vasculogenesis followed by sprouting angiogenesis	318
<b>7.19</b> New blood vessels are formed from pre-existing vessels in angiogenesis	319
<b>Summary</b>	320
<b>Cell migration</b>	320
<b>7.20</b> Embryonic neural crest gives rise to a wide range of different cell types	321
<b>7.21</b> Neural crest migration is controlled by environmental cues	321
<b>7.22</b> The formation of the lateral-line primordium in fishes is an example of collective cell migration	323
<b>7.23</b> Body wall closure occurs in <i>Drosophila</i> , <i>Caenorhabditis</i> , mammals, and chick	324
<b>Summary</b>	325
<b>Summary to Chapter 7</b>	326
<b>Chapter 8 Cell differentiation and stem cells</b>	<b>333</b>
■ <b>Box 8A</b> Conrad Waddington's 'epigenetic landscape' provides a framework for thinking about how cells differentiate	335
<b>The control of gene expression</b>	337
<b>8.1</b> Control of transcription involves both general and tissue-specific transcriptional regulators	338
<b>8.2</b> Gene expression is also controlled by epigenetic chemical modifications to DNA and histone proteins that alter chromatin structure	341
■ <b>Cell Biology Box 8B</b> Epigenetic control of gene expression by chromatin modification	344
<b>8.3</b> Patterns of gene activity can be inherited by persistence of gene-regulatory proteins or by maintenance of chromatin modifications	347
<b>8.4</b> Changes in patterns of gene activity during differentiation can be triggered by extracellular signals	348
<b>Summary</b>	349

<b>Cell differentiation and stem cells</b>	350	<b>Chapter 9 Germ cells, fertilization, and sex determination</b>	<b>397</b>
<b>8.5</b> Muscle differentiation is determined by the MyoD family of transcription factors	350	<b>The development of germ cells</b>	398
<b>8.6</b> The differentiation of muscle cells involves withdrawal from the cell cycle, but is reversible	352	<b>9.1</b> Germ cell fate is specified in some embryos by a distinct germplasm in the egg	399
<b>8.7</b> All blood cells are derived from multipotent stem cells	354	<b>9.2</b> In mammals germ cells are induced by cell-cell interactions during development	401
<b>8.8</b> Intrinsic and extrinsic changes control differentiation of the hematopoietic lineages	357	<b>9.3</b> Germ cells migrate from their site of origin to the gonad	402
■ <b>Experimental Box 8C</b> Single-cell analysis of cell-fate decisions	358	<b>9.4</b> Germ cells are guided to their destination by chemical signals	403
<b>8.9</b> Developmentally regulated globin gene expression is controlled by control regions far distant from the coding regions	361	<b>9.5</b> Germ cell differentiation involves a halving of chromosome number by meiosis	404
<b>8.10</b> The epidermis of adult mammalian skin is continually being replaced by derivatives of stem cells	363	■ <b>Box 9A</b> Polar bodies	405
■ <b>Medical Box 8D</b> Treatment of junctional epidermolysis bullosa with skin grown from genetically corrected stem cells	366	<b>9.6</b> Oocyte development can involve gene amplification and contributions from other cells	408
<b>8.11</b> Stem cells use different modes of division to maintain tissues	367	<b>9.7</b> Factors in the cytoplasm maintain the totipotency of the egg	408
<b>8.12</b> The lining of the gut is another epithelial tissue that requires continuous renewal	368	<b>9.8</b> In mammals some genes controlling embryonic growth are 'imprinted'	409
<b>8.13</b> Skeletal muscle and neural cells can be renewed from stem cells in adults	370	<b>Summary</b>	412
<b>8.14</b> Embryonic stem cells can proliferate and differentiate into many cell types in culture and contribute to normal development <i>in vivo</i>	372	<b>Fertilization</b>	412
■ <b>Experimental Box 8E</b> The derivation and culture of mouse embryonic stem cells	374	<b>9.9</b> Fertilization involves cell-surface interactions between egg and sperm	413
<b>Summary</b>	375	<b>9.10</b> Changes in the egg plasma membrane and enveloping layers at fertilization block polyspermy	415
<b>The plasticity of the differentiated state</b>	376	<b>9.11</b> Sperm-egg fusion causes a calcium wave that results in egg activation	415
<b>8.15</b> Nuclei of differentiated cells can support development	376	<b>Summary</b>	418
<b>8.16</b> Patterns of gene activity in differentiated cells can be changed by cell fusion	378	<b>Determination of the sexual phenotype</b>	419
<b>8.17</b> The differentiated state of a cell can change by transdifferentiation	379	<b>9.12</b> The primary sex-determining gene in mammals is on the Y chromosome	419
<b>8.18</b> Adult differentiated cells can be reprogrammed to form pluripotent stem cells	381	<b>9.13</b> Mammalian sexual phenotype is regulated by gonadal hormones	420
■ <b>Experimental Box 8F</b> Induced pluripotent stem cells	382	<b>9.14</b> The primary sex-determining factor in <i>Drosophila</i> is the number of X chromosomes and is cell autonomous	422
<b>8.19</b> Stem cells could be a key to regenerative medicine	382	<b>9.15</b> Somatic sexual development in <i>Caenorhabditis</i> is determined by the number of X chromosomes	424
■ <b>Experimental Box 8G</b> Stem cells can be cultured <i>in vitro</i> to produce 'organoids'—structures that mimic tissues and organs	386	<b>9.16</b> Determination of germ cell sex depends on both genetic constitution and intercellular signals	425
<b>8.20</b> Various approaches can be used to generate differentiated cells for cell-replacement therapies	388	<b>9.17</b> Various strategies are used for dosage compensation of X-linked genes	427
<b>Summary</b>	391	<b>Summary</b>	429
<b>Summary to Chapter 8</b>	391	<b>Summary to Chapter 9</b>	431
		<b>Chapter 10 Organogenesis</b>	<b>435</b>
		<b>The insect wing and leg</b>	436
		<b>10.1</b> Imaginal discs arise from the ectoderm in the early <i>Drosophila</i> embryo	437

<b>10.2</b> Imaginal discs arise across parasegment boundaries and are patterned by signaling at compartment boundaries	438
<b>10.3</b> The adult wing emerges at metamorphosis after folding and evagination of the wing imaginal disc	439
<b>10.4</b> A signaling center at the boundary between anterior and posterior compartments patterns the <i>Drosophila</i> wing disc along the antero-posterior axis	440
■ <b>Box 10A</b> Positional information and morphogen gradients	443
<b>10.5</b> A signaling center at the boundary between dorsal and ventral compartments patterns the <i>Drosophila</i> wing along the dorso-ventral axis	445
<b>10.6</b> Vestigial is a key regulator of wing development that acts to specify wing identity and control wing growth	445
<b>10.7</b> The <i>Drosophila</i> wing disc is also patterned along the proximo-distal axis	447
<b>10.8</b> The leg disc is patterned in a similar manner to the wing disc, except for the proximo-distal axis	448
<b>10.9</b> Different imaginal discs can have the same positional values	450
<b>Summary</b>	450
<b>The vertebrate limb</b>	452
<b>10.10</b> The vertebrate limb develops from a limb bud and its development illustrates general principles	452
<b>10.11</b> Genes expressed in the lateral plate mesoderm are involved in specifying limb position, polarity, and identity	454
<b>10.12</b> The apical ectodermal ridge is required for limb-bud outgrowth and the formation of structures along the proximo-distal axis of the limb	457
<b>10.13</b> Formation and outgrowth of the limb bud involves oriented cell behavior	458
<b>10.14</b> Positional value along the proximo-distal axis of the limb bud is specified by a combination of graded signaling and a timing mechanism	460
<b>10.15</b> The polarizing region specifies position along the limb's antero-posterior axis	462
<b>10.16</b> Sonic hedgehog is the polarizing region morphogen	464
■ <b>Medical Box 10B</b> Too many fingers: mutations that affect antero-posterior patterning can cause polydactyly	465
■ <b>Cell Biology Box 10C</b> Sonic hedgehog signaling and the primary cilium	466
<b>10.17</b> The dorso-ventral axis of the limb is controlled by the ectoderm	468
■ <b>Medical Box 10D</b> Teratogens and the consequences of damage to the developing embryo	470
<b>10.18</b> Development of the limb is integrated by interactions between signaling centers	470
<b>10.19</b> Hox genes have multiple inputs into the patterning of the limbs	472
<b>10.20</b> Self-organization may be involved in the development of the limb	475
■ <b>Box 10E</b> Reaction-diffusion mechanisms	476
<b>10.21</b> Limb muscle is patterned by the connective tissue	477
<b>10.22</b> The initial development of cartilage, muscles, and tendons is autonomous	478
<b>10.23</b> Joint formation involves secreted signals and mechanical stimuli	478
<b>10.24</b> Separation of the digits is the result of programmed cell death	479
<b>Summary</b>	480
<b>Teeth</b>	481
<b>10.25</b> Tooth development involves epithelial-mesenchymal interactions and a homeobox gene code specifies tooth identity	482
<b>Summary</b>	484
<b>Vertebrate lungs</b>	484
<b>10.26</b> The vertebrate lung develops from a bud of endoderm	484
■ <b>Medical Box 10F</b> What developmental biology can teach us about breast cancer	486
<b>10.27</b> Morphogenesis of the lung involves three modes of branching	488
<b>Summary</b>	489
<b>The vertebrate heart</b>	489
<b>10.28</b> The development of the vertebrate heart involves morphogenesis and patterning of a mesodermal tube	489
<b>The vertebrate eye</b>	492
<b>10.29</b> Development of the vertebrate eye involves interactions between an extension of the forebrain and the ectoderm of the head	493
<b>Summary</b>	497
<b>Summary to Chapter 10</b>	497
<b>Chapter 11 Development of the nervous system</b>	<b>505</b>
<b>Specification of cell identity in the nervous system</b>	507
<b>11.1</b> Initial regionalization of the vertebrate brain involves signals from local organizers	507
<b>11.2</b> Local signaling centers pattern the brain along the antero-posterior axis	508
<b>11.3</b> The cerebral cortex is patterned by signals from the anterior neural ridge	509
<b>11.4</b> The hindbrain is segmented into rhombomeres by boundaries of cell-lineage restriction	509
<b>11.5</b> Hox genes provide positional information in the developing hindbrain	512
<b>11.6</b> The pattern of differentiation of cells along the dorso-ventral axis of the spinal cord depends on ventral and dorsal signals	513



<b>11.7</b> Neuronal subtypes in the ventral spinal cord are specified by the ventral to dorsal gradient of Shh	515	<b>Chapter 12 Growth, post-embryonic development, and regeneration</b>	<b>553</b>
<b>11.8</b> Spinal cord motor neurons at different dorso-ventral positions project to different trunk and limb muscles	516	<b>Growth</b>	554
<b>11.9</b> Antero-posterior pattern in the spinal cord is determined in response to secreted signals from the node and adjacent mesoderm	517	<b>12.1</b> Tissues can grow by cell proliferation, cell enlargement, or accretion	555
<b>Summary</b>	518	<b>12.2</b> Cell proliferation is controlled by regulating entry into the cell cycle	556
<b>The formation and migration of neurons</b>	518	<b>12.3</b> Cell division in early development can be controlled by an intrinsic developmental program	557
<b>11.10</b> Neurons in <i>Drosophila</i> arise from proneural clusters	519	<b>12.4</b> Extrinsic signals coordinate cell division, cell growth, and cell death in the developing <i>Drosophila</i> wing	558
<b>11.11</b> The development of neurons in <i>Drosophila</i> involves asymmetric cell divisions and timed changes in gene expression	521	■ <b>Cell Biology Box 12A</b> The core Hippo signaling pathways in <i>Drosophila</i> and mammals	559
<b>11.12</b> The production of vertebrate neurons involves lateral inhibition, as in <i>Drosophila</i>	522	<b>12.5</b> Cancer can result from mutations in genes that control cell proliferation	560
■ <b>Box 11A</b> Specification of the sensory organs of adult <i>Drosophila</i>	523	<b>12.6</b> The relative contributions of intrinsic and extrinsic factors in controlling size differ in different mammalian organs	562
<b>11.13</b> Neurons are formed in the proliferative zone of the vertebrate neural tube and migrate outwards	524	<b>12.7</b> Overall body size depends on the extent and the duration of growth	564
■ <b>Experimental Box 11B</b> Timing the birth of cortical neurons	526	<b>12.8</b> Hormones and growth factors coordinate the growth of different tissues and organs and contribute to determining overall body size	565
<b>11.14</b> Many cortical interneurons migrate tangentially	528	<b>12.9</b> Elongation of the long bones illustrates how growth can be determined by a combination of an intrinsic growth program and extracellular factors	566
<b>Summary</b>	528	■ <b>Box 12B</b> Digit length ratio is determined in the embryo	568
<b>Axon navigation</b>	529	<b>12.10</b> The amount of nourishment an embryo receives can have profound effects in later life	570
<b>11.15</b> The growth cone controls the path taken by a growing axon	530	<b>Summary</b>	571
■ <b>Box 11C</b> The development of the neural circuit for the knee-jerk reflex	532	<b>Molting and metamorphosis</b>	572
<b>11.16</b> Motor neuron axons in the chick limb are guided by ephrin-Eph interactions	533	<b>12.11</b> Arthropods have to molt in order to grow	572
<b>11.17</b> Axons crossing the midline are both attracted and repelled	534	<b>12.12</b> Insect body size is determined by the rate and duration of larval growth	573
<b>11.18</b> Neurons from the retina make ordered connections with visual centers in the brain	535	<b>12.13</b> Metamorphosis in amphibians is under hormonal control	575
<b>Summary</b>	538	<b>Summary</b>	576
<b>Synapse formation and refinement</b>	539	<b>Regeneration</b>	577
<b>11.19</b> Synapse formation involves reciprocal interactions	539	<b>12.14</b> Regeneration involves repatterning of existing tissues and/or growth of new tissues	578
<b>11.20</b> Many motor neurons die during normal development	542	<b>12.15</b> Amphibian limb regeneration involves cell dedifferentiation and new growth	578
■ <b>Medical Box 11D</b> Autism: a developmental disorder that involves synapse dysfunction	543	■ <b>Box 12C</b> Regeneration in <i>Hydra</i>	580
<b>11.21</b> Neuronal cell death and survival involve both intrinsic and extrinsic factors	544	■ <b>Box 12D</b> Planarian regeneration	582
<b>11.22</b> The map from eye to brain is refined by neural activity	545	<b>12.16</b> Limb regeneration in amphibians depends on the presence of nerves	586
<b>Summary</b>	546	<b>12.17</b> The limb blastema gives rise to structures with positional values distal to the site of amputation	587
<b>Summary to Chapter 11</b>	547		

<b>12.18</b> Retinoic acid can change proximo-distal positional values in regenerating limbs	589	<b>13.12</b> The regular arrangement of leaves on a stem is generated by regulated auxin transport	629
<b>12.19</b> Mammals can regenerate the tips of the digits	590	<b>13.13</b> The outgrowth of secondary shoots is under hormonal control	630
<b>12.20</b> Insect limbs intercalate positional values by both proximo-distal and circumferential growth	591	<b>13.14</b> Root tissues are produced from <i>Arabidopsis</i> root apical meristems by a highly stereotyped pattern of cell divisions	633
■ <b>Box 12E</b> Why can't we regenerate our limbs?	592	<b>13.15</b> Root hairs are specified by a combination of positional information and lateral inhibition	635
<b>12.21</b> Heart regeneration in zebrafish involves the resumption of cell division by cardiomyocytes	594	<b>Summary</b>	636
<b>Summary</b>	596	<b>Flower development and control of flowering</b>	636
<b>Aging and senescence</b>	597	<b>13.16</b> Homeotic genes control organ identity in the flower	637
<b>12.22</b> Genes can alter the timing of senescence	598	■ <b>Box 13C</b> The basic model for the patterning of the <i>Arabidopsis</i> flower	639
<b>12.23</b> Cell senescence blocks cell proliferation	600	<b>13.17</b> The <i>Antirrhinum</i> flower is patterned dorso-ventrally, as well as radially	640
<b>12.24</b> Elimination of senescent cells in adult salamanders explains why regenerative ability does not diminish with age	601	<b>13.18</b> The internal meristem layer can specify floral meristem patterning	641
<b>Summary</b>	602	<b>13.19</b> The transition of a shoot meristem to a floral meristem is under environmental and genetic control	642
<b>Summary to Chapter 12</b>	602	■ <b>Box 13D</b> The circadian clock coordinates plant growth and development	643
<b>Chapter 13 Plant development</b>	<b>609</b>	<b>13.20</b> Vernalization reflects the epigenetic memory of winter	643
<b>13.1</b> The model plant <i>Arabidopsis thaliana</i> has a short life cycle and a small diploid genome	611	<b>13.21</b> Most flowering plants are hermaphrodites, but some produce unisexual flowers	645
<b>Embryonic development</b>	612	<b>Summary</b>	646
<b>13.2</b> Plant embryos develop through several distinct stages	612	<b>Summary to Chapter 13</b>	647
■ <b>Box 13A</b> Angiosperm embryogenesis	614	<b>Chapter 14 Evolution and development</b>	<b>651</b>
<b>13.3</b> Gradients of the signal molecule auxin establish the embryonic apical-basal axis	616	■ <b>Box 14A</b> Darwin's finches	654
<b>13.4</b> Plant somatic cells can give rise to embryos and seedlings	617	<b>The evolution of development</b>	655
<b>13.5</b> Cell enlargement is a major process in plant growth and morphogenesis	619	<b>14.1</b> Multicellular organisms evolved from single-celled ancestors	655
■ <b>Experimental Box 13B</b> Plant transformation and genome editing	620	<b>14.2</b> Genomic evidence is throwing light on the evolution of animals	657
<b>Summary</b>	621	■ <b>Box 14B</b> The metazoan family tree	658
<b>Meristems</b>	622	<b>14.3</b> How gastrulation evolved is not known	659
<b>13.6</b> A meristem contains a small, central zone of self-renewing stem cells	623	<b>14.4</b> More general characteristics of the body plan develop earlier than specializations	660
<b>13.7</b> The size of the stem cell area in the meristem is kept constant by a feedback loop to the organizing center	623	<b>14.5</b> Embryonic structures have acquired new functions during evolution	661
<b>13.8</b> The fate of cells from different meristem layers can be changed by changing their position	624	<b>14.6</b> Evolution of different types of eyes in different animal groups is an example of parallel evolution	663
<b>13.9</b> A fate map for the embryonic shoot meristem can be deduced using clonal analysis	626	<b>Summary</b>	664
<b>13.10</b> Meristem development is dependent on signals from other parts of the plant	627	<b>The diversification of body plans</b>	665
<b>13.11</b> Gene activity patterns the proximo-distal and adaxial-abaxial axes of leaves developing from the shoot meristem	628	<b>14.7</b> Hox gene complexes have evolved through gene duplication	665

**14.8** Differences in Hox gene expression determine the variation in position and type of paired appendages in arthropods 667

**14.9** Changes in Hox gene expression and their target genes contributed to the evolution of the vertebrate axial skeleton 671

**14.10** The basic body plan of arthropods and vertebrates is similar, but the dorso-ventral axis is inverted 672

**Summary** 673

**The evolutionary modification of specialized characters** 674

**14.11** Limbs evolved from fins 674

**14.12** Limbs have evolved to fulfill different specialized functions 678

**14.13** The evolution of limblessness in snakes is associated with changes in axial gene expression and mutations in a limb-specific enhancer 679

**14.14** Butterfly wing markings have evolved by redeployment of genes previously used for other functions 680

■ **Experimental Box 14C** Using CRISPR-Cas9 genome-editing techniques to test the functioning of the snake ZRS 681

**14.15** Adaptive evolution within the same species provides a way of studying the developmental basis for evolutionary change 684

■ **Experimental Box 14D** Pelvic reduction in sticklebacks is based on mutations in a gene control region 686

**Summary** 687

**Changes in the timing of developmental processes** 687

**14.16** Changes in growth can modify the basic body plan 687

■ **Box 14E** Origins of morphological diversity in dogs 689

**14.17** Evolution can be due to changes in the timing of developmental events 690

**14.18** The evolution of life histories has implications for development 692

■ **Box 14F** Long- and short-germ development in insects 693

**Summary** 695

**Summary to Chapter 14** 696

**Glossary** 702

**Index** 725