

Contents

Introduction

vii

1.	Magnetic clusters	1
1.1.	Dimers	1
1.1.1.	<i>Spin gap in cesium divanadate</i>	3
1.1.2.	<i>Mineral urusovite</i>	4
1.1.3.	<i>Copper trifluoroacetate</i>	9
1.2.	Trimmers	15
1.2.1.	<i>Trimers of octahedra in rubidium–copper diphosphate</i>	18
1.2.2.	<i>Trimer plaquettes in sodium–copper germanate</i>	24
1.3.	Tetramers	28
1.4.	Bose–Einstein condensation of magnons	33
1.4.1.	<i>Pigment of the Han Dynasty</i>	34
1.4.2.	<i>Barium–vanadium disilicate</i>	35
2.	Quasi-one dimensional magnetics	42
2.1.	Homogeneous chain of half-integer spins	42
2.1.1.	<i>Rubidium–copper molybdate</i>	44
2.1.2.	<i>Vanadyl diacetate</i>	45
2.2.	A homogeneous chain with competing interactions	54
2.2.1.	<i>Lithium–copper zirconate</i>	56
2.2.2.	<i>Isostructural cuprates of lithium and sodium</i>	58
2.2.3.	<i>Rubidium–copper–aluminium phosphate</i>	66
2.2.4.	<i>Cesium–copper vanadium-diphosphate</i>	72
2.2.5.	<i>Bismuth–iron selenite–oxochloride</i>	78
2.3.	Alternating chain of half-integer spins	83
2.4.	A homogeneous chain of integer spins	88
2.5.	Spin–Peierls transition	97
2.6.	Orbital mechanism of dimerization of the spin chain	101
3.	Spin ladders	106
3.1.	Spin ladders with an odd number of legs:	106
3.2.	Spin ladders with an even number of legs:a spin liquid with an energy gap in the spectrum of magnetic excitations	108
3.3.	Charge mechanism of dimerization of the spin ladder	112

Contents

3.4.	Combinations of spin chains and spin ladders	115
3.5.	Frame structures	117
4.	Quasi-two dimensional magnets with a square lattice	124
4.1.	Quantum ground state	124
4.2.	The Berezinsky–Kosterlitz–Thouless transition	126
4.3.	Manganese chromate	129
5.	Quasi-two dimensional magnetics with a triangular lattice	138
5.1.	Geometrical frustration	138
5.1.1.	Lithium–nickel tellurate	142
5.1.2.	Barium–cobalt antimonide	147
5.1.3.	Delafoseites	149
5.2.	BKT transition in two-dimensional magnetic materials with a triangular lattice	151
5.2.1.	Chiral and non-chiral polymorphs of manganese antimonate	153
5.2.2.	Lithium–iron antimonate	167
6.	Quasi-two dimensional magnets with a honeycomb magnetic lattice	178
6.1.	Frustration due to the competition of exchange interactions	178
6.2.	Kitaev model	181
6.3.	Experimental realization of Kitaev's model	182
6.4.	BKT transition in the honeycomb lattice	190
6.5.	Antimonates and tellurates of transition metals	192
6.5.1.	Nickel antimonates	201
6.5.2.	Cobalt antimonates	211
7.	Quasi-two-dimensional magnets with triangular motifs in the structure	219
7.1.	Frustration of exchange interactions in the lattice of kagome and in diamond chains	219
7.1.1.	Herbertsmithite and vésigniéite	221
7.1.2.	The combination of kagome and triangular layers in quasi-2D cobaltites	223
7.1.3.	Langasites	237
7.1.4.	Dugganites	248
7.2.	Plateau of magnetization in the lattice of diamond chains	257
7.3.	Shastry–Sutherland lattice	266
7.4.	Potassium carbonate–manganese vanadate	270
7.5.	Sodium–nickel phosphate hydroxide	275
Conclusion		283
References		285
Index		301