

Contents

| | |
|---|-----|
| Introduction | 5 |
| Chapter 1. | |
| Mössbauer spectroscopy | 12 |
| <i>1.1. Natural basics of Mössbauer spectroscopy</i> | 12 |
| <i>1.2. Doppler shift</i> | 20 |
| <i>1.3. Isomeric shift</i> | 21 |
| <i>1.4. Quadrupole splitting</i> | 22 |
| <i>1.5. Magnetic dipole splitting</i> | 25 |
| Chapter 2. | |
| Spatial distribution of electron defects in the $\text{R}\text{Ba}_2\text{Cu}_3\text{O}_{7-x}$ lattices | 27 |
| <i>2.1. Determination of properties of electric field gradient tensor at $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ lattice sites by emission Mössbauer spectroscopy</i> | 27 |
| <i>2.2. Parameters of the electric field gradient tensor at the copper sites in $\text{YBa}_2\text{Cu}_3\text{O}_{7-y}$. Problems of comparison of calculated and experimental values (EMS with the ^{67}Cu(^{67}Zn) isotope data)</i> | 33 |
| <i>2.3. Charge states of atoms in the superconductors $\text{R}\text{Ba}_2\text{Cu}_3\text{O}_7$</i> | 53 |
| <i>2.4. Electric field gradient tensor in positions of rare-earth metals in $\text{R}\text{Ba}_2\text{Cu}_3\text{O}_7$ lattice</i> | 79 |
| <i>2.5. Combined magnetic and electric quadrupole hyperfine interaction at the copper sites in high-temperature superconductor lattices investigated by ^{61}Cu(^{61}Ni) emission Mössbauer spectroscopy</i> | 89 |
| <i>2.6. Environment of Cu(I) sites in the tetragonal lattices $\text{Y}_{1-x}\text{Ca}_x\text{Ba}_{2-y}\text{La}_y\text{Cu}_3\text{O}_{6+z}$</i> | 95 |
| <i>2.7. Charge distribution in the $\text{Pr}\text{Ba}_2\text{Cu}_3\text{O}_7$ lattice</i> | 103 |
| <i>2.8 Superconductivity and magnetic ordering of the copper sublattice in $\text{YBa}_2\text{Cu}_{3-x}\text{Fe}_x\text{O}_{7+y}$ ceramics</i> | 106 |

Chapter 3.

| | |
|--|-----|
| Spatial distribution of electron defects in the $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$, $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_4$ lattices | 112 |
| 3.1. <i>Comparison of the experimental and calculated values of the parameters of the tensor describing the electric field gradient at iron impurity atoms in cupric oxide</i> | 112 |
| 3.2. <i>Using the $^{57m}\text{Fe}^{3+}$ Mössbauer probe to determine the EFG tensor parameters at CuO cation sites</i> | 122 |
| 3.3. <i>Mössbauer determination of the gradient of the electric-field at the cationic sites of $\text{La}_{2-x}(\text{Sr},\text{Ba})_x\text{CuO}_4$ lattice (EMS with the ^{57}Co(^{57m}Fe), ^{67}Cu(^{67}Zn) u ^{133}Ba(^{133}Cs) isotopes data)</i> | 130 |
| 3.4. <i>^{61}Cu(^{61}Ni) emission Mossbauer study of hyperfine interactions in copper-based oxides</i> | 140 |
| 3.5. <i>Lattice EFG tensors at the rare-earth metal sites in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$</i> | 146 |
| 3.6. <i>Spatial localization of holes in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$</i> | 152 |
| 3.7. <i>Electron density distribution in the superconductor $\text{Nd}_{1.85}\text{Ce}_{0.15}\text{CuO}_4$</i> | 164 |
| 3.8. <i>Hyperfine interactions of copper ions in the structure of high-temperature superconductors</i> | 172 |

Chapter 4.

| | |
|---|-----|
| Spatial distribution of electron defects in the $\text{RBa}_2\text{Cu}_4\text{O}_8$, $\text{Y}_2\text{Ba}_4\text{Cu}_7\text{O}_{15}$, $\text{Tl}_2\text{Ba}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$, $\text{Bi}_2\text{Sr}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$, and $\text{HgBa}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+2}$ lattices | 190 |
| 4.1. <i>Identification of the charge state of atoms in high-T_c superconductor lattices by the comparison of ^{63}Cu NQR/NMR and ^{67}Cu/^{67}Zn emission Mössbauer spectroscopy data</i> | 190 |

| | |
|---|------------|
| 4.2. Atomic charges in $R\text{Ba}_2\text{Cu}_4\text{O}_8$ lattices derived from a comparison of experimental and calculated nuclear-quadrupole coupling parameters..... | 198 |
| 4.3. Spatial distribution of electron defects in the $\text{Y}_2\text{Ba}_4\text{Cu}_7\text{O}_{15}$ lattice | 210 |
| 4.4. Charge states of atoms in the lattices of the high-temperature superconductors $\text{Tl}_2\text{Ba}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$ and $\text{Bi}_2\text{Sr}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$ | 221 |
| 4.5. Mössbauer study of the $\text{HgBa}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+2}$ ceramics..... | 238 |
| 4.6. Charge States of Atoms in Ceramic Superconductors $\text{HgBa}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+2}$, $\text{Tl}_2\text{Ba}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$, and $\text{Bi}_2\text{Sr}_2\text{Ca}_{n-1}\text{Cu}_n\text{O}_{2n+4}$ ($n = 1, 2, \text{ and } 3$)..... | 249 |
| 4.7 Determination of charge states of atoms in the lattices of superconducting copper metal oxide by NMR and emission Mössbauer spectroscopy..... | 265 |
| Chapter 5. | |
| Observation of localization-delocalization of electron pairs in high-temperature superconductors..... | 278 |
| 5.1. Experimental observation of localization-delocalization of Cooper pairs in $\text{Nd}_{1.85}\text{Ce}_{0.15}\text{CuO}_4$ | 278 |
| 5.2. Experimental observation of Bose condensation in high-temperature superconductors..... | 283 |
| 5.3. Spatial distribution of Bose condensate in high-temperature superconductors, determined by emission Mössbauer spectroscopy..... | 293 |
| 5.4. Mössbauer investigation into the influence of disordering of the electronic subsystem on the electron density distribution in copper metal oxides..... | 307 |
| References | 317 |