

B.Glowacka	
Microbiological control of the nun moth (<i>Lymantria monacha</i> L.) caterpillars	75
J.Kudler	
Susceptibility of forest pests to infection by preparations of <i>Bacillus thuringiensis</i> in dependence on the type and the outbreak phase	79
S.Zachov	
Results of many years measures in the biological control done in the reserves of main forest pests in Bulgaria	88
A.Samšišňáková	
Fungi attacking forest pests and their possibility in forest protection	95
V.Mařha - J.Weiser	
Effect of the fungus <i>Beauveria bassiana</i> on the adult of spruce bark beetle <i>Ips typographus</i>	103
Z.Mráček	
Nematoda and other natural enemies in biological control of <i>Cephalcia abietis</i>	107
E.A.Hernandes - Z.Mráček	
Nematoda and their use in the biological control of forest pests	114
J.Křístek - J.Urban	
The use of parasites for the reduction of the population density of willow sawflies <i>Euura laeta</i> (Zadd.) and <i>Euura mucronata</i> (Htg.)	119
J.Hruška - P.Bezděčka	
Protection of ants of the genus <i>Formica</i> and their rational use in forestry	126

SECOND THEMATIC CIRCLE: USE OF SEXUAL PHEROMONES OF LEPIDOPTERA
SPECIES IMPORTANT FROM SILVICULTURAL POINT OF VIEW

R.Hochmut - V.Skuhravý

Present state and prospects of the use of pheromones
for the control and prognosis of defoliators
in the ČSR 135

R.Hochmut - A.Kolk - J.Vrkoč - J.Žďárek

Results of the application of pheromones for the
development of the disorientation effect at various
level of the population density of the nun moth
(*Lymantria monacha*) 142

H.Bogenschütz - R.Albert

Monitoring populations of defoliators in Central
European pine forests with pheromone baited traps 151

W.Baltensweiler

On the problem of host races in the larch bud moth
(*Zeiraphera diniana* Gn., Tortricidae) and its outbreaks
on Norway spruce in Central Europe 162

A.Kolk

Application of pheromones of *Rhyacionia buoliana*
Schiff. in the forests of Poland 167

E.V.Arteaga - R.Hochmut

Possible use of pheromones in forest protection
on Cuba 175

THIRD THEMATIC CIRCLE: USE OF AGGREGATION PHEROMONES FOR THE
CONTROL AND THE REDUCTION OF THE SPRUCE BARK BEETLE POPULATION
(*IPS TYPOGRAPHUS* POPULATION)

M.Švestka - J.Balek

Results and prospects of the research and practical
use of aggregation pheromone of spruce bark beetle
in the ČSR 185

A.Bakke	
Potential use of synthetic pheromone of <i>Ips typographus</i> for monitoring, population suppression and timber protection	191
V.Zumr	
The flight activity and dispersal of the spruce bark beetle, <i>Ips typographus</i> in Norway spruce stands	199
H.H.Eidmann	
Control and monitoring of the spruce bark beetle <i>Ips typographus</i> in Sweden using pheromones	205
V.Novák	
Preliminary results of the research into the reduction of bark beetles <i>Ips typographus</i> L. and <i>Ips amitinus</i> Eichh.	222
G.E.Ozols	
System of the use of pheromone of spruce bark beetle - <i>Ips typographus</i> L.	230
D.Brutovský	
Evaluation of various types of pheromone traps for bark beetles (Coleoptera, Scolytidae)	234
A.Rodziewicz	
Studies on the range of activity of Pheroprax - an aggregation pheromone of the spruce bark beetle (<i>Ips typographus</i> L.)	239
A.Krištín - Š.Varkonda - Ľ.Kuruc - F.Benda	
The new findings in development of new pheromone preparations against the spruce bark beetle <i>Ips typographus</i> L. and notes to their selective effect	242
K.Opalički	
The use of aggregate pheromones for control and reduction of the bark beetle (<i>Ips typographus</i>) in the forests of the Socialist Republic of Croatia	261

E.Kula

Natural enemies of spruce bark beetle (*Ips
typographus* L.) 266

I.Ceianu - V.Mihalciuc - I.Oprean - L.Tautan - L.Ginsca
Aggregation pheromones in the control of the bark
beetles attacking conifers in Roumania 276