

Contents

| | |
|--|----|
| Introduction | 11 |
| Character of empirical types of ore mineralization in the Bohemian Massif | 15 |
| Metamorphic stratiform Cu-Fe-(Pb-Zn) sulfide mineralization in the Monotonous Group of the Moldanubicum (type Bodenmais) | 15 |
| Metamorphic W mineralization of greisen stockworks in orthogneisses and their mantle in the Moldanubicum (type Cetoraz) | 17 |
| Metamorphic Fe deposits of the skarn type, associated with amphibolite and marble, or at the contact of marble with orthogneiss in the Moldanubicum (type Vlastějovice n. Sázavou) | 17 |
| Metamorphic stratiform Pb-Zn sulfide mineralization in the Moldanubicum (type Český Šternberk-Malovidy) | 19 |
| Metamorphic stratiform scheelite mineralization of the Moldanubian erlans (type Deštná) | 20 |
| Metamorphic skarn Fe oxide and Cu sulfide mineralization in various crystalline rock series of the Bohemian Massif outside the region of the Moldanubicum proper (type Měděnec) | 21 |
| Upper Proterozoic partly metamorphic stratiform pyrite mineralization of "black shales" (iron sulfide shales or alum shales) (type Hromnice) | 23 |
| Upper Proterozoic partly metamorphic stratiform Mn carbonate and pyrite mineralization of "black shales" (type Chvaletice) | 26 |
| Upper Proterozoic partly metamorphic stratiform Fe-Cu-Zn-Pb sulfide mineralization (type Újezd) | 28 |
| Upper Proterozoic metamorphic stratiform barite mineralization (type Krhanice) | 30 |
| Upper Proterozoic metamorphic stratiform Fe-V-U mineralization (Fe-U-V-Zn-Cu Mo formation of Mrázek - Pouba 1976) (type Struhadlo) | 30 |
| Upper Proterozoic metamorphic stratiform Fe silicites (type Blovice) | 32 |
| Upper Proterozoic metamorphic stratiform scheelite mineralization in metatuffs (type Čelina - Mokrsko) | 32 |
| Upper Proterozoic metamorphic stratiform Fe-Pb-Zn sulfide mineralization of the Lúgicum (type Hraničná) | 33 |
| Intramagmatic ilmenite-magnetite and sulfide mineralization of basic and ultrabasic intrusive rocks (type Pocinovice) | 34 |
| Upper Proterozoic to early Paleozoic metamorphic stratiform Fe oxide-sulfide and Cu sulfide mineralization of the Varied Group (type Čerčany) | 35 |
| Upper Proterozoic or early Paleozoic metamorphic stratiform banded magnetite mineralization (type Švagrov) | 36 |
| Disseminated and stockwork Mo sulfide mineralization of the Upper Proterozoic (?) (type Všestary) | 36 |
| Disseminated Cu sulfide mineralization of Upper Proterozoic to early Paleozoic keratophyres (Maršovický vrch hill) | 37 |
| Upper Proterozoic (?) stratiform or impregnated Sn oxide mineralization (type Nové Město pod Smrkem) | 37 |

| | |
|---|----|
| Fissure-bound up to impregnated Mo sulfide mineralization of the Upper Proterozoic to Cambrian Brno massif (type Černá Hora) | 38 |
| Cambrian magmatic liquation Ni-Cu sulfide mineralization of basic-ultrabasic rocks (type Staré Ransko) | 39 |
| Ni-Cu sulfide mineralization of altered pre-Devonian diabase dykes (type Sohland) | 41 |
| Cambro-Lower Ordovician metamorphic stratiform Fe oxide mineralization (type Vrát) | 41 |
| Cambro-Lower Ordovician metamorphic stratiform Fe-Cu sulfide mineralization (type Tisová) | 42 |
| Ordovician stratiform Fe-Cu-Zn sulfide mineralization (type Kupferberg) | 44 |
| Ordovician unmetamorphosed stratiform oolitic Fe oxide, carbonate, and silicate mineralization (type Ejpovice) | 45 |
| Impregnated to vein Hg sulfide mineralization in mostly Ordovician (volcanic) sedimentary rocks (type Dědova hora hill) | 47 |
| Silurian or Devonian (?) metamorphic stratiform Fe-Cu sulfide mineralization (type Wleściszowice) | 48 |
| Early Paleozoic stratiform or disseminated pyrite mineralization of porphyroids (type Lukavice) | 49 |
| Early Paleozoic disseminated Fe-Cu-Zn sulfide mineralization in mylonites, or stratiform mineralization of xenoliths (type Včelákov) | 49 |
| Devonian metamorphic stratiform Fe-Cu-Zn-Pb-(Au) sulfide mineralization (type Zlaté Hory) | 50 |
| Devonian metamorphic stratiform Fe-Zn-Pb-(Ag) sulfide mineralization (type Horní Město) | 54 |
| Middle to Upper Devonian stratiform Fe oxide and silicate mineralization of the Lahn-Dill type (type Medlov) | 56 |
| Devonian to Carboniferous vein or strata-bound Fe-Zn sulfide mineralization in the altered spilite and tuff zone (type Hlásnice) | 57 |
| Devonian to Carboniferous metamorphic stockwork or vein Au mineralization, partly overlapping the Devonian metamorphic stratiform Fe-Zn-Pb-Cu sulfide mineralization (type Zlaté Hory-West) | 58 |
| Devonian to Lower Carboniferous (?) stratiform or disseminated Fe-Cu-Mo sulfide mineralization (type Vidly) | 59 |
| Post-Silurian or pre-Upper Permian stratiform Fe oxide skarn mineralization (boreholes at Ryczów) | 60 |
| Post-Silurian or pre-Upper Permian Au-Te mineralization in skarns (boreholes at Ryczów) | 61 |
| Post-Silurian or pre-Upper Permian contact metasomatic or vein Fe-Zn-Cu-Pb-Bi-Ag-Te mineralization (boreholes at Zawiercie) | 62 |
| Devonian to Lower Carboniferous metamorphogenic Au mineralization in the Moldanubicum (type Kašperské Hory) | 62 |
| Devonian or Carboniferous bedded or vein siderite mineralization of Upper Proterozoic carbonate rocks (type Arzberg) | 64 |
| Upper Devonian to Lower Carboniferous Au vein or stockwork mineralization (type Jílové u Prahy) | 64 |
| Lower Carboniferous Au vein and stockwork mineralization of "black shales" (type Suchá Rudná) | 68 |
| Lower Carboniferous (?) metamorphogenic Au stockwork mineralization in pre-Devonian metamorphic rocks (type Zlatý Chlum) | 69 |

| | |
|---|-----|
| Lower Carboniferous (?) disseminated Au and As sulfide mineralization (type Zlötí Stok) | 70 |
| Lower Carboniferous (?) Sb sulfide vein mineralization in the Saxo-Thuringicum (type Schleiz) | 71 |
| Massive partly metamorphic Zn-(Cu) sulfide mineralization in basic intrusives (type Staré Ransko) | 73 |
| Carboniferous metamorphic stratiform barite Fe-Zn-Pb sulfide mineralization of a mylonite zone (type Křižanovice) | 74 |
| Disseminated Mo-Cu sulfide mineralization in Upper Devonian to Lower Carboniferous intrusive rocks (type Dolní Heřmánky) | 75 |
| Au-(Pd, Ag) mineralization of fault zones in metamorphosed Paleozoic rocks (type Svoboda nad Úpou) | 76 |
| Carboniferous (?) polymetallic sulfide vein mineralization, mostly transecting stratiform sulfide deposits (type deposit Bayerland) | 77 |
| Upper Carboniferous or Permian disseminated or stockwork porphyry Cu-Mo sulfide mineralization (type Myszków) | 78 |
| Upper Carboniferous skarn Fe oxide mineralization, in part with Cu sulfides (type Pöhlá) | 78 |
| Upper Carboniferous W-Sn oxide vein mineralization (type Pechtelsgrün) | 79 |
| Upper Carboniferous Mo sulfide greisen mineralization (type Nebelstein Mt.) | 81 |
| Upper Carboniferous erlan (or skarn) scheelite mineralization (type Göpfersgrün) | 81 |
| Upper Carboniferous Mo sulfide vein or stockwork mineralization (type Hůrky near Čistá) | 82 |
| Upper Carboniferous (?) Au vein or stockwork mineralization (type Roudný) | 83 |
| Upper Carboniferous or Permian limnic strata-bound Cu sulfide and carbonate mineralization (type Horní Verneřovice) | 85 |
| Upper Carboniferous to Lower Permian stratiform U mineralization of coal basins (type Jedomělice) | 86 |
| Upper Carboniferous stratiform bauxite mineralization (type Nowa Ruda) | 87 |
| Upper Carboniferous to Permian strata-bound Au mineralization (type Křivce) | 88 |
| Upper Carboniferous or Permian strata-bound up to stockwork skarn Fe-Sn-Pb-Zn-Cu oxide and sulfide mineralization (type Breitenbrunn) | 88 |
| Upper-Carboniferous or Permian pseudo-stratiform Sn oxide mineralization in melanomylonite of the "felsite horizon" (type Grossschirma) | 90 |
| Lower Permian Sn-W oxide stockwork and greisen mineralization (type Cínovec - Altenberg) | 92 |
| Lower Permian vein to vein zone Fe-Zn-Pb-Ag-(Cu) sulfide mineralization (type Kutná Hora and the $kb+eb$ Erzformation of Freiberg) | 96 |
| Lower Permian (?) vein quartz-fluorite mineralization of the Moldanubicum (type Kožlí) | 98 |
| Lower Permian vein, exceptionally metasomatic, Fe sulfide-poor Pb-Zn-Cu-Ag sulfide mineralization (type Ratibořské Hory) | 99 |
| Lower Permian (?) Sb sulfide vein mineralization (type Milešov) | 102 |
| Lower Permian vein, exceptionally stockwork uraninite mineralization (Plíbram - Jáchymov) | 104 |
| Lower Permian post-uranium vein mineralization with selenides, sulfides, arsenides, etc. (type Bukov) | 107 |
| Permian (?) Fe-Mn oxide vein mineralization (type Ilmenau) | 108 |
| Upper Permian fluorite-paradoxite-illite-quartz vein mineralization (type Schönbrunn) | 109 |

| | |
|---|-----|
| Permian or Triassic (?) coffinite-uraninite-brannerite mineralization of tectonic zones associated with silicate metasomatism (type Zadní Chodov - Höhensteinweg) | 111 |
| Lower Permian limnic strata-bound Cu-sulfide mineralization of "black shales" (type Okrzeszyn) | 113 |
| Upper Permian strata-bound Cu sulfide mineralization of "black shales" of marine environment (type Lubin) | 113 |
| Pre-Cretaceous lateritic Fe-Al oxide mineralization (type Lukavice) | 115 |
| Triassic (?) Pb-Zn sulfide vein mineralization (type Stříbro) | 115 |
| Triassic (?) hematite-barite vein mineralization (type Ruhla) | 118 |
| Triassic to Jurassic fluorite-barite vein mineralization (type Harrachov - Halsbrücke - Nabburg) | 119 |
| Triassic to Tertiary stratiform Zn-Pb sulfide mineralization of Middle Triassic dolomites of Upper Silesia (type Olkusz) | 125 |
| Post-Permian (?) Cu sulfide vein mineralization (type Mutěnín) | 127 |
| Upper Cretaceous to Oligocene U mineralization of weathered rocks of the crystalline complex (type Fojtov) | 128 |
| Upper Cretaceous to Oligocene Fe hydroxide-Ni hydrosilicate mineralization of weathered ultrabasic rocks (type Szklary) | 129 |
| Upper Cretaceous to Miocene peneconcordant U-Zr oxide mineralization of basal Upper Cretaceous psammites (type Hamr) | 130 |
| Tertiary strata-bound U mineralization of the Tertiary North Bohemian Basins (type Hroznětín) | 132 |
| Tertiary strata-bound Ti-Al oxide mineralization in Tertiary basins (type Most-Bílina) | 133 |
| Mesozoic or Tertiary vein Ag-As-Bi-Co-Ni mineralization (type Jáchymov - Schneeburg) | 134 |
| Mesozoic or Tertiary vein Ag-Sb-As sulfide mineralization (type Jáchymov - Aue) | 136 |
| Tertiary to subrecent metasomatic and vein fluorite-barite mineralization (type Jílové near Děčín and Teplice) | 137 |
| Tertiary vein Pb-Zn-Cu mineralization (type Roztoky nad Labem) | 138 |
| Tertiary clastic Sn oxide mineralization of lacustrine and fluvial sediments (type Ryžovna near Boží Dar) | 139 |
| Quaternary clastic Sn-W oxide mineralization of fluvial sediments (type river Ohře) | 140 |
| Quaternary clastic Au mineralization of fluvial sediments and of glacial and glaciofluvial sediments (type Zlaté Hory) | 141 |
| Less important ore mineralizations of the Bohemian Massif | 142 |
| References | 149 |
| Index | 173 |