

CONTENTS

| | |
|---|----|
| FOREWORD | |
| CONTENTS | 1 |
| Weiser J.: | 5 |
| Microbial insecticides: New dimensions | |
| Roberts D.W. and St.Leger. R.J.: | 11 |
| Entomopathogenic fungi: Recent basic and applied research | |
| Badowska-Czubik T., Ziemnicka J., and Lipa J.J.: | 31 |
| Healthy rearing of the codling moth (<i>Carpocapsa pomonella</i> L.) for virus production | |
| David L. and Weiser J.: | 36 |
| Ultrastructures of <i>Nosema lymantriae</i> and <i>Thelohania similis</i> , two microsporidian pathogens of the gypsy moth | |
| Deseő K.V., Burchi C., and Lazzari L.: | 40 |
| Application of entomopathogenic nematodes in Italian protected crops | |
| Grubhoffer L., Mařhová A., and Mařha V.: | 45 |
| Lectins of <i>Galleria mellonella</i> larvae and their role in physiological and defence processes | |
| Grubhoffer L., Mařha V., Pultar O., and Weiser J.: | 54 |
| Detection of the granulosis virus of the codling moth by some immunochemical techniques | |
| Jegorov A., Mařha V., Sedmera P., and Roberts D.W.: | 64 |
| Destruxins from <i>Metarhizium anisopliae</i> | |

- Jegorov A., Kadlec Z., Novák J., Mařha V., Sedmera P.,
Tříska J., and Zahradníčková H.: 71
Are the depsipeptides of *Beauveria brongniartii* involved
in the entomopathogenic process ?
(Paper dedicated to Prof. J. Weiser at occasion of his 70th birthday).
- Jizba J., Blumauerová M., Kryštůfek V., Žiřka Z., Weiser J.,
Kandybin N.V., Samoukina I., Kiyko I., and Orekhova N.: 82
Pesticides produced by *Streptomyces globisporus* and *S. griseus*
- Kamionek M. and Bednarek A.: 87
Using entomophilous nematodes in control of tenthredinid
Acantholyda nemoralis
- Keller S.: 91
The use of blastospores of *Beauveria brongniartii* to control
Melolontha melolontha in Switzerland
- Kuzmanova J., Terziiski D., and Karageorgiev St.: 98
Electron-microscopic investigation of histopathological changes
in the midgut epithelium of *Lepinotarsa decemlineata* Say larvae
treated with a formulation containing beta-exotoxin of *Bacillus*
thuringiensis
- Landa Z., Jegorov A., Mařha V., and Novák J.: 110
Light induced production of carotenoids by the entomo-
pathogeneous fungus *Aschersonia aleyrodís*
- Landa Z. and Jiranová R.: 120
Entomopathogenic fungi as an additional selective pest suppressing
agents of greenhouse whitefly populations on greenhouse cucumber

| | |
|--|-----|
| Lipa J.: | 131 |
| Spiroplasmas - a new group of insect pathogens | |
| Mařha V., Jegorov A., and Weiser J.: | 137 |
| The role of secondary metabolites in entomopathogenic processes of fungi | |
| Mráček Z. and Arteaga E.M.: | 143 |
| The incidence of the genus <i>Steinernema</i> (Nematoda : Steinernematidae) in Cuba | |
| Mráček Z. and Bednarek A.: | 147 |
| Lateral fields of infective larvae of the family Steinernematidae (Nematoda) and their possible significance in taxonomy | |
| Mráček Z. and Kahounová L.: | 154 |
| A field evaluation of steinernematid nematodes as biological control agents of insect pests | |
| Mráček Z. and Weyda F.: | 159 |
| The unique observation of the natural penetration of the apterygote host by unknown nematode | |
| Novotný J.: | 164 |
| Microsporidian diseases in the control of forest defoliating moth | |
| Peeters H., Mařha V., and Roberts D.W.: | 169 |
| Enzymes involved in the synthesis of fungal toxins | |
| Prenerová E.: | 183 |
| A method for rapid stimulation of germination of the blastospores in entomopathogenic fungi | |

| | |
|---|-----|
| Rovesti L.: | 186 |
| Response of <i>Steinernema</i> spp. and <i>Heterorhabditis</i> spp. to chemical pesticides | |
| Van der Schaaf D.A., Malais M., and Ravensberg W.J.: | 191 |
| Experiments with <i>Verticillium lecanii</i> in cucumber against whitefly and thrips | |
| Smits P.H.: | 196 |
| Control of grass grubs, <i>Phyllopertha horticola</i> and <i>Amphimallon solstitialis</i> (Coleoptera : Scarabaeidae) with entomopathogenic nematodes | |
| Surli J.A., Jegorov A., Zachardová D., and Landa Z.: | 203 |
| Comparison of various isolates of <i>Hirsutella thompsonii</i> | |
| Vey A.: | 213 |
| Toxins of fungi pathogenic for invertebrates: Chemical composition and biological effects | |
| Videnova E.: | 223 |
| Some aspects of the application of <i>Bacillus thuringiensis</i> against insect pests | |
| Weiser J., Mařha V., and Jegorov A.: | 227 |
| <i>Tolypocladium terricola</i> sp. nov., a new entomopathogenic species of the genus <i>Tolypocladium</i> Gams (Hyphomycetes) | |
| Wegensteiner R.: | 238 |
| The efficacy of <i>Beauveria bassiana</i> , <i>B. brongniartii</i> and <i>B. tenella</i> against <i>Hylobius abietis</i> (Col., Curcul.) | |
| AUTHOR INDEX | 243 |