Contents

| Acknowledgementsi |
|--|
| Introduction1 |
| Report of the International Expert Meeting to review and analyse clinical reports on combination treatment for SARS5 |
| Report 1: Clinical research on treatment of SARS with integrated Traditional Chinese medicine and Western Medicine11 |
| Report 2: Clinical efficacy of the treatment of SARS with integrated Traditional Chinese medicine and Western medicine: an analysis of 524 cases |
| Report 3: Manifestation of symptoms in patients with SARS and analysis of the curative effect of treatment with integrated Traditional Chinese medicine and Western medicine |
| Report 4: Clinical study on 103 inpatients undergoing therapy with integrated Traditional Chinese medicine and Western medicine65 |
| Report 5: Clinical observations of 11 patients with SARS treated with Traditional Chinese medicine |
| Report 6: Effects of applying integrated therapy with Traditional Chinese medicine and Western medicine on liver and kidney functions in patients with SARS |
| Report 7: Clinical research on 63 patients with SARS treated with integrated Traditional Chinese medicine and Western medicine111 |
| Report 8: Influence of integrated therapy with Traditional Chinese medicine and Western medicine on lymphocytes and T-lymphocyte subpopulations of patients with SARS |
| Report 9: Analysis of the clinical curative effects on patients with SARS of treatment with Traditional Chinese medicine and Western medicine 131 |
| Report 10: Evaluation of clinical curative effects of Traditional Chinese medicine in treatment of patients convalescing from SARS143 |
| Report A: A herbal formula for the prevention of transmission of SARS during the SARS epidemic in Hong Kong Special Administrative Region — a prospective cohort study |
| Report B: Effects of Chinese medicine on patients convalescing from SARS in Hong Kong special administrative region — a prospective non-randomized controlled trial |
| Report C: Traditional Chinese medicine in the management of patients with SARS in Hong Kong Special Administrative Region — a case-control study of 24 patients |
| Annex: |