CONTENTS

Preface Introduction

1.	Lea	of Cells Consist of Several Metabolic Compartments		1
	1.1	The Cell Wall Gives the Plant Cell Mechanical Stability		5
		The cell wall consists mainly of carbohydrates and proteins		5
		Plasmodesmata connect neighboring cells		10
	1.2	Vacuoles Have Multiple Functions		12
	1.3	Plastids Have Evolved From Cyanobacteria		13
		Functionally intact cell organelles can be isolated from plant cells		18
	1.4	Mitochondria Originate From Endosymbionts		20
	1.5	Peroxisomes Perform Fatty Acid Oxidation, Respiration, and Reactive		
		Oxygen Species Metabolism		22
	1.6	The Endoplasmic Reticulum and Golgi Apparatus Form a Network for the Distribution of Biosynthesis Products	6	23
	1.7	Various Transport Processes Facilitate the Exchange of Metabolites Between Different Compartments	14	27
		Translocators catalyze the specific transport of metabolic substrates and products		28
		Metabolite transport is achieved by a conformational change to the translocator		30
		Aquaporins make cell membranes not only permeable for water	٤	33
		Ion channels exhibit a very high transport capacity		34
		Porins consist of β-sheet structures		38
2.	Sola	ar Power and Photosynthesis Are the Basis of Life on Earth	4	13
	2.1	The Origin of Photosynthesis	A .	44
				1-

xv xvii

2.2 Pigments Capture Energy From Sunlight 45 The energy content of light depends on its wavelength 45 2.3 Chlorophyll is the main photosynthetic pigment 47 Light Absorption Excites the Chlorophyll Molecule 50 2.4 Antennas Are Required to Capture Light 54 How the excitation energy of the photons is captured in the antennas and transferred to the reaction centers 55 The function of an antenna is illustrated by the antenna of photosystem II 57 Phycobilisomes enable cyanobacteria and red algae to carry out photosynthesis in dim light 59

vi Contents

3.	Pho	otosynthesis Is an Electron Transport Process	63
	3.1	The Photosynthetic Machinery Is Constructed From Complex Modules	63
	3.2	A Reductant and an Oxidant Are Formed During Photosynthesis	66
	3.3	Elucidation of the 3D Structure of a Photosynthetic Reaction Center of	
		Purple Bacteria Was Resolved by X-Ray Crystallography	67
		X-ray structure analysis	69
		The reaction center has a symmetric structure	71
		How does a reaction center function?	72
	3.4	Two Photosynthetic Reaction Centers Are Arranged in Tandem in	
		Photosynthesis of Plants	74
	3.5	Water Is Oxidized by Photosystem II	77
	3.6	Herbicides used in mechanized agriculture target photosystems	83
	3.7	The Cytochrome b ₆ /f Complex Mediates Electron Transport Between	05
		Photosystem II and Photosystem I	65
		from atoms in cytochromes and iron—sulfur centers function as redox carriers	85
		Electron transport by the cytochrome b_6/t complex is coupled to a proton transport	89
		The number of protons pumped through the cyt b_6/f complex can be	02
		doubled by a Q-cycle The Dreduct of Dhotes store Lie Deduced Niestinerside Adenine	95
	3.8	Disucleatide Phosphate (NADPH)	94
		The light operaw driven cyclic electron transport of photosystem produces	24
		only adenosine triphosphate	96
		In the absence of other accentors electrons can be transferred from	20
		photosystem I to oxygen	97
	3.9	Regulatory Processes Control the Distribution of the Captured Photons	
	nsm	Between the Two Photosystems	101
		Excess light energy is eliminated as heat	103
4.	Ade	enosine Triphosphate Is Generated by Photosynthesis	107
	4.1	A Proton Gradient Serves as an Energy-Rich Intermediate State During Adenosine	
		Triphosphate Biosynthesis	107
	4.2	The Electron Chemical Proton Gradient Can Be Dissipated by Uncouplers to Heat	111
	4.3	H ⁺ -ATP Biosynthases From Bacteria, Chloroplasts, and Mitochondria Have a	
		Common Basic Structure	113
		H ⁺ -ATP synthase of chloroplasts is regulated by light	117
		V-ATPase Is related to the F-ATP synthase	117
	4.4	The Biosynthesis of Adenosine Triphosphate Depends on Conformation	
		Change of the Protein	118
5	Mit	ochondria Are the Power Station of the Cell	123
		Biological Oxidation Is Preceded by Degradation of Substrates to Form	
	5.1	protograd and the record of begradener of beneditated to remin	
	5.1	Bound Hydrogen and CO ₂	123

Contents vii

144

145

147

148

149

149

151

157

158

160

167

5.3	Biological Oxidation Takes Place in the Matrix	125
	Pyruvate is oxidized by a multienzyme complex	125
	Acetate is completely oxidized in the tricarboxylic acid cycle	129
	A loss of intermediates of the tricarboxylic acid cycle is replenished by	
	anaplerotic reactions	131
5.4	How Much Energy Can Be Gained by the Oxidation of NADH?	133
5.5	The Mitochondrial Respiratory Chain Shares Common Features With the	
	Photosynthetic Electron Transport Chain	134
	The complexes of the mitochondrial respiratory chain	136
	Electron transport of the respiratory chain Is coupled to the biosynthesis of	
	ATP via proton transport	140
	Mitochondrial proton transport results in the formation of a membrane potential	141
	Mitochondrial ATP biosynthesis serves the energy demand of the cytosol	143

- white children and the biosynthesis serves the childry actually actually of the cytosol
- 5.6 Plant Mitochondria Fulfill Special Metabolic Functions Mitochondria can oxidize surplus NADH without the formation of ATP Cytosolic NADH and NADPH can be oxidized by the respiratory chain
- 5.7 Compartmentation of Mitochondrial Metabolism Requires Specific Membrane Translocators
- 6. The Calvin–Benson–Bassham Cycle Catalyzes Photosynthetic CO₂ Assimilation
 - 6.1 Carboxylation, Reduction, and Acceptor Regeneration Are Three Major Steps in CO₂ Assimilation
 - 6.2 Ribulose Bisphosphate Carboxylase/Oxygenase (Rubisco) Catalyzes Two Reactions Simultaneously

There is no net CO₂ fixation at the compensation point

6.3 The Reduction of 3-Phosphoglycerate Yields Triose Phosphate

6.4 The CO₂ Acceptor Ribulose 1,5-Bisphosphate Is Regenerated From Triose Phosphate6.5 Reductive and Oxidative Pentose Phosphate Pathways Are Present in Chloroplasts

6.6 Reduced Thioredoxins Transmit the Signal "Illumination" and Activate or Inactivate

		Enzymes	170
		Multiple regulatory processes tune the reactions of the reductive pentose phosphate pathway	174
7.	Pho	otorespiratory Pathway Recycles Phosphoglycolate	177
	7.1	Ribulose 1,5-Bisphosphate Is Recovered by Recycling 2-Phosphoglycolate	177
	7.2	Peroxisomes Need External Reducing Equivalents for the Reduction of Hydroxypyruvate	183
	7.3	Reassimilation of NH ₄ ⁺ Released in the Photorespiratory Pathway in Chloroplasts	186
	7.4	The Peroxisomal Matrix Eliminates Toxic Metabolites	188
8.	Pho	otosynthesis Needs the Consumption of Water	191
	8.1	The Uptake of CO ₂ Into the Leaf Is Accompanied by an Escape of Water Vapor	191

8.2	Stomata Regulate the Gas Exchange of Leaves	192
8.3	The Diffusive Flux of CO ₂ Into a Plant Cell	197
8.4	C4 Plants Perform CO2 Assimilation With Lower Water Consumption Than C3 Plants	199
	The CO ₂ pump in C ₄ plants	201
	C4 metabolism in plants of the NADP malic enzyme type	204
	C4 metabolism in plants of the NAD malic enzyme type	206
	C4 metabolism in plants of the phosphoenolpyruvate carboxykinase type	209
	Enzymes of C ₄ metabolism are regulated by light	209
	C4 plants have advantages in climates with higher temperatures	211
8.5	Crassulacean Acid Metabolism Allows Plants to Survive Severe Water Shortages	211
	CO ₂ fixed during the night is stored as malic acid	213
	Photosynthesis proceeds with closed stomata	214

9.	Pol	ysaccharides Are Storage and Transport Forms of Carbohydrates duced by Photosynthesis	217
	9.1	Sucrose Biosynthesis Takes Place in the Cytosol	219
	92	The Utilization of the Photosynthesis Product Triose Phosphate Is Strictly Regulated	221
		Fructose 1,6-bisphosphatase is an entrance valve for the sucrose biosynthesis pathway	221
		covalent modification	225
	9.3	Large Quantities of Carbohydrate Can Be Stored as Starch to Support Metabolism and Growth at Night	227
		Starch is synthesized via ADP-glucose	230
		Photosynthesis products can be stored temporarily in chloroplasts as starch	231
		Starch accumulation is regulated to provide carbohydrate for the plant at night	234
	9.4	Degradation of starch proceeds in two different ways	235
		Assimilatory starch is degraded by a combination of hydrolytic degradation	
		in the plastid, maltose export, and phosphorolysis in the cytosol	239
		Starch degradation is regulated by the circadian clock to ensure that plants do not starve at night	241
		Trehalose 6-phosphate is an important signal mediating the carbohydrate supply	242
	9.5	In Some Plants Photoassimilates From the Leaves Are Exported as Sugar Alcohols or Oligosaccharides of the Raffinose Family	244
	9.6	Fructans Are Deposited as Storage Compounds in the Vacuole	247
	9.7	Cellulose Is Synthesized by Enzymes Located in the Plasma Membrane	250
		Most cell wall polysaccharides are synthesized in the Golgi apparatus	252
		Biosynthesis of callose is often induced by wounding	253
10.	Niti	rate Assimilation Is Essential for the Biosynthesis of Organic Matter	255
	10.1	The Reduction of Nitrate to NH ₃ Proceeds in Two Reactions	255
		Nitrate is reduced to nitrite in the cytosol	257
		The reduction of nitrite to ammonia proceeds in plastids	259

100

Contents ix

	The assimilation of NH ₄ ⁺ proceeds via two reactions of GS/GOGAT	261
	The oxidative pentose phosphate pathway in leucoplasts provides reducing	
	equivalents for nitrite reduction	262
0.2	Nitrate Assimilation Is Controlled at Various Levels	263
0.3	The End Product of Nitrate Assimilation Is a Whole Spectrum of Amino Acids	266
	CO2 assimilation provides the carbon skeletons for the biosynthesis of amino acids	267
	The biosynthesis of glutamate requires the participation of mitochondrial metabolism	269
	Biosynthesis of proline and arginine	269
	Aspartate is the precursor of five amino acids	271
	Acetolactate synthase participates in the biosynthesis of hydrophobic amino acids	274
	Aromatic amino acids are synthesized via the shikimate pathway	277
	Glyphosate acts as a herbicide in the shikimate pathway	279

1

11.	Nitro	ogen Fixation Enables Plants to Use the Nitrogen in the	
	Air f	or Growth	285
	11.1	Legumes Form a Symbiosis With Nodule-Inducing Bacteria	286
		The nodule formation depends on balanced bacterial and plant gene expression	288
		Metabolic products are exchanged between bacteroids and host cells	290
	11.2	Dinitrogenase Reductase Delivers Electrons for the Dinitrogenase Reaction	291
		N_2 as well as H^+ are reduced by dinitrogenase	292
		N ₂ fixation can proceed only at very low oxygen concentrations	294
		The energy costs for utilizing N_2 as a nitrogen source are much higher than for NO_3^-	296
	11.3	Plants Improve Their Nutrition by Symbiosis With Fungi	296
		The arbuscular mycorrhiza is widespread	297
		Ectomycorrhiza supply trees with nutrients	298
		A cetyl-Coal carboxydase is the the the Werkerie of the bissivit one stored by bissivit one stored	
12.	Proc	lucts of Nitrogen Fixation and Nitrate Assimilation Are Deposited	
	as St	torage Proteins	299
	12.1	Globulins Are the Most Abundant Storage Proteins	300
	12.2	2S-Proteins Are Present in Seeds of Oilseed Plants	301
	12.3	Prolamins Are Formed as Storage Proteins in Grasses	301
	12.4	Special Proteins Protect Seeds From Being Eaten by Animals	301
	12.5	Biosynthesis of Storage Proteins Is Performed at the Rough Endoplasmic Reticulum	302
	12.6	Proteinases Mobilize the Amino Acids Deposited in Storage Proteins	305
13.	Sulfa	ate Assimilation Enables the Synthesis of Sulfur-Containing	
	Com	pounds	307
	13.1	Sulfate Assimilation Proceeds Primarily by Photosynthesis	307
		Sulfate is activated prior to reduction	309
		Sulfite reductase is similar to nitrite reductase	311
		H ₂ S is fixed in the amino acid cysteine	312

280

x Contents

	13.2	Glutathione Serves the Cell as an Antioxidant and Is an Agent for the Detoxification of Pollutants	314
		Xenobiotics are detoxified by conjugation	315
		Phytochelatins protect the plant against heavy metals	316
		Excessive concentrations of sulfur dioxide in the air are toxic for plants	318
	13.3	Methionine Is Synthesized From Cysteine	319
		S-Adenosylmethionine is a universal methylation reagent	319
	13.4	Sulfate Plays an Important Role During Drought Stress	319
14.	Phlo	em Transport Distributes Photoassimilates to Various Sites	
	of C	onsumption and Storage	323
	14.1	There Are Different Modes of Phloem Loading	325
	14.2	Phloem Transport Proceeds by Mass Flow	327
	14.3	Sink Tissues Are Supplied by Phloem Unloading	327
		Starch is deposited in plastids	328
		The glycolysis pathway utilizes carbohydrates	330
15.	Lipic	Is Are Membrane Constituents and Function as Carbon Stores	335
	15.1	Polar Lipids Are Important Membrane Constituents	335
		The fluidity of the membrane is governed by the proportion of unsaturated	
		fatty acids and the content of sterols	338
		Membrane lipids contain a variety of hydrophilic head groups	339
		Sphingolipids are important constituents of the plasma membrane	342
	15.2	Triacylglycerols Are Storage Compounds: their de novo biosynthesis takes place in plastids	342
		Acetyl-CoA is a precursor for the biosynthesis of fatty acids	344
		Acetyl-CoA carboxylase is the first enzyme of fatty acid biosynthesis	345
		Fatty acid biosynthesis is catalyzed by a multienzyme complex	347
		The first double bond in a newly synthesized fatty acid is introduced by a	
		soluble acyl-ACP-dependent desaturase	350
	15.3	Glycerol 3-Phosphate Is a Precursor for the Biosynthesis of Glycerolipids	353
		Triacylglycerol biosynthesis	355
		The endoplasmic reticulum membrane is the site of fatty acid elongation and desaturation	357
		Some plactidial membrane lipide are synthesized via the eukanyotic pathway	357
	15 /	During Seed Cormination Storage Lipids Are Mobilized in the Glyovysomes for the	557
	13.4	Production of Carbohydrates	360
		The glyoxylate cycle enables plants to synthesize hexoses from acetyl-CoA	362
		Reactions with toxic fatty acid intermediates take place in peroxisomes	364
	15.5	Specialized Fatty Acids and Derivatives	365
		Plant fat is used for human nutrition and also as a raw material in industry	368



16.	Spec	ial Metabolites Fulfill Specific Biological and Ecological	
	Fund	tions in Plants	373
	16.1	Specialized Metabolites Often Protect Plants From Pathogenic Microorganisms	
		and Herbivores	373
		Plants protect themselves by tricking herbivores with false amino acids	374
		Pathogens produce aggressive compounds to ensure their survival	375
	16.2	Plants React to Pathogen Attack by the Release of Immunomodulatory Agents, Phytoalexins, and Phytoanticipins	376
	16.3	Degradation of Cyanogenic Glycosides Releases Toxic Hydrogen Cyanide (HCN)	380
	16.4	Glucosinolate Degradation Releases Toxic Volatile Mustard Oils	383
	16.5	Alkaloids Comprise a Variety of Heterocyclic Specialized Metabolites	386
	16.6	Transport, Storage, and Turnover of Specialized Metabolites	388
	Pla	ant defense compounds can also be a risk to humans	389
		Auten diartes shoot elenative grander and the left of the state and the sets and the	
17.	A La	rge Diversity of Isoprenoids Has Multiple Functions in Plants	391
	17.1	Higher Plants Have Two Different Biosynthesis Pathways for Isoprenoids	393
		Acetyl CoA is a precursor for the biosynthesis of isoprenoids in the cytosol	393
		Pyruvate and p-glyceraldehyde-3-phosphate are the precursors for the	
		biosynthesis of isopentyl pyrophosphate in plastids	395
		Prenyl transferases catalyze the fusion of isoprene units	396
		Some plants emit isoprenes into the air	398
	17.2	Monoterpenes Derive From Geranyl Pyrophosphate	399
	17.3	Farnesyl Pyrophosphate Is the Precursor for the Biosynthesis of Sesquiterpenes	400
		Steroids are synthesized from farnesyl pyrophosphate	402
	17.4	Geranylgeranyl Pyrophosphate Is the Precursor for Diterpenes and Polyterpenes	404
		Carotene biosynthesis	405
		A prenyl chain renders compounds lipid soluble	407
	17.5	The Regulation of Isoprenoid Biosynthesis	409
		Isoprenoids are very stable and persistent compounds	410

18.	Pher Meta	nylpropanoids Comprise a Multitude of Plant-Specialized abolites and Cell Wall Components	411
	18.1	Phenylalanine Ammonia Lyase and Monooxygenases Are Involved in the Biosynthesis of Phenolic Compounds	411
	18.2	Phenylpropanoid Compounds Polymerize to Macromolecules	417
		Lignans act as defense substances	417
		Lignin is formed by radical polymerization of phenylpropanoid derivatives	419
		Suberins and cutins form gas- and water-impermeable layers of cells	421
	18.3	The Biosynthesis of Flavonoids and Stilbenes Also Requires Acetate for	
		Aromatic Ring Formation	422
		Some stilbenes are very potent natural fungicides	423
		Flavonoids have multiple functions in plants	423

20

Contents

xiii

	20.4	The Mitochondrial Genome of Plants Varies Greatly in Size	494
		in hybrid breeding	497
21.	Bios	ynthesis, Processing, and Degradation of Plant Proteins	503
	21.1	Protein Biosynthesis Is Catalyzed by Ribosomes	503
		A peptide chain is synthesized	507
		Specific inhibitors of the translation can be used to decide whether a protein	
		is synthesized in the cytosol, plastids, or mitochondria	510
		The translation is regulated	512
	21.2	Proteins Attain Their Three-Dimensional Structure by Controlled Folding	513
		The folding of a protein is a multistep process	513
		Heat shock proteins protect against heat damage	516
		Molecular chaperones bind to unfolded proteins	516
	21.3	Nuclear-Encoded Proteins Are Targeted to Various Cell Compartments	519
		Most of the proteins imported into the mitochondria must cross two membranes	520
		The import of proteins into chloroplasts requires several translocation complexes	523
		Protein import into peroxisomes	525
		Nuclear localization signals direct protein import into the nucleus	527
	21.4	Proteins Are Degraded by Proteasomes in a Strictly Controlled Manner	528
22.	Biote	echnology Alters Plants to Meet the Requirements of Agriculture,	
22.	Biote	echnology Alters Plants to Meet the Requirements of Agriculture, ition, and Industry	533
22.	Biote Nutr 22.1	echnology Alters Plants to Meet the Requirements of Agriculture, ition, and Industry A Gene Is Isolated and Archived in a Gene Library	533 533
22.	Biote Nutr 22.1	echnology Alters Plants to Meet the Requirements of Agriculture, ition, and Industry A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies,	533 533
22.	Biote Nutr 22.1	echnology Alters Plants to Meet the Requirements of Agriculture, ition, and Industry A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA	533 533
22.	Biote Nutr 22.1	 A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned 	533 533 537 542
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding	533 533 537 542 542
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length	533 533 537 542 542
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism	533 533 537 542 542
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism	533 533 542 542 544
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism	533 533 537 542 542 544 547
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker	533 533 537 542 542 542 542
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene IsIsolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker Agrobacteria Can Transform Plant Cells	533 533 537 542 542 542 542 549 549
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker Agrobacteria Can Transform Plant Cells The Ti-plasmid contains the genetic information for tumor formation	533 533 537 542 542 542 549 549 549 549
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker Agrobacteria Can Transform Plant Cells The Ti-plasmid contains the genetic information for tumor formation Ti-plasmids are used as transformation vectors	533 533 537 542 542 542 549 549 549 549 549 549
22.	Biote Nutr 22.1 22.3	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker Agrobacteria Can Transform Plant Cells The Ti-plasmid contains the genetic information for tumor formation Ti-plasmids are used as transformation vectors Plant cells are transformed by the uptake of DNA	533 533 537 542 542 542 543 549 549 549 549 549 549
22.	Biote Nutr 22.1 22.3	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker Agrobacteria Can Transform Plant Cells The Ti-plasmid contains the genetic information for tumor formation Ti-plasmids are used as transformation vectors Plant cells are transformed by the uptake of DNA A new plant is regenerated from callus tissue	533 533 537 542 542 542 543 544 549 549 549 551 553 553
22.	Biote Nutr 22.1 22.2	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker Agrobacteria Can Transform Plant Cells The Ti-plasmid contains the genetic information for tumor formation Ti-plasmids are used as transformation vectors Plant cells are transformed by the uptake of DNA A new plant is regenerated from callus tissue Selected promoters and reporter genes enable defined expression studies of foreign capes	533 533 537 542 542 544 549 549 549 551 553 554 555
22.	Biote Nutr 22.1 22.2 22.3	A Gene Is Isolated and Archived in a Gene Library A gene library is screened for a certain gene with DNA probes, antibodies, transposons, or T-DNA Genes encoding unknown proteins can be functionally assigned DNA Polymorphism Yields Genetic Markers for Plant Breeding Individuals of the same species can be differentiated by restriction fragment length polymorphism The random amplified polymorphic DNA technique is a simple method for investigating DNA polymorphism The polymorphism of microsatellite DNA is used as a genetic marker Agrobacteria Can Transform Plant Cells The Ti-plasmid contains the genetic information for tumor formation Ti-plasmids are used as transformation vectors Plant cells are transformed by the uptake of DNA A new plant is regenerated from callus tissue Selected promoters and reporter genes enable defined expression studies of foreign genes	533 533 537 542 542 542 549 549 549 549 549 551 553 554 553

xiv Contents

22.5	Plant Genetic Engineering Can Be Used for Many Different Purposes	564
	Plant protection via Bt protein	565
	Plant protection against viruses	566
	Fungus-resistant plants	567
	Herbicide-resistant plants	567
	Improvement of yield and quality of crop products	568
	Renewable resources for industry	568
	Protection of crop plants against environmental stress	569



571

Moleculer charactores bind to unfolde effective trachter that the solution and the solution of the induct of proteins are carbined to be active trackateries instantion and the induct of proteins are carbined to be inducted proteins and the frequence of the induct of proteins and the frequence of the induct of proteins and the frequence of the induct of proteins and the induct of proteins and the induct of proteins and the induct of proteins are carbined to be inducted of the induct of proteins and the induct of the induct o

